

APPENDIX C



US006824777B1

(12) **United States Patent**
Alitalo et al.

(10) **Patent No.:** **US 6,824,777 B1**
(45) Date of Patent: **Nov. 30, 2004**

(54) **FLT4 (VEGFR-3) AS A TARGET FOR TUMOR IMAGING AND ANTI-TUMOR THERAPY**

(75) Inventors: **Kari Alitalo**, Espoo (FI); **Arja Kaipainen**, Boston, MA (US); **Reija Valtola**, Helsinki (FI); **Lotta Jussila**, Helsinki (FI)

(73) Assignees: **Licentia Ltd.**, Helsinki (FI); **Ludwig Institute for Cancer Research**, New York, NY (US)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

(21) Appl. No.: **09/169,079**

(22) Filed: **Oct. 9, 1998**

Related U.S. Application Data

(63) Continuation of application No. 08/901,710, filed on Jul. 28, 1997, now Pat. No. 6,107,046, and a continuation of application No. 08/340,011, filed on Nov. 14, 1994, now Pat. No. 5,776,755, and a continuation of application No. 08/257,754, filed on Jun. 9, 1994, now abandoned, which is a continuation-in-part of application No. 07/959,951, filed on Oct. 9, 1992, now abandoned.

(51) **Int. Cl.**⁷ **A61K 39/395**

(52) **U.S. Cl.** **424/143.1; 424/133.1; 424/141.1; 514/2**

(58) **Field of Search** **424/133.1, 141.1, 424/143.1; 514/2**

(56) **References Cited**

U.S. PATENT DOCUMENTS

4,543,439 A	9/1985	Fracketton, Jr. et al.
4,652,639 A	3/1987	Stabinsky 536/27
4,933,294 A	6/1990	Waterfield et al. 436/501
5,183,884 A	2/1993	Kraus et al. 536/23.5
5,185,438 A	2/1993	Lemischka 536/23.2
5,198,359 A	3/1993	Taniguchi et al. 435/252.3
5,231,001 A	7/1993	Kaplan et al. 435/7.21
5,256,766 A	10/1993	Coughlin 530/327
5,270,458 A	12/1993	Lemischka 536/23.5
5,283,354 A	2/1994	Lemischka 536/23.5
5,367,057 A	11/1994	Lemischka 530/350
5,635,177 A	6/1997	Bennett et al. 424/143.1
5,643,759 A	7/1997	Pfreundschuh 435/70.21
5,693,762 A	12/1997	Queen et al. 530/387.3
5,700,822 A	12/1997	Hirth et al. 514/380
5,712,395 A	1/1998	App et al. 544/344
5,747,651 A	5/1998	Lemischka 530/387.9
5,750,078 A	5/1998	Shitara et al. 424/133.1
5,763,441 A	6/1998	App et al. 514/249
5,763,733 A	6/1998	Whitlow et al. 530/387.3
5,776,427 A	7/1998	Thorpe et al. 424/1.49
5,776,755 A	7/1998	Alitalo et al. 435/194
5,798,097 A	8/1998	McKenzie et al. 424/181.1
5,807,548 A	9/1998	Shitara et al. 424/133.1
5,952,199 A	9/1999	Davis-Smyth et al. 435/69.7
6,011,003 A	1/2000	Charnock-Jones et al. 514/2
6,331,302 B1	12/2001	Bennett et al. 424/146.1
6,451,764 B1	9/2002	Lee et al. 514/12

FOREIGN PATENT DOCUMENTS

EP	0 325 224 A2	7/1989
WO	WO 90/14425	11/1990
WO	WO 92/14748	3/1992
WO	WO 92/13867	8/1992
WO	WO 93/14124	7/1993
WO	WO 93/15201	8/1993
WO	WO 94/10202	5/1994
WO	WO 95/24473	9/1995
WO	WO 95/33772	12/1995
WO	WO 96/39515	12/1996
WO	WO 97/05250	2/1997
WO	WO 97/09427	3/1997
WO	98/07832	2/1998
WO	WO 98/33917	8/1998
WO	99/33485	7/1999

OTHER PUBLICATIONS

Mikayama T. Molecular cloning and functional expression of a cDNA encoding glycosylation-inhibiting factor. *Proc. Natl. Acad. Sci. USA* vol. 90, pp. 10056-10060, 1993.*

Voet et al. *Biochemistry*. 1990. John Wiley & Sons, Inc., pp. 126-128 and 228-234.*

Beers and Berkow eds. The Merck Manual of Diagnosis and Therapy. Seventeenth Edition, 1999. pp. 986-99 Co., Merck & Co., Inc. Whitehouse Station, N.J.*

Achen, M.G. et al., "Monoclonal antibodies to vascular endothelial growth factor-D block its interactions with both VEGF receptor-2 and VEGF receptor-3," *Eur. J. Biochem*, 267(9):2505-15 (May, 2000).

Akagi, K., et al., "Vascular endothelial growth factor-C (VEGF-C) expression in colorectal cancer tissues," *Br. J. Cancer*, 83(7):887-91 (Oct., 2000).

Andre, T., et al., "Vegf, Vegf-B, Vegf-C and their receptors KDR, FLT-1 and FL the neoplastic progression of human colonic mucosa," *Int. J. Cancer*, 86(2):174-81 (Apr. 15, 2000).

Bunone, G., et al., "Expression of Angiogenesis Stimulators and Inhibitors in Human Thyroid Tumors and Correlation with Clinical Pathological Features," *Am. J. Pathol*, 155(6):1967-76 (Dec. 1999).

Cao, Y., et al., "Vascular endothelial growth factor C induces angiogenesis in vivo," *Proc. Natl. Acad. Sci. USA*, 95:14389-94 (Nov. 1998).

(List continued on next page.)

Primary Examiner—Yvonne Eyer

Assistant Examiner—Joseph F. Murphy

(74) *Attorney, Agent, or Firm*—Marshall, Gerstein & Borun LLP

(57) **ABSTRACT**

The present invention provide purified Flt4 receptor tyrosine kinase polypeptides and fragments thereof, polynucleotides encoding such polypeptides, antibodies that specifically bind such polypeptides, and uses therefor.

54 Claims, 11 Drawing Sheets

(2 of 11 Drawing Sheet(s) Filed in Color)

OTHER PUBLICATIONS

- Eggert, A., et al., "High-Level Expression of Angiogenic Factors Is Associated with Advanced Tumor Stage in Human Neuroblastomas," *Clin. Cancer Res.*, 6(5):1900-8 (May 2000).
- Fellmer, P.T., et al., "Vascular endothelial growth factor-C gene expression in papillary and follicular thyroid carcinomas," *Surgery*, 126(6):1056-61 (Dec. 1999).
- Fielder W., et al., "Expression of FLT4 and its ligand VEGF-C in Acute Myeloid Leukemia," *Leukemia*, 8:1234-7 (Aug. 1997) (Abstract).
- Folpe, A.L., et al., "Vascular Endothelial Growth Factor Receptor-3 (VEGFR-3): A Marker of Vascular Tumors with Presumed Lymphatic Differentiation, Including Kaposi's Sarcoma, Kaposiform and Dabska-Type Hemangioendotheliomas, and a Subset of Angiosarcomas," *Mod. Pathol.*, 13(2):180-185 (2000).
- Gunningham, S.P., et al., "The Short Form of the Alternatively Spliced flt-4 but not Its Ligand Vascular Endothelial Growth Factor C Is Related to Lymph Node Metastasis in Human Breast Cancers," *Clin. Cancer Res.*, 6(11):4278-86 (Nov. 2000).
- Hatva, E., et al., "Vascular Growth Factors and Receptors in Capillary Hemangioblastomas and Hemangiopericytomas," *Am. J. Pathol.*, 148(3):763-75 (Mar. 1996).
- Hu, Jing-Shan, et al., "A Novel Regulatory Function of Proteolytically Cleaved VEGF-2 for Vascular Endothelial and Smooth Muscle Cells," *The FASEB Journal*, 11:498-504 (May 1997).
- Jacquemier, J., et al., "Prognosis of Breast-Carcinoma Lymphagenesis Evaluated by Immunohistochemical Investigation of Vascular-Endothelial-Growth-Factor Receptor 3," *Int. J. Cancer (Pred. Oncol.)*, 89:69-73 (2000).
- Jones, A., et al., "Angiogenesis and lymphangiogenesis in stage 1 germ cell tumours of the testis," *BJU International*, 86:80-86 (2000).
- Karkkainen, M.J., et al., "Vascular endothelial growth factor receptors in the regulation of angiogenesis and lymphangiogenesis," *Oncogene*, 19(49):5598-605 (Nov. 20, 2000).
- Kubo, H., et al., "Involvement of vascular endothelial growth factor receptor-3 in maintenance of integrity of endothelial cell lining during tumor angiogenesis," *Blood*, 96(2):546-553 (Jul. 15, 2000).
- Kurebayashi, J., et al., "Expression of Vascular Endothelial Growth Factor (VEGF) Family Members in Breast Cancer," *Jpn. J. Cancer Res.*, 90(9):977-81 (Sep. 1999).
- Marchio, S., et al., "Vascular Endothelial Growth Factor-C Stimulates the Migration and Proliferation of Kaposi's Sarcoma Cells," *J. Biol. Chem.*, 274(39):27627-22 (Sep. 24, 1999).
- Marconcini, L., et al., "c-fos-induced growth factor/vascular endothelial growth factor D induces angiogenesis in vivo and in vitro," *Proc. Natl. Acad. Sci. USA*, 96:9671-76 (Aug. 1999).
- Moriyama, M., et al., "Immunohistochemical Study of Tumour Angiogenesis in Oral Squamous Cell Carcinoma," *Oral Oncol.*, 33(5):369-74 (Sep. 1997).
- Nathanson, S.D., et al., "Microvessels That Predict Axillary Lymph Node Metastases in Patients With Breast Cancer," *Arch Surg*, 135(5):586-93 (May 2000).
- Nicosia, R.F., "What Is the Role of Vascular Endothelial Growth Factor-Related Molecules in Tumor Angiogenesis?" *Am. J. Pathol.*, 153(1):11-6 (Jul. 1998).
- Niki, T., et al., "Expression of Vascular Endothelial Growth Factors A, B, C, and D and Their Relationships to Lymph Node Status in Lung Adenocarcinoma," *Clin. Cancer Res.*, 6(6):2431-9 (Jun. 2000).
- Ohta, Y., et al., "Increased Vascular Endothelial Growth Factor and Vascular Endothelial Growth Factor-C and Decreased NM23 Expression Associated with Microdissemination in the Lymph Nodes in Stage 1 Non-Small Cell Lung Cancer," *J. Thorac Cardiovasc Surg*, 119(4 Pt 1):804-13 (Apr. 2000).
- Ohta, Y., et al., "VEGF and VEGF type C play an important role in angiogenesis and lymphangiogenesis in human malignant mesothelioma tumours," *Br. J. Cancer*, 81(1):54-61 (Sep. 1999).
- Partanen, T.A., et al., "Lack of Lymphatic Vascular Specificity of Vascular Endothelial Growth Factor Receptor 3 in 185 Vascular Tumors," *Cancer*, 86(11):406-12 (Dec. 1, 1999).
- Saaristo, A., et al., "Vascular Endothelial Growth Factor-C and its Receptor VEGFR-3 in the Nasal Mucosa and in Nasopharyngeal Tumors," *Am. J. Pathology*, 157(1):7-14 (Jul. 2000).
- Shushanov, S., et al., "VEGFc and VEGFR3 Expression in Human Thyroid Pathologies," *Int. J. Cancer*, 86:47-52 (2000).
- Skobe, M., et al., "Vascular Endothelial Growth Factor-C (VEGF-C) and its Receptors KDR and flt-4 are Expressed in AIDS-Associated Kaposi's Sarcoma," *J. Invest. Dermatology*, 113:1047-1053 (1999).
- Sowter, H.M., et al., "Expression and Localization of the Vascular Endothelial Growth Factor Family in Ovarian Epithelial Tumors," *Lab. Invest.*, 77(6):607-14 (Dec. 1997).
- Tsurasaki, T., et al., "Vascular endothelial growth factor-C expression in human prostatic carcinoma and its relationship to lymph node metastasis," *Br. J. Cancer*, 80(2):309-313 (1999).
- Valtola, R., et al., "VEGFR-3 and Its Ligand VEGF-C Are Associated with Angiogenesis in Breast Cancer," *Am. J. Pathol.*, 154(5):1381-90 (May 1999).
- Weninger, W., et al., "Expression of Vascular Endothelial Growth Factor Receptor-3 and Podoplanin Suggests a Lymphatic Endothelial Cell Origin of Kaposi's Sarcoma Tumor Cells," *Lab. Invest.*, 79(2):243-51 (Feb. 1999).
- Witzenbichler, B., et al., "Vascular Endothelial Growth Factor-C (VEGF-C/VEGF-2) Promotes Angiogenesis in the Setting of Tissue Ischemia," *American Journal of Pathology*, vol. 153, No. 2:381-394 (Aug. 1998).
- Yokoyama, Y., et al., "Prognostic Significance of Vascular Endothelial Growth Factor and Its Receptors in Endometrial Carcinoma," *Gyn. Oncology* 77:413-418 (2000).
- Yonemura, Y., et al., "Role of Vascular Endothelial Growth Factor C Expression in the Development of Lymph Node Metastasis in Gastric Cancer," *Clin. Cancer Res.*, 5:1823-1829 (Jul. 1999).
- Achen, M.G. et al., "Vascular endothelial growth factor D (VEGF-F) is a ligand for the tyrosine kinases VEGF receptor 2 (Flk1) and VEGF receptor 3 (Flt4)," *Proc. Natl. Acad. Sci., USA*, 95(2):548-553 (Jan., 1998).
- Andersson et al., "Structural and Functional Markers During Induced Differentiation in Human Leukemia Cell Lines," In R. F. Revoltella (ed.), *Expression of Differentiated Functions in Cancer Cells*. 239-245, Raven Press, New York (1982).

- Aprelikova et al., "FLT4, A Novel Class III Receptor Tyrosine Kinase in Chromosome 5q33-qter," *Cancer Research*, 52(3):746-748 (Feb. 1, 1992).
- Aujame, L. et al., "High affinity human antibodies by phage display," *Human Antibodies*, 8(4):155-168 (1997).
- Beckstead, J.H. et al., "Evidence for the Origin of Kaposi's Sarcoma From Lymphatic Endothelium," *Am. J. Pathol.*, 119(2):294-300 (May, 1985).
- Berridge et al., "Cell-Lineage Antigens of the Stem Cell-Megakaryocyte-Platelet Linkage are Associated with the Platelet IIb-IIIa Glycoprotein Complex," *Blood*, 66(1):76-85 (Jul., 1985).
- Bolen, J.B., "Nonreceptor Tyrosine Protein Kinases," *Oncogene*, 8:2025-2031 (1993).
- Bolhuis, R.L. et al., "T cell targeting in cancer therapy," *Cancer Immunology Immunotherapy*, 34(1):1-8 (1991).
- Borg et al., "Biochemical Characterization of Two Isoforms of FLT4, and VEGF Receptor-Related Tyrosin Kinase," *Oncogene*, 10:973-984 (1995).
- Brown, L.F. et al., "Expression of Vascular Permeability Factor (Vascular Endothelial Growth Factor) and Its Receptors in Breast Cancer," *Human Pathology*, 26(1):86-91 (Jan., 1995).
- Brüggemann, M. et al., "Production of human antibody repertoires in transgenic mice," *Curr. Opin. Biotechnol.*, 8:455-458 (1997).
- Brüggemann, M. et al., "Strategies for expressing human antibody repertoires in transgenic mice," *Immunol. Today*, 17(8):391-397 (Aug., 1996).
- Cantley et al., "Oncogenes and Signal Transduction," *Cell*, 64:281-302 (Jan. 25, 1991).
- Carter, P. et al., "Toward the Production of Bispecific Antibody Fragments for Clinical Applications," *Journal of Hematotherapy*, 4:463-470 (1995).
- Catoretti et al., "Monoclonal Antibodies Against Recombinant Parts of the Ki-67 Antigen (MIB 1 and MIB 3) Detect Proliferating Cells in Microwave-Processed FormalinFixed Paraffin Section," *J. of Pathol.*, 168:357-363 (1992).
- Cheng & Flanagan, "Identification and Cloning of ELF-1, a Developmentally Expressed Ligand for the Mek4 and Sek Receptor Tyrosine Kinases," *Cell*, 79:157-168 (Oct. 7, 1994).
- Cole et al., "The EVB-Hybridoma Technique and Its Application to Human Lung Cancer," *Monoclonal Antibodies and Cancer Therapy*, Alan R Liss, Inc., pp. 77-96 (1985).
- Collins et al., "Continuous Growth and Differentiation of Human Myeloid Leukaemic Cells in Suspension Culture," *Nature*, 270:347-349 (1977).
- De Gast, G.C. et al., "Clinical perspectives of bispecific antibodies in cancer," *Cancer Immuno Immunother*, 45:121-123 (1997).
- De Vries et al., "The fms-Like Tyrosine Kinase, a Receptor for Vascular Endothelial Growth Factor," *Science*, 255:989-991 (Feb. 21, 1992).
- De Waal, R. et al., "Technical Advance: Lack of Lymphangiogenesis in Human Primary Cutaneous Melanoma," *American Journal of Pathology*, 150(6):1951-1957 (Jun., 1997).
- Devereux et al., "A Comprehensive Set of Sequence Analysis Programs for the VAX," *Nucleic Acids Res.*, 12(1):387-395 (1984).
- Dictor, M. et al., "Lymphaticovenous Differentiation in Kaposi's Sarcoma: Cellular Phenotypes by Stage," *American Journal of Pathology*, 130(2):411-417 (Feb., 1988).
- Edgell et al., "Permanent Cell Line Expressing Human Factor VIII-Related Antigen Established by Hybridization," *Proc. Nat'l. Acad. Sci. USA*, 50:3734-3737 (Jun., 1983).
- Eichmann et al., "Molecular cloning of Quack 1 and 2, two quail vascular endothelial growth factor (VEGF) receptor-like molecules," *Gene*, 174(1):3-8 (1996).
- Enholm, B. et al., "Vascular Endothelial Growth Factor-C, a Growth Factor for Lymphatic and Blood Vascular Endothelial Cells," *TCM*, 8(7):292-297 (1998).
- Fanger, M.W. et al., "Bispecific Antibodies," *Critical Reviews in Immunology*, 12(3,4):101-124 (1992).
- Ferrara, N. et al., "The Biology of Vascular Endothelial Growth Factor," *Endocrine Reviews*, 18(1):4-25 (1997).
- Finnerty et al., "Molecular Cloning of Murine FLT and FLT4," *Oncogene*, 8(11):2293-2298 (1993).
- Flanagan & Leder, "The kit Ligand: A Cell Surface Molecule Altered in Steel Mutant Fibroblasts," *Cell*, 63:185-194 (Oct. 5, 1990).
- Folkman et al., "Long-term culture of capillary endothelial cells," *Proc. Nat'l Acad. Sci., USA*, 76(10):5217-5221 (Oct., 1979).
- Foot, J. et al., "Antibody Framework Residues Affecting the Conformation of the Hypervariable Loops," *J. Mol. Biol.*, 224:487-499 (1992).
- Fortkamp et al., "Cloning and Expression in *Escherichia coli* of a Synthetic DNA for Hirudin, the Blood Coagulation Inhibitor in the Leech," *DNA*, 5(6):511-517 (1986).
- Fournier et al., "Mutation at Tyrosine Residue 1337 Abrogates Ligand-Dependent Transforming Capacity of the FLT4 Receptor," *Oncogene*, 11:921-931 (1995).
- Fournier et al., "Interaction with the Phosphotyrosine Binding Domain/Phosphotyrosine Interacting Domain of SHC Is Required for the Transforming Activity of the FLT4/VEGFR3 Receptor Tyrosine Kinase," *J. Biological Chemistry*, 271(22):12956-12963 (1996).
- Gahmberg et al., "Membrane Glycosylation During Cell Differentiation," In L. C. Andersson, et al. (ed.), *Gene Expression During Normal and Malignant Differentiation*, 107-123, Academic Press, London (1985).
- Galland et al., "Chromosomal Localization of FLT4, a Novel Receptor-Type Tyrosine Kinase Gene," *Genomics*, 13:475-478 (1992).
- Galland et al., "The FLT4 Gene Encodes a Transmembrane Tyrosine Kinase Related to the Vascular Endothelial Growth Factor Receptor," *Oncogene*, 8(11):1233-1240 (1993).
- Gasparini, G. et al., "Clinical Importance of the Determination of Tumor Angiogenesis in Breast Carcinoma: Much More Than a New Prognostic Tool," *Journal of Clinical Oncology*, 13(3):765-782 (Mar., 1995).
- Genbank Accession X51602 Human flt mRNA for receptor-related tyrosine kinase, deposited by Shibuya, M.
- Genbank Accession X60280 plasmid pLTRpoly, deposited by Maekelae et al.
- Genbank Accession X68203 *H. sapiens* mRNA for FLT4, Class III receptor tyrosine kinase, deposited by Aprelikova, O.
- Genbank Accession X83287 *C. coturnix* Quack2 mRNA for vascular endothelial growth factor receptor, deposited by Eichmann, et al.
- GenBank Accession No. AJ000185, *Homo sapiens* mRNA for vascular endothelial growth factor-D, deposited by Achen, M.G.

- Greenberg et al., "Characterization of a New Megakaryocyte Cell Line: The Dami Cell," *Blood*, 72(6):1968-1977 (Dec., 1988).
- Harlow et al., *Antibodies: A Laboratory Manual*, pp. 72-137, 141-157, 287 & 321-358 (1988).
- Heldin et al., "Platelet-Derived Growth Factor: Mechanisms of Action and Possible in Vivo Function," *Cell Regulation*, 1:555-566 (Jul., 1990).
- Hemmila et al., "Europium as a Label in Time-Resolved Immunofluorometric Assays," *Anal. Biochem.*, 137:335-343 (1984).
- Hewett, P.W. et al., "Coexpression of flt-1, flt-4 and KDR in Freshly Isolated and Cultured Human Endothelial Cells," *Biochemical and Biophysical Research Communications*, 221:697-702 (1996).
- Hirai et al., "A Novel Putative Tyrosine Kinase Receptor Encoded by the eph Gene," *Science*, 238:1717-1720 (1987).
- Hoogenboom, H.R., "Designing and optimizing library selection strategies for generating high-affinity antibodies," *TIBTECH*, 15:62-70 (1997).
- Huang et al., "The Hematopoietic Growth Factor KL is Encoded by the SI Locus and is the Ligand of the c-kit Receptor, the Gene Product of the W Locus," *Cell*, 63:225-33 (Oct. 5, 1990).
- Hunter & Greenwood, "Preparation of Iodine-131 Labelled Human Growth Hormone of High Specific Activity," *Nature*, 194(4827):495-496 (May 5, 1962).
- Jeltsch, "Hyperplasia of Lymphatic Vessels of VEGF-C Transgenic Mice," *Science*, 276:1423-1425 (May 30, 1997).
- Jones, P.T. et al., "Replacing the complementarity-determining regions in a human antibody with those from a mouse," *Nature*, 321:522-525 (May, 1986).
- Joukov, V. et al., "Proteolytic processing regulates receptor specificity and activity of VEGF-C," *EMBO Journal*, 16(13):3898-3911 (Jun., 1997).
- Joukov et al., "A Novel Vascular Endothelial Growth Factor, VEGF-C, is a Ligand for the Flt4 (VEGFR-3) and KDR (VEGFR-2) Receptor Tyrosine Kinases," *EMBO J.*, 15(2):290-298 (Jan. 15, 1996).
- Jussila, L. et al., "Lymphatic Endothelium and Kaposi's Sarcoma Spindle Cells Detected by Antibodies against the Vascular Endothelial Growth Factor Receptor-3," *Cancer Res.*, 58:1599-1604 (Apr., 1998).
- Kaipainen et al., "Expression of the FMS-Like Tyrosine Kinase 4 Gene Becomes Restricted to Lymphatic Endothelium During Development," *Proc. Nat'l Acad. Sci., USA*, 92:3566-3570 (Apr., 1995).
- Kaipainen et al., "The Related FLT4, FLT1, and KDR Receptor Tyrosine Kinases Show Distinct Expression Patterns in Human Fetal Endothelial Cells," *J. Exp. Med.*, 178:2077-2088 (Dec., 1993).
- Kerstens, H. et al., "A Novel In Situ Hybridization Signal Amplification Method Based on the Deposition of Biotinylated Tyramine," *Journal Histochemistry and Cytochemistry*, 43(4):347-352 (1995).
- Kettleborough, C.A. et al., "Humanization of a mouse monoclonal antibody by CDR-grafting: the importance of framework residues on loop conformation," *Protein Engineering*, 4(7):773-783 (1991).
- Kieffer et al., "Uncoupling in the Expression of Platelet GP IIb/IIIa in Human Endothelial Cells and K562 Cells: Absence of Immunologic Crossreactivity Between Platelet GP IIb and the Vitronectin Receptor Alpha Chain," *Blood*, 72(4):1209-1215 (Oct., 1988).
- Kim, K.J. et al., "Inhibition of vascular endothelial growth factor-induced angiogenesis suppresses tumour growth in vivo," *Nature*, 362(6423):841-844 (Apr., 1993).
- Koeffler et al., "Acute Myelogenous Leukemia: A Human Cell Line Responsive to Colony-Stimulating Activity," *Science*, 200:1153-1154 (1978).
- Köhler et al., "Continuous Cultures of Fused Cells Secreting Antibody of Predefined Specificity," *Nature*, 256:495-497 (Aug. 7, 1975).
- Korhonen et al., "Enhanced Expression of the tie Receptor Tyrosine Kinase in Endothelial Cells During Neovascularization," *Blood*, 80(10):2548-2555 (Nov. 15, 1992).
- Korhonen et al., "The Mouse Tie Receptor Tyrosine Kinase Gene: Expression During Embryonic Angiogenesis," *Oncogene (England)*, 9(2):395-403 (Feb., 1994).
- Kozbor et al., "The Production of Monoclonal Antibodies From Human Lymphocytes," *Immunology Today*, 4(3):72-79 (1983).
- Kukk et al., "VEGF-C Receptor Binding and Pattern of Expression with VEGFR-3 Suggests A Role in Lymphatic Vascular Development," *Development*, 122:3829-3837 (Dec., 1996).
- Lee et al., "Vascular Endothelial Growth Factor-Related Protein: A Ligand and Specific Activator of the Tyrosine Kinase Receptor Flt4," *Proc. Natl. Acad. Sci., USA*, 93:1988-1992 (Mar., 1996).
- Lhotak et al., "Characterization of Elk, a Brain-Specific Receptor Tyrosine Kinase," *Mol. Cell. Biol.*, 11:2496-2502 (May, 1991).
- Lindberg et al., "cDNA Cloning and Characterization of eck, an Epithelial Cell Receptor Protein-Tyrosine Kinase in the eph/elk Family of Protein Kinases," *Mol. Cell. Biol.*, 10:6316-6324 (Dec., 1990).
- Lovgren et al., "Time-Resolved Fluorometry in Immunoassay," In: Collins W.P. (Ed.) *Alternative Immunoassays*, John Wiley & Sons Ltd., pp. 203-217 (1985).
- Lozzio et al., "Human Chronic Myelogenous Leukemia Cell-Line With Positive Philadelphia Chromosome," *Blood*, 45(3):321-334 (Mar., 1975).
- Lyman et al., "Molecular Cloning of a Ligand for the flt3/flt-2 Tyrosine Kinase Receptor: A Proliferative Factor for Primitive Hematopoietic Cells," *Cell*, 75:1157-1167 (Dec. 17, 1993).
- Lymboussaki, A. et al., "Expression of the Vascular Endothelial Growth Factor C Receptor VEGFR-3 in Lymphatic Endothelium of the Skin and in Vascular Tumors," *American Journal of Pathology*, 153(2):395-403 (Aug., 1998).
- Makela et al., "Plasmid pLTRpoly: a Versatile High-Efficiency Mammalian Expression Vector," *Gene*, 118:293-294 (1992).
- Martin et al., "HEL Cells: A New Human Erythroleukemia Cell Line With Spontaneous Induced Globin Expression," *Science*, 216:1233-1235 (1982).
- Matthews et al., "A Receptor Tyrosine Kinase cDNA Isolated From a Population of Enriched Primitive Hematopoietic Cells and Exhibiting Close Genetic Linkage to c-kit," *Proc. Natl. Acad. Sci. USA*, 88(20):9026-9030 (Oct., 1991).
- Matthews et al., "A Receptor Tyrosine Kinase Specific to Hematopoietic Stem and Progenitor Cell-Enriched Populations," *Cell*, 65(7):1143-1152 (Jun. 28, 1991).
- McCutchan et al., "Enhancement of the Infectivity of Simian Virus 40 Deoxyribonucleic Acid with Diethylaminoethyl-Dextran," *J. Natl. Cancer Inst.*, 41:351-357 (1968).

- Metzelaar et al., "CD63 Antigen," *J. Biol. Chem.*, 266(5):3239-3245 (Feb. 15, 1991).
- Millauer et al., High Affinity VEGF Binding and Development Expression Suggest Flk-1 as a Major Regulator of Vasculogenesis and Angiogenesis, *Cell*, 72:835-846 (Mar. 26, 1993).
- Minowada et al., "Brief Communication: Rosette-Forming Human Lymphoid Cell Lines: Establishment and Evidence for Origin of Thymus-Derived Lymphocytes," *J. Natl. Cancer Inst.*, 49:891-895 (1972).
- Mollinedo et al., "Early and Selective Induction of Apoptosis in Human Leukemic Cells By the Alkyl-Lysophospholipid ET-18-OCH₃," *Biochem. & Biophys. Res. Comm.*, 192(2):603-609 (Apr. 30, 1993).
- Moroni et al., "EGF-R Antisense RNA Blocks Expression of the Epidermal Growth Factor Receptor and Suppresses the Transforming Phenotype of a Human Carcinoma Cell Line," *J. Biol. Chem.*, 267(5):2714-2722 (Feb. 5, 1992).
- Morrison, S.L. et al., "Genetically Engineered Antibody Molecules," *Advances in Immunology*, 44:65-92 (1989).
- Moshakis, V. et al., "Localization of human breast-carcinoma xenografts using antibodies to carcinoembryonic antigen," *Br. J. Cancer*, 43:575-581 (1981).
- Mukkala et al., "The Synthesis and Use of Activated N-Benzyl Derivatives of Diethylenetriaminetetraacetic Acids: Alternative Reagents for Labeling of Antibodies with Metal Ions," *Annal. Biochem.*, 176:319-325 (1989).
- Mustonen et al., "Endothelial Receptor Tyrosine Kinases Involved in Angiogenesis," *J. Cell Biology*, 129(4):895-898 (1995).
- Nowell et al., "Chromosome Studies in Preleukemic States: Myeloproliferative versus Cytopenic Disorders," *Cancer*, 42:2254-2260 (1978).
- Oerlich et al., "NYK/FLK-1: A Putative Receptor Protein Tyrosine Kinase Isolated From E10 Embryonic Neuroepithelium is Expressed in Endothelial Cells of the Developing Embryo," *Oncogene*, 8(1):11-18 (Jan., 1993).
- O'Reilly, M.S. et al., "Angiostatin: A Novel Angiogenesis Inhibitor That Mediates the Suppression of Metastases by a Lewis Lung Carcinoma," *Cell*, 79(2):315-328 (Oct., 1994).
- O'Reilly, M.S. et al., "Endostatin: An Endogenous Inhibitor of Angiogenesis and Tumor Growth," *Cell*, 88(2):277-285 (Jan., 1997).
- Oh et al., "VEGF and VEGF-C: Specific Induction of Angiogenesis and Lymphangiogenesis in the Differentiated Avian Chorioallantoic Membrane," *Developmental Biology*, 188:96-109 (1997).
- Padlan, E.A., "A Possible Procedure For Reducing The Immunogenicity Of Antibody Variable Domains While Preserving Their Ligand-Binding Properties," *Molecular Immunology*, 28(4/5):489-498 (1991).
- Pajusola et al., "FLT4 Receptor Tyrosine Kinase Contains Seven Immunoglobulin-Like Loops and is Expressed in Multiple Human Tissues and Cell Lines," *Cancer Research*, 52(20):5783-5743 (Oct. 15, 1992).
- Pajusola et al., "Two Human FLT4 Receptor Tyrosine Kinase Isoforms With Distinct Carboxy Terminal Tails are Produced by Alternative Processing of Primary Transcripts," *Oncogene*, 8:2931-2937 (1993).
- Pajusola, "Cloning and Characterization of a New Endothelial Receptor Tyrosine Kinase FLT4 and Two Novel VEGF-Like Growth Factors VEGF-B and VEGF-C," Molecular/Cancer Biology Laboratory and Department of Pathology, Haartman Institute and Department of Biosciences, Division of Genetics, University of Helsinki, Academic Dissertation, Helsinki 1996.
- Pajusola et al., "Signalling Properties of FLT4, a Proteolytically Processed Receptor Tyrosine Kinase Related To Two VEGF Receptors," *Oncogene*, 9:3545-3555 (1994).
- Partanen et al., "A Novel Endothelial Cell Surface Receptor Tyrosine Kinase with Extracellular Epidermal Growth Factor Homology Domains," *Mol. Cell. Biol.*, 12(4):1698-1707 (Apr. 1992).
- Partanen et al., "Putative Tyrosine Kinases Expressed in K-562 Human Leukemia Cells," *Proc. Nat'l Acad. Sci., USA*, 87(22):8913-8917 (Nov., 1990).
- Perumov et al., "Influence of Antisense RNA's of Interleukin-1 β and Interleukin-1 Receptor Antagonist on Interleukin-1 β Production," *J. Cell. Biochem., Supplement*, 16 pt B:285 (ABSTRACT J216) (1992).
- Peters et al., "Vascular Endothelial Growth Factor Receptor Expression During Embryogenesis and Tissue Repair Suggests a Role in Endothelial Differentiation and Blood Vessel Growth," *Proc. Nat'l Acad. Sci., USA*, 90:8915-8919 (Oct., 1993).
- Pietersz et al., "Antibody Conjugates for the Treatment of Cancer," *Immunological Reviews*, 129:57-80 (1992).
- Plückthun, A. et al., "New protein engineering approaches to multivalent and bispecific antibody fragments," *Immunotechnology*, 3:83-105 (1997).
- Poncz et al., "Cloning and Characterization of Platelet Factor 4 cDNA Derived From a Human Erythroleukemic Cell Line," *Blood*, 69(1):219-223 (Jan., 1987).
- Rader, C. et al., "Phage display of combinatorial antibody libraries," *Curr. Opin. Biotech.*, 8:503-508 (1997).
- Reedijk et al., "Tyr721 Regulates Specific Binding of the CSF-1 Receptor Kinase Insert to P1 3'-Kinase SH2 Domains: a Model for SH2-Mediated Receptor-Target Interactions," *EMBO J.*, 11(4):1365-1372 (1992).
- Relf, M. et al., "Expression of the Angiogenic Factors Vascular Endothelial Cell Growth Factor, Acidic and Basic Fibroblast Growth Factor, Tumor Growth Factor β -1, Platelet-derived Endothelial Cell Growth Factor, Placenta Growth Factor, and Pleiotrophin in Human Primary Breast Cancer and Its Relation to Angiogenesis," *Cancer Research*, 57:963-969 (Mar. 1997).
- Renner, C. et al., "Tumor Therapy by Immune Recruitment with Bispecific Antibodies," *Immunological Reviews*, No. 145, pp. 179-209 (1995).
- Riechmann, L. et al., "Reshaping human antibodies for therapy," *Nature*, 332(6162):323-327 (Mar., 1988).
- Roitt, M., "Essential Immunology," Blackwell Scientific Pub., Oxford, pp. 65-68 & 74 (1991).
- Rosnet et al., "Isolation and Chromosomal Localization of a Novel FMS-Like Tyrosine Kinase Gene," *Oncogene*, 6(9):1641-1650 (1991).
- Rosnet et al., "Murine Flt3, a Gene Encoding a Novel Tyrosine Kinase Receptor of the PDGF/CSF1R Family," *Genomics*, 9:380-385 (1991).
- Sabin, F.R., "The Lymphatic System in Human Embryos, with Consideration of the Morphology of the System as a Whole," *Am. J. Anat.*, 9(1):43-91 (1909).

- Salven, P. et al., "Vascular Endothelial Growth Factors VEGF-B and VEGF-C Are Expressed in Human Tumors," *American J. Pathology*, 153(1):103-108 (Jul., 1998).
- Sambrook et al., *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, pp. 2.60-2.70, 4.21-4.32, 7.3-7.36 (1989).
- Sato and Seiki, "Regulatory mechanism of 92 kDa type IV collagenase gene expression which is associated with invasiveness of tumor cells," *Oncogene*, 8:395-405 (1993).
- Satoh et al., "Regional Localization of the Human c-ros-1 on 6q22 and flt on 13q12," *Jpn. J. Cancer Res.*, 78:772-775 (1987).
- Schneider et al., "A One-step Purification of Membrane Proteins Using a High Efficiency Immunomatrix," *J. Biol. Chem.*, 257(18):10766-10769 (Sep. 25, 1982).
- Schreiber et al., "Interaction of endothelial cell growth factor with heparin: Characterization by receptor and antibody recognition," *Proc. Nat'l Acad. Sci.*, 82:6138-6142 (Sep., 1985).
- Schwenk et al., "Cell Cycle Dependency of a T-Cell Marker on Lymphoblasts," *Blut*, 31:299-306 (1975).
- Scott, P.A.E. et al., "Laboratory-Clinic Interface: Current approaches to targeting cancer using antiangiogenesis therapies," *Cancer Treatment Reviews*, 20:393-412 (1994).
- Segal, D.M. et al., "Alternative Triggering Molecules and Single Chain Bispecific Antibodies," *Journal of Hematology*, 4:377-382 (1995).
- Segal, D.M. et al., "Targeting of Anti-Tumor Responses with Bispecific Antibodies," *Immunobiology*, 185(2-4):390-402 (Aug., 1992).
- Sherr et al., "The c-fms Proto-Oncogene Product is Related to the Receptor for the Mononuclear Phagocyte Growth Factor, CSF-1," *Cell*, 41:665-676 (Jul., 1985).
- Shi et al., "16-Kilodalton Heparin Binding (Fibroblast) Growth Factor Type One Appears in a Stable 40-Kilodalton Complex After Receptor-Dependent Internalization," *J. Biol. Chem.*, 266(9):5774-5779 (Mar. 25, 1991).
- Shibuya et al., "Nucleotide Sequence and Expression of a Novel Human Receptor-Type Tyrosine Kinase Gene (flt) Closely Related to the fms Family," *Oncogene*, 5:519-524 (1990).
- Shibuya, M., "Role of VEGF-FLT Receptor System in Normal and Tumor Angiogenesis," *Advances in Cancer Research*, 67:281-316 (1995).
- Stacey et al., "SVpoly: a Versatile Mammalian Expression Vector," *Nucl. Acids Res.*, 18(9):2829 (1990).
- Staunton et al., "The Arrangement of the Immunoglobulin-Like Domains of ICAM-1 and the Binding Sites for LFA-1 and Rhinovirus," *Cell*, 61:243-254 (Apr. 20, 1990).
- Stenman et al., "Human PDGFA Receptor Gene Maps to the Same Region on Chromosome 4 as the KIT Oncogene," *Genes, Chromosomes, Cancer*, 1:155-158 (1989).
- Sundström et al., "Establishment and Characterization of a Human Histiocytic Lymphoma Cell Line (U-937)," *Int. J. Cancer*, 17:565-577 (1976).
- Swolin et al., "On the 5q-Deletion: Clinical and Cytogenetic Observations in Ten Patients and Review of the Literature," *Blood*, 58:986-993 (1981).
- Tekmal, R.R. et al., "A novel in vitro and in vivo breast cancer model for testing inhibitors of estrogen biosynthesis and its action using mammary tumor cells with an activated int-5/aromatase gene," *Cancer Letters*, 118(1):21-28 (Sep., 1997).
- Tempest, P.R. et al., "Reshaping a Human Monoclonal Antibody to Inhibit Human Respiratory Syncytial Virus Infection in Vivo," *BioTechnology*, 9(3):266-271 (Mar., 1991).
- Terman et al., "Identification of a New Endothelial Cell growth Factor Receptor Tyrosine Kinase," *Oncogene*, 6(9):1677-1683 (1991).
- Terman et al., "Identification of the KDR Tyrosine Kinase as a Receptor for Vascular Endothelial Cell Growth Factor," *Biochem & Biophys. Res. Comm.*, 187(3):1579-1586 (Sep. 30, 1992).
- Thompson et al., "Cloned Human Teratoma Cells Differentiate into Neuron-Like Cells and Other Cell Types in Retinoic Acid," *J. Cell Sci.*, 72:37-64 (1984).
- Tomiyasu et al., "Long Arm Deletion of Chromosome No. 5 in a Case of Philadelphia Chromosome-Positive Chronic Myelocytic Leukemia," *Cancer Genet. Cytogenet.*, 2:309-315 (1980).
- Tortora et al., "Differentiation of HL-60 Leukemia By Type I Regulatory Subunit Antisense Oligodeoxynucleotide of a cAMP-Dependent Protein Kinase," *Proc. Nat'l Acad. Sci., USA*, 88(5):2011-2015 (Mar., 1991).
- Trautnecker, A. et al., "Myeloma based expression system for production of large mammalian proteins," *Trends in Biotechnology*, 9(4):109-113 (Apr., 1991).
- Ullrich et al., "Signal Transduction By Receptors with Tyrosine Kinase Activity," *Cell*, 61:203-212 (Apr. 20, 1990).
- van der Putte, S.C.J., "The Development of the Lymphatic System in Man," *Adv. Anat. Embryol. Cell Biol.*, 51:3 (1975).
- Van Den Berghe et al., "Distinct Haematological Disorder with Deletion of Long Arm of No. 5 Chromosome," *Nature*, 251:437-439 (1974).
- Van Den Berghe et al., "Transformation of Polycythemia Vera to Myelofibrosis and Late Appearance of a 5q-Chromosome Anomaly," *Cancer Genet. Cytogenet.*, 1:157-162 (1979).
- Van Hinsberg et al., "Effect of Thrombin on the Production of Plasminogen Activators and PA Inhibitor-1 by Human Foreskin Microvascular Endothelial Cells," *Thromb. Haemostas.*, 57(2):148-153 (1987).
- Van Hinsberg et al., "Production of Plasminogen Activators and Inhibitors by Serially Propagated Endothelial Cells From Adult Human Blood Vessels," *Arteriosclerosis*, 7:389-400 (Jul./Aug., 1987).
- Verhoeven, M. et al., "Reshaping Human Antibodies: Grafting an Antilysozyme Activity," *Science*, 239:1534-1536 (Mar., 1988).
- Wang, "Signal Transduction in Human Hematopoietic Cells by Vascular Endothelial Growth Factor Related Protein, a Novel Ligand for the FLT4 Receptor," *Blood*, 90(9):3507-3515 (Nov., 1997).
- Warrington et al., "Radiation of Hybrid Map of 13 Loci on the Long Arm of Chromosome 5," *Genomics*, 11:701-708 (1991).
- Weidner, N. et al., "Tumor Angiogenesis: A New Significant and Independent Prognostic Indicator in Early-Stage Breast Carcinoma," *J. Natl. Cancer Inst.*, 84(24):1875-1887 (Dec., 1992).
- Wen, D. et al., "New Differentiation Factor: A Transmembrane Glycoprotein Containing an EGF Domain and an Immunoglobulin Homology Unit," *Cell*, 69:559-572 (May 1, 1992).

- Whang-Feng et al., "Cytogenic Studies in Patients With Myelofibrosis and Myeloid Metaplasia," *Leuk. Res.*, 2:41-48 (1978).
- Wilkinson et al., "Expression of the Proto-Oncogene int-1 is Restricted to Specific Neural Cells in the Developing Mouse Embryo," *Cell*, 50:79-88 (1987).
- Williams et al., "The Immunoglobulin Superfamily-Domains for Cell Surface Recognition," *Ann. Rev. Immunol.*, 6:381-405 (1988).
- Williams, J.C. et al., "N-methyl-N-nitrosourea-Induced Rat Mammary Tumors. Hormonal Responsiveness but Lack of Spontaneous Metastasis," *J. Nat. Cancer Inst.*, 66(1):147-155 (Jan., 1981).
- Yamaguchi et al., "Flk-1, an Flt-Related Tyrosine Kinase is an Early Marker for Endothelial Cell Precursors," *Development*, 118:489-498 (1993).
- Yarden et al., "Human Proto-Oncogene c-kit: A New Cell Surface Receptor Tyrosine Kinase for an Unidentified Ligand," *EMBO J.*, 6(11):3341-3351 (1987).
- Ylänne et al., "Platelet Glycoprotein IIb/IIIa Complex in Cultured Cells: Localization in Focal Adhesion Sites in Spreading HEL Cells," *Blood*, 72: 1478-1486 (1988).
- Hatva et al., "Expression of Endothelial Cell-Specific Receptor Tyrosine Kinases and Growth Factors in Human Brain Tumors," *Am. J. Pathol.*, 146:368-378 (1995).
- Fielder et al., "Expression of FLT4 and its ligand VEGF-C in acute myeloid leukemia," *Leukemia*, 11(8):1234-1237 (Aug., 1977).
- Dias, et al., "Vascular endothelial growth factor (VEGF)-C signaling through Flt-4 (VEGFR-3) mediates leukemic cell proliferation, survival, and resistance to chemotherapy," *Blood*, 99:2179-2184 (2002).
- Karkkainen, et al., "Lymphatic endothelium: a new frontier of metastasis research," *Nat. Cell. Biol.* 4:E2-E5 (2002).
- Karpanen, et al., "Vascular endothelial growth factor C promotes tumor lymphangiogenesis and intralymphatic tumor growth," *Cancer Res.* 61:1786-90 (2001).
- Leu, et al., "Absence of functional lymphatics within a murine sarcoma: a molecular and functional evaluation," *Cancer Res.* 60:4324-7 (2000).
- Makinen, et al., "Inhibition of lymphangiogenesis with resulting lymphedema in transgenic mice expressing soluble VEGF receptor-3," *Nat. Med.* 7:199-205 (2001).
- Mattila, et al., "VEGF-C induced lymphangiogenesis is associated with lymph node metastasis in orthotopic MCF-7 tumors," *Int. J. Cancer.* 98:946-951 (2002).
- Siemeister, et al., "Two independent mechanisms essential for tumor angiogenesis: inhibition of human melanoma xenograft growth by interfering with either the vascular endothelial growth factor receptor pathway or the Tie-2 pathway," *Cancer Res.* 59:3185-91 (1999).
- Witmer, et al., "VEGFR-3 in adult angiogenesis," *J. Path.* 195:490-497 (2001).

* cited by examiner

FIG. 1A




FIG. 1B




FIG. 2A

FLT4	1	MQ..RGAALCLRLWLCGLLDGLVSGYSMTPTLNITEESHVIDTGDSLS	48	SS
FLT1	1	MVSYWDTGVLLCALLSCLLLTGSSSGSKLKDPELSLKGTOHIMQAGQTLH	50	
FLT4	49	ISCRGQHPLEWAWPGAQEAAPATGDKDSEDTCVVRDCEGTDARPYCKVLLL	98	
FLT1	51	LQCRGEAAHKWSLPE.....MVSKESERLSITKSACGRNGKQFCSTLT	94	Ig I
FLT4	99	HEVHANDTGSYVCCYKYIKARIEGTTAASSYVFVRDFEQPFINK...PD	144	
FLT1	95	NTAQANHTGFYSCKYLAVPSTSKKKETESAIYIFISDTGRPFVEMYSEIPE	144	
FLT4	145	TLLVNRKDAMWVPCCLVSIPGLNVTLS.QSSVLWPDGQEVVWDDRRGMLV	193	
FLT1	145	IIHMTGRELVIPCRVTSFNITVTLKKFPLDTLIPDGKRIIWDSRKGFII	194	Ig II
FLT4	194	STPLLHDALYLQCEETTWGDQDFLSNPFLVHITGNELYDIQLLPKRSLELL	243	
FLT1	195	SNATYKEIGLLTCEATVNGHLYKTN.YLTHRQTNTIIDVQISTPRPVKLL	243	

FIG. 2B

FLT4	244	VGEKLVLC	TVWAEFNSGVTFDWDYPGKQAE	RGKWPERR...	SQQTHT	289	
				
FLT1	244	RGHTLVLC	TATPLNTRVQMTWSYPD...	EKNKRASVRRRIDQSN	SHAN	290	
				
FLT4	290	ELSSILTIHNV	SOHDLGSYVC	KANNGIQRFRE	STEVIVHENPFIS	VEWLK	339
		:		
FLT1	291	IFYSVLTIDK	MQNKDKGLYTC	RVRS	SGPSFKSVNTSVHIYDKAFITV	KHRK	340
		
FLT4	340	GPILEATAGDE	LVKLPVKLAAYPPPEFQWKDGKAL	SGRHS...	PHAL	384	
		
FLT1	341	QQVLETVAGK	RSYRLSMKVKAFPSPEVVLKDG	LPATEKSARYL	TRGYSL	390	
		
FLT4	385	VLKEVTEASTG	TYTLALWN	SAAGLRRNISLE	LVNVVPPQIHE	KEASS...	431
		:		
FLT1	391	IIKDVT	EEDAGNYTILL	SIKQSNVFNLTATLIVNVK	PKPIYEKAV	SSFPD	440
		

Ig III

Ig IV

FIG. 2C

FLT4	432	PSIYSRHSRQALTC	●	CTAYGVPLPLSIQWHWRPWPCKMFAQSLRRRQQD	481	
FLT1	441	PALYPLGSRQILTCT		AYGIQP.TIKWFHPCNHNHSEARCDFCSNNEES	489	
FLT4	482	LMPOCRDWRAVTTQDAVNPIESLDTWTEFVEGKNKTVSKLVIQNAVVSAM		531		
FLT1	490	FILD.....ADSNMGNRIESITQRMATIEGKNKMASTLVVADSRI		532		
FLT4	532	YKCVVSNKVGQDERLIYFYVTTIPDGFTIESKPSSEELLEGPVLLSC		581		
FLT1	533	YICIASNKVGTVGRNISFYITDVPNGFHVNLEKMPT..EGEDLKLSC		580		
FLT4	582	SYKYEHLRWYRLNLSTLHDAHGNPLLLDCKNVHLFATPLAASLEEVAPGA		631		
FLT1	581	KFLYRDVTWILL.....RTVNNRMTMHYSISKQKMAITK		613		
FLT4	632	RHA.TLSLSIPRVAPHEGHHYV	●	CEVQDRRSHDKHCHKKYLVSQALEAPRL	680	
FLT1	614	EHSITLNLTIMNVSLQDSGTYAC		RARNVYTGEELQKKEITIRDQEAPYL	663	

Ig V

Ig VI

FIG. 2D

FLT4	681	TQNLTDLLVNVSDSLEMQ	CLVAGAHAPSIVWYKDERLLEEKSGVDLADSN	730	Ig VII
FLT1	664	LRNLS DHTVAISSSTLDC	CHANGVPEPQITWFKNNHKIQQEPGIILGPS	713	
FLT4	731	QKLSIQRVREEDAGRYL	CSVCNAKGCNVSSASVAVEGSEDKGSMEIVLV	780	
FLT1	714	STLFIERVTEDEGVYHC	KATNQKGSVESSAYLTVQGTSDKSNLELITLT	763	
FLT4	781	GTGVIAVFFWVLLLLIFC	NMRRPAHADIKTGYSIIMDPGEVPLEEQCEY	830	TM
FLT1	764	CTCVAATLEFWLLLLIRKMKRSS	.SEIKTDYLSIIMDPDEVPLDEQCER	812	
FLT4	831	LSYDASQWEFFRERLHGRVL	GYGAFGKVVVEASAFGIHKGSSCDTVAVKM	880	TK 1
FLT1	813	LPYDASKWEFARERLKLKSLGRGAFGKVVQASAFGIKKSPTCRTVAVKM	862		
FLT4	881	LKEGATASEHRALMSELKILIHIGNHLNVNLLGACTKPKQGPLMVIVEFC	930		
FLT1	863	LKEGATASEYKALMTELKILTHIGHHLNVNLLGACTKQGGPLMVIVEYC	912		
FLT4	931	KYGNLSNFLRAKRD	AF...SPCAEKSPQGRFRAMVELARLDRRRPGSS	977	KI
FLT1	913	KYGNLSNYLKS KRD	LFFLNKNDAALHMEPKKEKMEPGLEQGKKPRLDSVTS	962	

FIG. 2E

[illegible]

FIG. 2F

FLT4	1178	LQEEEEVC	MAPRSSQ	SSSEEGS	FQVSTMA	LHIAQADA	EDSPPSL	QQRHSLA	1227
FLT1	1163	QQDGKDY	IPINAIL	TGNSGFT	YSTPAF	SEDFFKES	ISAPKF	NSGSSDDVR	1212
FLT4	1228	ARYYNWV	SFPGCL	ARGAET	RGSSRM	KTFEEFP	MTPTTY	KGSVDNQ	1277
FLT1	1213	YVNAFK	FMSLER	IKTFEE	LLPNAT	SMFDDY	QGDSS	TLLASP	1262
FLT4	1278	VLASEE	FEQIES	RHRQES	CGER				
FLT1	1263	SKPKAS	LKIEV						

FIG. 3A




FIG. 3B




FIG. 3C





FIG. 4

FIG. 5E



FIG. 5F




FIG. 5G




FIG. 5H




FIG. 5A




FIG. 5B




FIG. 5C





FIG. 5D



FLT4 (VEGFR-3) AS A TARGET FOR TUMOR IMAGING AND ANTI-TUMOR THERAPY

This application is a continuation-in-part of U.S. patent application Ser. No. 08/901,710, filed Jul. 28, 1997, now U.S. Pat. No. 6,107,046; which is a continuation-in-part of both U.S. patent application Ser. No. 08/340,011, filed Nov. 14, 1994, now U.S. Pat. No. 5,776,755; and U.S. patent application Ser. No. 08/257,754, filed Jun. 9, 1994, now abandoned; the latter two of which in turn are continuations-in-part of U.S. patent application Ser. No. 07/959,951, filed on Oct. 9, 1992, now abandoned. All of these applications are incorporated herein by reference, in their entireties.

FIELD OF THE INVENTION

The present invention relates generally to genes for receptors, specifically genes for receptor tyrosine kinases, their insertion into recombinant DNA vectors, and the production of the resulting proteins in host strains of microorganisms and host eukaryotic cells. More specifically, the present invention is directed to Flt4, a receptor tyrosine kinase; to nucleotide sequences encoding Flt4; to methods for the generation of DNAs encoding Flt4 and their gene products; to nucleic acid probes which specifically recognize (hybridize to) nucleic acids encoding such receptors; to antibodies that specifically recognize such receptors; and to methods of using such probes and antibodies, e.g., for identifying lymphatic vessels and high endothelial venules (HEV) in animal and human tissues and augmenting or preventing their growth in pathological conditions.

BACKGROUND

The cellular behavior responsible for the development, maintenance and repair of differentiated cells and tissues is regulated, in large part, by intercellular signals conveyed via growth factors and similar ligands and their receptors. The receptors are located on the cell surface of responding cells and they bind peptides or polypeptides known as growth factors as well as other hormone-like ligands. The results of this interaction are rapid biochemical changes in the responding cells, as well as a rapid and a long term readjustment of cellular gene expression. Several receptors associated with various cell surfaces can bind specific growth factors.

Tyrosine phosphorylation is one of the key modes of signal transduction across the plasma membrane. Several tyrosine kinase genes encode transmembrane receptors for polypeptide growth factors and hormones, such as epidermal growth factor (EGF), insulin, insulin-like growth factor-I (IGF-I), platelet derived growth factors (PDGF-A and -B) and fibroblast growth factors (FGFs) [Heldin et al., *Cell Regulation*, 1: 555-566 (1990); Ullrich et al., *Cell*, 61: 243-54 (1990)]. The receptors of several hematopoietic growth factors are tyrosine kinases; these include c-fms, which is the colony stimulating factor 1 receptor [Sherr et al., *Cell*, 41: 665-676 (1985)] and c-kit, a primitive hematopoietic growth factor receptor [Huang et al., *Cell*, 63: 225-33 (1990)].

These receptors differ in their specificity and affinity. In general, receptor tyrosine kinases are glycoproteins, which consist of an extracellular domain capable of binding a specific growth factor(s), a transmembrane domain which is usually an alpha-helical portion of the protein, a juxtamembrane domain (where the receptor may be regulated by, e.g., protein phosphorylation), a tyrosine kinase domain (which is the enzymatic component of the receptor), and a carboxy-

terminal tail, which in many receptors is involved in recognition and binding of the substrates for the tyrosine kinase.

In several receptor tyrosine kinases, the processes of alternative splicing and alternative polyadenylation are capable of producing several distinct polypeptides from the same gene. These may or may not contain the various domains listed above. As a consequence, some extracellular domains may be expressed as separate proteins secreted by the cells and some forms of the receptors may lack the tyrosine kinase domain and contain only the extracellular domain inserted into the plasma membrane via the transmembrane domain plus a short carboxy-terminal tail.

The physiology of the vascular system, embryonic vasculogenesis and angiogenesis, blood clotting, wound healing and reproduction, as well as several diseases, involve the vascular endothelium lining the blood vessels. The development of the vascular tree occurs through angiogenesis, and, according to some theories, the formation of the lymphatic system starts shortly after arterial and venous development by sprouting from veins. See Sabin, F. R., *Am. J. Anat.*, 9:43 (1909); and van der Putte, S. C. J., *Adv. Anat. Embryol. Cell Biol.*, 51:3 (1975).

After the fetal period, endothelial cells proliferate very slowly, except during angiogenesis associated with neovascularization. Growth factors stimulating angiogenesis exert their effects via specific endothelial cell surface receptor tyrosine kinases.

Among ligands for receptor tyrosine kinases, the Platelet Derived Growth Factor (PDGF) has been shown to be angiogenic, albeit weakly, in the chick chorioallantoic membrane. Transforming Growth Factor α (TGF α) is an angiogenic factor secreted by several tumor cell types and by macrophages. Hepatocyte Growth Factor (HGF), the ligand of the c-met proto-oncogene-encoded receptor, is also strongly angiogenic, inducing similar responses to those of TGF α in cultured endothelial cells.

Striking new evidence shows that there are endothelial cell specific growth factors and receptors that may be primarily responsible for the stimulation of endothelial cell growth, differentiation, as well as certain of differentiated functions. The most-widely studied growth factor is Vascular Endothelial Growth Factor (VEGF), a member of the PDGF family. Vascular endothelial growth factor is a dimeric glycoprotein of disulfide-linked 23 kDa subunits, discovered because of its mitogenic activity toward endothelial cells and its ability to induce vessel permeability (hence its alternative name vascular permeability factor). Other reported effects of VEGF include the mobilization of intracellular Ca^{2+} , the induction of plasminogen activator and plasminogen activator inhibitor-1 synthesis, stimulation of hexose transport in endothelial cells, and promotion of monocyte migration in vitro. Four VEGF isoforms, encoded by distinct mRNA splicing variants, appear to be equally capable of stimulating mitogenesis of endothelial cells. The 121 and 165 amino acid isoforms of VEGF are secreted in a soluble form, whereas the isoforms of 189 and 206 amino acid residues remain associated with the cell surface and have a strong affinity for heparin. Soluble non-heparin-binding and heparin-binding forms have also been described for the related placenta growth factor (PlGF; 131 and 152 amino acids, respectively), which is expressed in placenta, trophoblastic tumors, and cultured human endothelial cells.

The pattern of VEGF expression suggests its involvement in the development and maintenance of the normal vascular system and in tumor angiogenesis. During murine development, the entire 7.5 day post-coital endoderm

expresses VEGF and the ventricular neuroectoderm produces VEGF at the capillary ingrowth stage. On day two of quail development, the vascularized area of the yolk sac as well as the whole embryo show expression of VEGF. In addition, epithelial cells next to fenestrated endothelia in adult mice show persistent VEGF expression, suggesting a role in the maintenance of this specific endothelial phenotype and function.

Two high affinity receptors for VEGF have been characterized, VEGFR-1/Flt1 (fms-like tyrosine kinase-1) and VEGFR-2/Kdr/Flk-1 (kinase insert domain containing receptor/fetal liver kinase-1). These receptors are classified in the PDGF-receptor family. However, the VEGF receptors have seven immunoglobulin-like loops in their extracellular domains (as opposed to five in other members of the PDGF family) and a longer kinase insert. The expression of VEGF receptors occurs mainly in vascular endothelial cells, although some may also be present on monocytes and on melanoma cell lines. Only endothelial cells have been reported to proliferate in response to VEGF, and endothelial cells from different sources show different responses. Thus, the signals mediated through VEGFR-1 and VEGFR-2 appear to be cell type specific.

VEGFR-1 and VEGFR-2 bind VEGF 165 with high affinity (K_d about 20 pM and 200 pM, respectively). Flk-1 receptor has also been shown to undergo autophosphorylation in response to VEGF, but phosphorylation of Flt1 was barely detectable. VEGFR-2 mediated signals cause striking changes in the morphology, actin reorganization and membrane ruffling of porcine aortic endothelial cells overexpressing this receptor. In these cells, VEGFR-2 also mediated ligand-induced chemotaxis and mitogenicity; whereas VEGFR-1 transfected cells lacked mitogenic responses to VEGF. In contrast, VEGF had a strong growth stimulatory effect on rat sinusoidal endothelial cells expressing VEGFR-1. Phosphoproteins co-precipitating with VEGFR-1 and VEGFR-2 are distinct, suggesting that different signalling molecules interact with receptor specific intracellular sequences.

In situ hybridization studies, mouse VEGFR-2 mRNA expression was found in yolk sac and intraembryonic mesoderm (estimated 7.5 day post-coitum (p.c.) embryos, from which the endothelium is derived, and later in presumptive angioblasts, endocardium and large and small vessel endothelium (12.5 days p.c.). Abundant VEGFR-2 mRNA in proliferating endothelial cells of vascular sprouts and branching vessels of embryonic and early postnatal brain and decreased expression in adult brain suggested that VEGFR-2 is a major regulator of vasculogenesis and angiogenesis. VEGFR-1 expression was similarly associated with early vascular development in mouse embryos and with neovascularization in healing skin wounds. However, high levels of VEGFR-1 expression were detected in adult organs, suggesting that VEGFR-1 has a function in quiescent endothelium of mature vessels not related to cell growth. The avian homologue of VEGFR-2 was observed in the mesoderm from the onset of gastrulation, whereas the VEGFR-1 homologue was first found in cells co-expressing endothelial markers. In vitro quail epiblast cultures, FGF-2, which is required for vasculogenic differentiation of these cells, upregulated VEGFR-2 expression. The expression of both VEGF receptors was found to become more restricted later in development. In human fetal tissues VEGFR-1 and VEGFR-2 showed overlapping, but slightly different, expression patterns. These data suggest that VEGF and its receptors act in a paracrine manner to regulate the differentiation of endothelial cells and neovascularization of tissues.

VEGF recently has been shown to be a hypoxia-induced stimulator of endothelial cell growth and angiogenesis, and inhibition of VEGF activity using specific monoclonal antibodies has been shown to reduce the growth of experimental tumors and their blood vessel density. [Ferrara et al., *Endocrine Reviews*, 18: 4-25 (1997); Shibuya et al., *Adv. Cancer Res.*, 67: 281-316 (1995); Kim et al., *Nature*, 362: 841-844 (1993).]

Growth of solid tumors beyond a few cubic millimeters in size is dependent on vascular supply, making angiogenesis an attractive target for anti-cancer therapy. Encouraging results have been reported with endogenous angiogenic inhibitors or "statins" which include angiostatin, a fragment of plasminogen, and endostatin, a fragment of collagen 18. [O'Reilly et al., *Cell*, 79: 315-328 (1994); O'Reilly et al., *Cell*, 88: 277-85 (1997).]. Both factors are normally produced by primary tumors and keep metastasis dormant. Systemic administration of either statin has been shown to also induce and sustain dormancy of primary tumors in animal models. The receptors and signalling by statins, as well as the proteases which activate them, remain to be identified. A need exists for additional therapeutic molecules for controlling angiogenesis in the treatment of cancer and other pathological disease states.

Primary breast cancers have been shown to express several angiogenic polypeptides, of which VEGF was the most abundant. [See, e.g., Relf et al., *Cancer Res.*, 57: 963-969 (1997).] Tumor cells contained high levels of VEGF mRNA in both invasive and non-invasive, ductal (in situ) breast carcinomas. [Brown et al., *Hum. Pathol.*, 26: 86-91 (1995).] The endothelial cells adjacent to the in situ carcinomas expressed VEGFR-1 and VEGFR-2 mRNA. VEGF and its receptors may contribute to the angiogenic progression of malignant breast tumors, because in several independent studies, correlations have been found between tumor vascular density and the prognosis of the disease. [Weidner et al., *J. Natl. Cancer Inst.*, 84: 1875-1887 (1992).] A need exists for additional markers for breast cancer and breast cancer-related angiogenesis, to improve diagnosis and screening and to serve as a target for therapeutic intervention.

A major function of the lymphatic system is to provide fluid return from tissues and to transport many extravascular substances back to the blood. In addition, during the process of maturation, lymphocytes leave the blood, migrate through lymphoid organs and other tissues, and enter the lymphatic vessels, and return to the blood through the thoracic duct. Specialized venules, high endothelial venules (HEVs), bind lymphocytes again and cause their extravasation into tissues. The lymphatic vessels, and especially the lymph nodes, thus play an important role in immunology and in the development of metastasis of different tumors.

Since the beginning of the 20th century, three different theories concerning the embryonic origin of the lymphatic system have been presented. However, lymphatic vessels have been difficult to identify, due to the absence of known specific markers available for them.

Lymphatic vessels are most commonly studied with the aid of lymphography. In lymphography, X-ray contrast medium is injected directly into a lymphatic vessel. That contrast medium is distributed along the efferent drainage vessels of the lymphatic system. The contrast medium is collected in lymph nodes, where it stays for up to half a year, during which time X-ray analyses allow the follow-up of lymph node size and architecture. This diagnostic is especially important in cancer patients with metastases in the

lymph nodes and in lymphatic malignancies, such as lymphoma. However, improved materials and methods for imaging lymphatic tissues are needed in the art.

SUMMARY OF THE INVENTION

The present invention addresses a gene for a novel receptor tyrosine kinase located on chromosome 5, identified as an unknown tyrosine kinase-homologous PCR-cDNA fragment from human leukemia cells [Aprelikova et al., *Cancer Res.*, 52: 746-748 (1992)]. This gene and its encoded protein are called Flt4. This abbreviation comes from the words fms-like tyrosine kinase 4.

Flt4 is a receptor tyrosine kinase closely related in structure to the products of the VEGFR-1 and VEGFR-2 genes. By virtue of this similarity and subsequently-discovered similarities between ligands for these three receptors, the Flt4 receptor has additionally been named VEGFR-3. The names Flt4 and VEGFR-3 are used interchangeably herein. Despite the similarity between these three receptors, the mature form of Flt4 differs from the VEGFRs in that it is proteolytically cleaved in the extracellular domain into two disulfide-linked polypeptides of 125/120 kD and 75 kD. The Flt4 gene encodes 4.5 and 5.8 kb mRNAs which exhibit alternative 3' exons and encode polypeptides of 190 kD and 195 kD, respectively.

Further evidence of a distinction is that VEGF does not show specific binding to Flt4 and doesn't induce its autophosphorylation.

A comparison of Flt4, Flt1, and KDR/Flk-1 receptor mRNA signals showed overlapping, but distinct expression patterns in the tissues studied. Kaipainen, et al., *J. Exp. Med.*, 178:2077 (1993). Flt4 gene expression appears to be more restricted than the expression of VEGFR-1 or VEGFR-2. The expression of Flt4 first becomes detectable by in situ hybridization in the angioblasts of head mesenchyme, the cardinal vein and extraembryonically in the allantois of 8.5 day post-coital mouse embryos. In 12.5 day post-coital embryos the Flt4 signal is observed on developing venous and presumptive lymphatic endothelia, but arterial endothelia appear to be negative. During later stages of development, Flt4 mRNA becomes restricted to developing lymphatic vessels. Only the lymphatic endothelia and some high endothelial venules express Flt4 mRNA in adult human tissues and increased expression occurs in lymphatic sinuses in metastatic lymph nodes and in lymphangioma. The results support the theory of the venous origin of lymphatic vessels.

The protein product of the Flt4 receptor tyrosine kinase cDNA, cloned from a human erythroleukemia cell line, is N-glycosylated and contains seven immunoglobulin-like loops in its extracellular domain. The cytoplasmic tyrosine kinase domain of Flt4 is about 80% identical at the amino acid level with the corresponding domains of Flt1 and KDR and about 60% identical with the receptors for platelet-derived growth factor, colony stimulating factor-1, stem cell factor, and the Flt3 receptor. See Pajusola et al., *Cancer Res.*, 52:5738 (1992).

The present invention provides isolated polynucleotides (e.g., DNA or RNA segments of defined structure) encoding an Flt4 receptor tyrosine kinase useful in the production of Flt4 protein and peptide fragments thereof and in recovery of related genes from other sources.

The present invention provides a recombinant DNA vector containing a heterologous segment encoding the Flt4 receptor tyrosine kinase or a related protein that is capable of being inserted into a microorganism or eukaryotic cell and that is capable of expressing the encoded protein.

The present invention provides eukaryotic cells capable of producing useful quantities of the Flt4 receptor tyrosine kinase and proteins of similar function from many species.

The present invention provides peptides that may be produced synthetically in a laboratory or by microorganisms, which peptides mimic the activity of the natural Flt4 receptor tyrosine kinase protein. In another embodiment, the invention is directed to peptides which inhibit the activity of Flt4 receptor tyrosine kinase protein.

Particularly preferred are peptides selected from the group consisting of: (a) a Flt4-short form, the nucleotide and deduced amino acid sequence of which appear in SEQ. ID NOs. 1 and 2; and (b) a second form with different nucleotide and corresponding amino acid residues at its carboxyl terminal, i.e., an Flt4-long form, the nucleotide and deduced amino acid sequence of which appear in SEQ. ID NOs. 3 and 4. The Flt4 long form has a length of 1363 amino acid residues.

DNA and RNA molecules, recombinant DNA vectors, and modified microorganisms or eukaryotic cells comprising a nucleotide sequence that encodes any of the proteins or peptides indicated above are also part of the present invention. In particular, sequences comprising all or part of the following two DNA sequences, a complementary DNA or RNA sequence, or a corresponding RNA sequence are especially preferred: (a) a DNA sequence such as SEQ ID NO: 1, encoding Flt4-short form [SEQ ID NO: 2], and (b) a DNA sequence such as SEQ ID NO: 3, encoding a Flt4 wherein nucleotides 3913-4416 of SEQ ID NO: 1 are changed, encoding Flt4-long form [SEQ ID NO: 4].

DNA and RNA molecules containing segments of the larger sequence are also provided for use in carrying out preferred aspects of the invention relating to the production of such peptides by the techniques of genetic engineering and the production of oligonucleotide probes.

Because the DNA sequence encoding the Flt4 protein is identified herein, DNA encoding the Flt4 protein may be produced by, e.g., polymerase chain reaction or by synthetic chemistry using commercially available equipment, after which the gene may be inserted into any of the many available DNA vectors using known techniques of recombinant DNA technology. Furthermore, automated equipment is also available that makes direct synthesis of any of the peptides disclosed herein readily available.

The present invention also is directed to Flt4 peptides and other constructs and to the use of Flt4 as a specific marker for lymphatic endothelial cells.

In a specific embodiment, the invention is directed to nucleic acid probes and antibodies recognizing Flt4, especially to monoclonal antibodies, and compositions containing such antibodies.

Also in a specific embodiment, the invention is directed to a method for monitoring lymphatic vessels in tissue samples and in organisms. Further, is it an object of the present invention to provide clinical detection methods describing the state of lymphatic tissue, and especially lymphatic vessels (inflammation, infection, traumas, growth, etc.), and to provide methods for detecting lymphatic vessels, and thus lymphatic vascularization, in an organism.

It is a further object of the present invention to provide monoclonal antibodies which specifically recognize the Flt4 receptor protein or various epitopes thereof. It is an object of the invention to use these monoclonal antibodies for diagnostic purposes for detecting and measuring the amount of Flt4 receptors in tissues, especially in lymphatic tissues.

Another aspect of the present invention relates to a method of determining the presence of Flt4-receptors in a

cell sample, comprising the steps of: (a) exposing a cell sample to an antibody, especially a monoclonal antibody, of the present invention; and (b) detecting the binding of said monoclonal antibody to Flt4 receptors.

The invention is further directed to a method of modulating (e.g., antagonizing or augmenting) the function of Flt4 in lymphatic vascularization and in inflammatory, infectious and immunological conditions. For example, in one embodiment, such a method comprises inhibiting the Flt4-mediated lymphatic vascularization by providing amounts of a Flt4-binding compound sufficient to block the Flt4 endothelial cell sites participating in such reaction, especially where Flt4 function is associated with a disease such as metastatic cancers, lymphomas, inflammation (chronic or acute), infections and immunological diseases.

The invention is further directed to a specific Flt4-stimulating ligand and monoclonal antibodies and their use for stimulating lymphatic endothelia and fragments and peptides as well as antibodies derived from research on the ligand to inhibit Flt4 function when desirable, such as in various disease states involving Flt4 function.

The invention provides a cell line source for the ligand of the Flt4 receptor tyrosine kinase. Using the conditioned medium from these cells, the Flt4 ligand may be purified and cloned by using methods standard in the art. Using this conditioned medium or the purified ligand, an assay system for Flt4 ligand and dimerization inhibitors as well as inhibitors of Flt4 signal transduction are obtained, which allow for identification and preparation of such inhibitors.

In a preferred embodiment of the invention, conditioned medium from the PC-3 cell line comprises a protein or a fragment thereof, which is capable of stimulating the Flt4 receptor and regulating the growth and differentiation as well as the differentiated functions of certain endothelial cells. The Flt4 ligand or its peptides or derivatives are useful in the regulation of endothelial cell growth, differentiation and their differentiated functions and in the generation of agonists and antagonists for the ligand. Particularly, the Flt4 ligand is useful in regulating lymphatic endothelia. However, the Flt4 ligand, when purified, or produced from a recombinant source, may also stimulate the related KDR/Flk-1 receptor.

The identification of Flt4-stimulating ligand makes it directly possible to assay for inhibitors of this ligand or inhibitors of Flt4 function. Such inhibitors are simply added to the conditioned media containing the Flt4 ligand and if they inhibit autophosphorylation, they act as Flt4 signalling inhibitors. For example, recombinant or synthetic peptides (including but not limited to fragments of the Flt4 extracellular domain) may be assayed for inhibition of Flt4-ligand interaction or Flt4 dimerization. Such putative inhibitors of Flt4 and, in addition, antibodies against the Flt4 ligand, peptides or other compounds blocking Flt4 receptor-ligand interaction, as well as antisense oligonucleotides complementary to the sequence of mRNA encoding the Flt4 ligand are useful in the regulation of endothelial cells and in the treatment of diseases associated with endothelial cell function.

A detailed characterization of the Flt4 ligand, designated VEGF-C, is provided in PCT Patent Application No. PCT/US98/01973, filed Feb. 2, 1998, and published as International Publication No. WO 98/33917; in PCT Patent Application PCT/EP96/00427, filed Aug. 1, 1996, and published as International Publication WO 97/05250; and in the U.S. Patent Application priority documents relied upon therein for priority, all of which are incorporated herein by refer-

ence. The deduced amino acid sequence for prepro-VEGF-C is set forth herein in SEQ ID NO: 21.

A detailed description of a second Flt4 ligand, designated VEGF-D, is provided in Achen, et al., *Proc. Nat'l Acad. Sci. U.S.A.*, 95(2): 548-553 (1998), and in Genbank Accession No. AJ000185, both of which are incorporated herein by reference. The deduced amino acid sequence for prepro-VEGF-D is set forth herein in SEQ ID NO: 22.

The invention also is directed to a method of treating a mammalian organism suffering from a disease characterized by expression of Flt4 tyrosine kinase (Flt4) in cells, comprising the step of administering to the mammalian organism a composition, the composition comprising a compound effective to inhibit the binding of an Flt4 ligand protein to Flt4 expressed in cells of the organism, thereby inhibiting Flt4 function. The disease may be diseases already mentioned above, such as diseases characterized by undesirable lymphatic vascularization. Additionally, it has been discovered that Flt4 expression also occurs in blood vessel vasculature associated with at least some breast cancers, and possibly other cancers (i.e., at a level greatly exceeding the barely detectable or undetectable levels of expression in blood vessel vasculature of corresponding normal (healthy) tissue). Thus, in a preferred embodiment, the cells comprise endothelial cells (lymphatic or vascular). In another embodiment, the cells comprise neoplastic cells such as certain lymphomas that express Flt4. Treatment of humans is specifically contemplated.

By "compound effective to inhibit the binding of an Flt4 ligand protein to Flt4 expressed in cells of the organism" is meant any compound that inhibits the binding of the Flt4 ligand described herein as vascular endothelial growth factor C, as isolatable from PC-3 conditioned medium. It is contemplated that such compounds also will be effective for inhibiting the binding of vascular endothelial growth factor D to Flt4. Exemplary compounds include the following polypeptides: (a) a polypeptide comprising an antigen-binding fragment of an anti-Flt4 antibody; (b) a polypeptide comprising a soluble Flt4 fragment (e.g., an extracellular domain fragment), wherein the fragment and the polypeptide are capable of binding to an Flt4 ligand; (c) a polypeptide comprising a fragment or analog of a vertebrate vascular endothelial growth factor C (VEGF-C) polypeptide, wherein the polypeptide and the fragment or analog bind, but fail to activate, the Flt4 expressed on native host cells; and (d) a polypeptide comprising a fragment or analog of a vertebrate vascular endothelial growth factor-D (VEGF-D) polypeptide, wherein the polypeptide and the fragment or analog bind, but fail to activate, the Flt4 expressed on native host cells. Small molecule inhibitors identifiable by standard in vitro screening assays, e.g., using VEGF-C and recombinantly-expressed Flt4 also are contemplated. Polypeptides comprising an antigen-binding fragment of an anti-Flt4 antibody are highly preferred. Such polypeptides include, e.g., polyclonal and monoclonal antibodies that specifically bind Flt4; fragments of such antibodies; chimeric and humanized antibodies; bispecific antibodies that specifically bind to Flt4 and also specifically bind to another antigen, and the like.

In a preferred variation, the compound further comprises a detectable label as described elsewhere herein, or a cytotoxic agent. Exemplary cytotoxic agents include plant toxins (e.g., ricin, saporin), bacterial or fungal toxins, radioisotopes (e.g., 211-Astatine, 212-Bismuth, 90-Yttrium, 131-Iodine, 99 m-Technetium, and others described herein), anti-metabolite drugs (e.g., methotrexate, 5-fluorodeoxyuridine), alkylating agents (e.g., chlorambucil), anti-mitotic agents (e.g., vinca alkaloids), and DNA intercalating agents (e.g., adriamycin).

Likewise, to improve administration, the composition preferably further comprises a pharmaceutically acceptable diluent, adjuvant, or carrier medium.

As explained in detail herein, Flt4 expression, while largely restricted to the lymphatic endothelia of healthy adults, has been identified in the blood vasculature surrounding at least certain tumors. Thus, the invention further includes a method of treating a mammalian organism suffering from a neoplastic disease characterized by expression of Flt4 tyrosine kinase (Flt4) in vascular endothelial cells, comprising the steps of: administering to a mammalian organism in need of such treatment a composition, the composition comprising a compound effective to inhibit the binding of an Flt4 ligand protein to Flt4 expressed in vascular endothelial cells of the organism, thereby inhibiting Flt4-mediated proliferation of the vascular endothelial cells. Treatment of neoplastic diseases selected from carcinomas (e.g., breast carcinomas), squamous cell carcinomas, lymphomas, melanomas, and sarcomas, are specifically contemplated. However, it will be readily apparent that the screening techniques described herein in detail will identify other tumors characterized by Flt4 expression in vascular endothelial cells, which tumors are candidates susceptible to the anti-Flt4 treatment regimens described herein. Treatment of breast carcinomas characterized by expression of Flt4 in vascular endothelial cells is specifically contemplated. By neoplastic disease characterized by expression of Flt4 tyrosine kinase in vascular endothelial cells is meant a disease wherein Flt4 is identifiable in blood vasculature at a much higher level than the undetectable or barely detectable levels normally observed in the blood vascular of healthy tissue, as exemplified herein.

Therapeutically effective amounts of compounds are empirically determined using art-recognized dose-escalation and dose-response assays. By therapeutically effective for treatment of tumors is meant an amount effective to reduce tumor growth, or an amount effective to stop tumor growth, or an amount effective to shrink or eliminate tumors altogether, without unacceptable levels of side effects for patients undergoing cancer therapy. Where the compound comprises an antibody or other polypeptide, doses on the order of 0.1 to 100 mg antibody per kilogram body weight, and more preferably 1 to 10 mg/kg, are specifically contemplated. For humanized antibodies, which typically exhibit a long circulating half-life, dosing at intervals ranging from daily to every other month, and more preferably every week, or every other week, or every third week, are specifically contemplated. Monitoring the progression of the therapy, patient side effects, and circulating antibody levels will provide additional guidance for an optimal dosing regimen. Data from published and ongoing clinical trials for other antibody-based cancer therapeutics (e.g., anti-HER2, anti-EGF receptor) also provide useful dosing regimen guidance.

For therapeutic methods described herein, preferred compounds include polypeptides comprising an antigen-binding fragment of an anti-Flt4 antibody, and polypeptides comprising a soluble Flt4 extracellular domain fragment. Human and humanized anti-Flt4 antibodies are highly preferred.

An expected advantage of the therapeutic methods of the invention lies in the fact that Flt4 is normally not expressed at any significant level in the blood vasculature of healthy tissues. In a highly preferred embodiment, the therapeutic compound comprises a bispecific antibody, or fragment thereof, wherein the antibody or fragment specifically binds Flt4 and specifically binds a blood vascular endothelial marker antigen. By "blood vascular endothelial marker

antigen" is meant any cell surface antigen that is expressed on proliferating vascular endothelial cells, and, preferably, that is not expressed on lymphatic endothelial cells. Exemplary blood vascular endothelial markers include PAL-E [deWaal, et al., *Am. J. Pathol.*, 150:1951-1957 (1994)], VEGFR-1 and VEGFR-2 [Ferrara et al., *Endocrine Reviews*, 18:4-25 (1997)], Tie [Partanen et al., *Mol. Cell. Biol.*, 12:1698-1707 (1992)], endoglin [U.S. Pat. No. 5,776,427, incorporated herein by reference in its entirety], and von Willebrandt Factor. Such bispecific antibodies are expected to preferentially locate to the tumor-associated vasculature that expresses both Flt4 and the blood vascular endothelial marker. In a highly preferred embodiment, the compound further comprises an anti-neoplastic or cytotoxic agent conjugated to the bispecific antibody, for the purposes of killing the tumor cells and/or killing the vasculature supply to the tumor cells. Exemplary agents include those described above, and also therapeutic proteins, such as statins, cytokines, chemokines, and the like, to stimulate an immune response to the tumor in the host.

In an alternative embodiment, the compound comprises an antibody (or bispecific antibody) that recognizes an epitope (or epitopes) comprised of an Flt4/Flt4 ligand complex (e.g., a complex comprised of Flt4 bound to VEGF-C or VEGF-D).

It is further contemplated that the therapeutic compound will be conjugated or co-administered with broad spectrum agents that have potential to inhibit angiogenic factors. Such agents include, e.g., heparin binding drugs (such as pentosan and suramin analogs) that may inhibit angiogenic factors that bind heparin; and chemical agents that block endothelial cell growth and migration, such as fumagillin analogs.

Conjugation of the anti-Flt4 compound to a prodrug that would be targeted to tumor vessels by the anti-Flt4 compound and then activated (e.g., by irradiation) locally at sites of tumor growth also is contemplated. Use of such prodrug strategy has the expected advantage of minimizing side effects of the drug upon healthy lymphatic vessels that express Flt4.

Similarly, the invention includes a method of treating a mammalian organism suffering from a neoplastic disease characterized by expression of Flt4 tyrosine kinase (Flt4) in vascular endothelial cells, comprising the steps of: identifying a mammalian organism suffering from a neoplastic disease state characterized by expression of Flt4 in vascular endothelial cells, and administering to the mammalian organism in need of such treatment a composition, the composition comprising a compound effective to inhibit the binding of an Flt4 ligand protein to Flt4 expressed in vascular endothelial cells of the organism, thereby inhibiting Flt4-mediated proliferation of the vascular endothelial cells.

The invention also provides a method for screening a biological sample for the presence of Flt4 receptor tyrosine kinase protein (Flt4), comprising the steps of: (a) contacting a biological sample suspected of containing Flt4 with a composition comprising an Flt4 binding compound, under conditions wherein the compound will bind to Flt4 in the biological sample; (b) washing the biological sample under conditions that will remove Flt4 binding compound that is not bound to Flt4 in the sample; and (c) screening the sample for the presence of Flt4 by detecting Flt4 binding compound bound to Flt4 receptor tyrosine kinase in the sample after the washing step. Preferably, the compound comprises a polypeptide selected from the group consisting of: (a) a polypeptide comprising an antigen-binding fragment of an anti-Flt4 antibody; and (b) a polypeptide comprising an Flt4

ligand or Flt4 binding fragment thereof. Antibodies that specifically bind Flt4, and that further comprise a detectable label, are highly preferred.

The invention also is directed to a method for imaging vertebrate tissue suspected of containing cells that express Flt4 receptor tyrosine kinase protein (Flt4), comprising the steps of: (a) contacting vertebrate tissue with a composition comprising an Flt4 binding compound; and (b) imaging the tissue by detecting the Flt4 binding compound bound to the tissue. Preferably, the tissue is human tissue, and the method further comprises the step of washing the tissue, after the contacting step and before the imaging step, under conditions that remove from the tissue Flt4 compound that is not bound to Flt4 in the tissue.

The invention is further directed to a method of screening for a neoplastic disease state, comprising the steps of: (a) contacting tissue from a mammalian organism suspected of having a neoplastic disease state with a composition comprising an antibody or antibody fragment that specifically binds Flt4 receptor tyrosine kinase; (b) detecting the antibody or antibody fragment bound to cells in the mammalian organism; and (c) screening for a neoplastic disease from the quantity or distribution of the antibody bound to cells in the mammalian organism. As described herein, Flt4 (which usually is undetectable or barely detectable in the blood vasculature) is strongly stained in the blood vasculature of at least some tumors. Thus, in one embodiment, in the screening step, the detection of the antibody or antibody fragment bound to blood vessel endothelial cells is correlated with the presence of a neoplastic disease. In this method, it will be understood that "detection" means detection at a level significantly higher than the barely detectable or undetectable levels that would occur in corresponding normal (healthy) tissue, as described herein. Such differential expression can be confirmed by comparison to a control performed with tissue from a healthy organism. Screening mammary tissue for neoplasms is specifically contemplated.

BRIEF DESCRIPTION OF THE DRAWINGS

The patent or application file contains at least one drawing executed in color. Copies of this patent or patent application publication with color drawing(s) will be provided by the Office upon request and payment of the necessary fee.

FIG. 1A is a schematic depiction of the structure of Flt4 cDNA clones;

FIG. 1B is a photographic reproduction of a Northern hybridization gel;

FIGS. 2A-F present a schematic depiction of structural features of Flt4 and a comparison with the Flt1 tyrosine kinase sequence;

FIG. 3A is a schematic depiction of the 3' ends of the cDNA inserts of clones J.1.1 and I.1.1;

FIG. 3B is a photographic reproduction of autoradiograms of hybridizations with anti-sense RNA probe and the long and short forms of Flt4 RNA;

FIG. 3C is a photographic reproduction of autoradiograms of hybridizations with anti-sense RNA probe and the long and short forms of Flt4 RNA;

FIG. 4 is a photographic reproduction of a gel illustrating a hybridization analysis of Flt4 sequences in DNA samples from different species;

FIGS. 5A-5H depict immunohistochemical characterization of VEGFR-3-expressing vessels in intraductal carcinoma. In adjacent sections (FIGS. 5A, B), VEGFR-3 and PAL-E decorate a similar pattern of "necklace" vessels

(arrowheads) around the duct filled with carcinoma cells. Another set of adjacent sections was compared with staining for VEGFR-3 (FIG. 5C), laminin (FIG. 5D), collagen XVIII (FIG. 5E) and SMA (FIG. 5F). Double staining for PAL-E and VEGFR-3 (FIG. 5G) and comparison with adjacent section stained for VEGFR-3 only (FIG. 5H). The vessels adjacent to the affected ducts are double-positive (arrowheads), whereas a VEGFR-3 positive vessel is present a short distance away from the affected duct in the inter-ductal stroma (arrows). Note that basal lamina is positive for PAL-E in the double staining procedure. Magnifications: FIGS. 5A,B 400 x. FIGS. 5C, D, E, F 320 x. FIGS. 5E,F 480 x.

DETAILED DESCRIPTION

The cloning, sequencing and expression of a novel receptor tyrosine kinase, termed Flt4, is described below. The Flt4 gene maps to chromosomal region 5q35 where many growth factors and growth factor receptors are located. The extracellular domain of Flt4 consists of seven immunoglobulin-like loops including twelve potential glycosylation sites. On the basis of structural similarities, Flt4 and the previously known Flt1 and KDR/FLK1 receptors may constitute a subfamily of class III tyrosine kinases. The Flt4 gene is expressed as 5.8 kb and 4.5 kb mRNAs which were found to differ in their 3' sequences and to be differentially expressed in HEL and DAMI leukemia cells.

A Wilm's tumor cell line, a retinoblastoma cell line, and a nondifferentiated teratocarcinoma cell line expressed Flt4; whereas differentiated teratocarcinoma cells were negative. Most fetal tissues also expressed the Flt4 mRNA, with spleen, brain intermediate zone and lung showing the highest levels. In human adult tissues the highest expression level was found in placenta, lung, kidney, heart and liver in decreasing order of expression. In situ hybridization, the Flt4 autoradiographic grains decorated endothelial cells of fetal lung. Immunohistochemical staining of Flt4 in fetal tissues confirmed staining of the endothelial cells. The expression pattern of Flt4 in comparison to Flt1 and KDR differs greatly in tissues of 18-week-old human fetuses. See Kaipainen et al., *J. Exp. Med.*, 178:2077 (1993).

Expression vectors containing the Flt4 cDNA have been produced and expressed in COS and NIH3T3 cells as described in Examples 4 and 11.

The Flt4 DNAs and polypeptides of the invention may be useful in the purification of the Flt4 ligand, and in the regulation of growth and differentiation of endothelial cells in various organs. They may also prove valuable in the diagnosis/treatment of certain diseases.

In the description that follows, a number of terms used in recombinant DNA (rDNA) technology are extensively utilized. In order to provide a clear and consistent understanding of the specification and claims, including the scope to be given to such terms, the following definitions are provided.

Gene. A DNA sequence containing a template for a RNA polymerase. The RNA transcribed from a gene may or may not code for a protein. RNA that codes for a protein is termed messenger RNA (mRNA) and, in eukaryotes, is transcribed by RNA polymerase II. However, it is also known to construct a gene containing a RNA polymerase II template wherein a RNA sequence is transcribed which has a sequence complementary to that of a specific mRNA but is not normally translated. Such a gene construct is herein termed an "antisense RNA gene" and such a RNA transcript is termed an "antisense RNA." Antisense RNAs are not normally translatable due to the presence of translational stop codons in the antisense RNA sequence.

A "complementary DNA" or "cDNA" gene includes recombinant genes synthesized by reverse transcription of mRNA lacking intervening sequences (introns).

Cloning vehicle. A plasmid or phage DNA or other DNA sequence which is able to replicate autonomously in a host cell, and which is characterized by one or a small number of endonuclease recognition sites at which such DNA sequences may be cut in a determinable fashion without loss of an essential biological function of the vehicle, and into which DNA may be spliced in order to bring about its replication and cloning. The cloning vehicle may further contain a marker suitable for use in the identification of cells transformed with the cloning vehicle. Markers, for example, are tetracycline resistance or ampicillin resistance. The word "vector" is sometimes used for "cloning vehicle."

Expression vector. A vehicle or vector similar to a cloning vehicle and which is capable of expressing a gene which has been cloned into it, after transformation into a host. The cloned gene is usually placed under the control of (i.e., operably linked to) certain control sequences such as promoter sequences. Expression control sequences vary depending on whether the vector is designed to express the operably linked gene in a prokaryotic or eukaryotic host and may additionally contain transcriptional elements such as enhancer elements, termination sequences, tissue-specificity elements, and/or translational initiation and termination sites.

The present invention pertains to both expression of recombinant Flt4 proteins (short and long forms), and to the functional derivatives of these proteins.

Functional Derivative. A "functional derivative" of Flt4 proteins is a protein which possesses a biological activity (either functional or structural) that is substantially similar to a biological activity of non-recombinant Flt4 proteins. A functional derivative of the Flt4 protein may or may not contain post-translational modifications such as covalently linked carbohydrate, depending on the necessity of such modifications for the performance of a specific function. The term "functional derivative" is intended to include the "fragments," "variants," "analogues," and "chemical derivatives" of a molecule.

As used herein, a molecule is said to be a "chemical derivative" of another molecule when it contains additional chemical moieties not normally a part of the molecule. Such moieties may improve the molecule's solubility, absorption, biological half-life, etc. The moieties may alternatively decrease the toxicity of the molecule and eliminate or attenuate any undesirable side effect of the molecule, etc. Moieties capable of mediating such effects are disclosed in Remington's Pharmaceutical Sciences (1980). Procedure for coupling such moieties to a molecule are well known in the art.

Fragment. A "fragment" of a molecule such as Flt4 protein is meant to refer to any portion of the molecule, such as the peptide core, or a variant of the peptide core.

Variant. A "variant" of a molecule such as Flt4 protein is meant to refer to a molecule substantially similar in structure and biological activity to either the entire molecule, or to a fragment thereof. Thus, provided that two molecules possess a similar activity, they are considered variants as that term is used herein even if the composition or secondary, tertiary, or quaternary structure of one of the molecules is not identical to that found in the other, or if the sequence of amino acid residues is not identical.

Analogue. An "analogue" of Flt4 protein or genetic sequence is meant to refer to a protein or genetic sequence

substantially similar in function to the Flt4 protein or genetic sequence herein.

DESCRIPTION OF PREFERRED EMBODIMENTS

The present invention is directed to what applicants have termed "Flt4," a receptor for tyrosine kinase, Flt4-encoding nucleic acid molecules (e.g. cDNAs, genomic DNAs, RNAs, anti-sense RNAs, etc.), production of Flt4 peptides or Flt4 protein from a Flt4 gene sequence and its product, recombinant Flt4 expression vectors, Flt4 analogues and derivatives, and diagnostic and/or therapeutic uses of Flt4 and related proteins, Flt4 ligands, Flt4 antagonists and anti-Flt4 antibodies.

PRODUCTION OF RECOMBINANT Flt4

Biologically active Flt4 may be produced by the cloning and expression of the Flt4-encoding sequence or its functional equivalent in a suitable host cell.

Production of Flt4 using recombinant DNA technology may be divided into a step-wise process for the purpose of description: (1) isolating or generating the coding sequence (gene) for the desired Flt4; (2) constructing an expression vector capable of directing the synthesis of the desired Flt4; (3) transfecting or transforming appropriate host cells capable of replicating and expressing the Flt4 gene and/or processing the gene product to produce the desired Flt4; and (4) identifying and purifying the desired Flt4 product.

ISOLATION OR GENERATION OF THE Flt4 GENE

The nucleotide coding sequence of Flt4 or functional equivalents thereof, may be used to construct recombinant expression vectors which will direct the expression of the desired Flt4 product. In the practice of the method of the invention, the nucleotide sequence depicted therein, or fragments or functional equivalents thereof, may be used to generate the recombinant molecules which will direct the expression of the recombinant Flt4 product in appropriate host cells. Flt4-encoding nucleotide sequences may be obtained from a variety of cell sources which produce Flt4-like activities and/or which express Flt4-encoding mRNA. Applicants have identified a number of suitable human cell sources for Flt4, including human placenta, leukemia cells and some tumor cell lines.

The Flt4 coding sequence may be obtained by cDNA cloning from RNA isolated and purified from such cell sources or by genomic cloning. The Flt4 sequence may be for example amplified by polymerase chain reaction from cDNA or genomic DNA material using techniques well known in the art. Either cDNA or genomic libraries of clones may be prepared using techniques well known in the art and may be screened for particular Flt4 DNAs with nucleotide probes which are substantially complementary to any portion of the Flt4 gene. Full length clones, i.e., those containing the entire coding region of the desired Flt4, may be selected for constructing expression vectors. Alternatively, Flt4 encoding DNAs may be synthesized in whole or in part by chemical synthesis using techniques standard in the art. Due to the inherent degeneracy of nucleotide coding sequences, other DNA sequences which encode substantially the same or a functionally equivalent amino acid sequence may be used in the practice of the method of the invention. Such alterations of Flt4 nucleotide sequences include deletions, additions or substitutions of different nucleotides resulting in a sequence that encodes the same or

15

a functionally equivalent gene product. The gene product may contain deletions, additions or substitutions of amino acid residues within the sequence which result in silent changes thus producing a bioactive product. Such amino acid substitutions may be made on the basis of similarity in polarity, charge, solubility, hydrophobicity, hydrophilicity and/or the amphipathic nature of the residues involved. For example, negatively charged amino acids include aspartic acid and glutamic acid; positively charged amino acids include lysine and arginine; amino acids with uncharged polar head groups or nonpolar head groups having similar hydrophilicity values include the following: leucine, isoleucine, valine; glycine, alanine; asparagine, glutamine; serine, threonine; phenylalanine, tyrosine.

CONSTRUCTION OF Flt4 EXPRESSION VECTORS

Using this information, a variety of recombinant DNA vectors capable of providing the Flt4 receptor tyrosine kinase in reasonable quantities are provided. Additional recombinant DNA vectors of related structure that code for synthetic proteins having the key structural features identified herein as well as for proteins of the same family from other sources can be produced from the Flt4 receptor tyrosine kinase cDNA using standard techniques of recombinant DNA technology. A transformant expressing the Flt4 receptor tyrosine kinase has been produced as an example of this technology (see EXAMPLES 3 and 4). The newly discovered sequence and structure information can be used, through transfection of eukaryotic cells, to prepare the Flt4 receptor tyrosine kinase and its various domains for biological purposes.

IDENTIFICATION OF TRANSFECTANTS OR TRANSFORMANTS EXPRESSING Flt4 GENE PRODUCTS

The host cells which contain the recombinant coding sequence and which express the biologically active, mature product may be identified by at least four general approaches: (a) DNA-DNA, DNA-RNA or RNA-antisense RNA hybridization; (b) the presence or absence of "marker" gene functions; (c) assessing the level of transcription as measured by the expression of Flt4 mRNA transcripts in the host cell; and (d) detection of the mature gene product as measured by immunoassay and, ultimately, by its biological activities.

In the first approach, the presence of Flt4 coding sequences inserted into expression vectors may be detected by DNA-DNA hybridization using probes comprising nucleotide sequences that are homologous to the Flt4 coding sequence.

In the second approach, the recombinant expression vector/host system may be identified and selected based upon the presence or absence of certain "marker" gene functions (e.g., thymidine kinase activity, resistance to antibiotics, resistance to methotrexate, transformation phenotype, occlusion body formation in baculovirus, etc.). For example, if the Flt4 coding sequence is inserted within a marker gene sequence of the vector, recombinants containing that coding sequence can be identified by the absence of the marker gene function. Alternatively, a marker gene can be placed in tandem with the Flt4 sequence under the control of the same or different promoter used to control the expression of the Flt4 coding sequence. Expression of the marker in response to induction or selection indicates expression of the Flt4 coding sequence.

16

In the third approach, transcriptional activity for the Flt4 coding region may be assessed by hybridization assays. For example, polyadenylated RNA can be isolated and analyzed by Northern blotting using a probe homologous to the Flt4 coding sequence or particular portions thereof. Alternatively, total nucleic acids of the host cell may be extracted and assayed for hybridization to such probes.

In the fourth approach, the expression of Flt4 can be assessed immunologically, for example by Western blots, immunoassays such as radioimmunoprecipitation, enzyme-linked immunoassays and the like. The ultimate test of the success of the expression system, however, involves the detection of the biologically active Flt4 gene product. Where the host cell secretes the gene product, the cell free media obtained from the cultured transfectant host cell may be assayed for Flt4 activity. Where the gene product is not secreted, cell lysates may be assayed for such activity. In either case, assays which measure ligand binding to Flt4 or other bioactivities of Flt4 may be used.

Flt4 DERIVATIVES, ANALOGUES AND PEPTIDES

The production and use of derivatives, analogues, and peptides related to Flt4 are also envisioned and are within the scope of the invention. Such derivatives, analogues, or peptides may have enhanced or diminished biological activities in comparison to native Flt4, depending on the particular application. Flt4 related derivatives, analogues, and peptides of the invention may be produced by a variety of means known in the art. Procedures and manipulations at the genetic and protein levels are within the scope of the invention. Peptide synthesis, which is standard in the art, may be used to obtain Flt4 peptides. At the protein level, numerous chemical modifications may be used to produce Flt4-like derivatives, analogues, or peptides by techniques known in the art, including but not limited to specific chemical cleavage by endopeptidases (e.g. cyanogen bromides, trypsin, chymotrypsin, V8 protease, and the like) or exopeptidases, acetylation, formylation, oxidation, etc.

Preferred derivatives, analogs, and peptides are those which retain Flt4 ligand binding activity. Those derivatives, analogs, and peptides which bind Flt4 ligand but do not transduce a signal in response thereto are useful as Flt4 inhibitors. Those derivatives, analogs, and peptides which bind Flt4 ligand and transduce a signal in response thereto, e.g., through a process involving intracellular Flt4 autophosphorylation, are useful in the same manner as native Flt4. A preferred Flt4 ligand for use in such binding and/or autophosphorylation assays is a ligand comprising an approximately 23 kd polypeptide that is isolatable from a PC-3 conditioned medium as described herein. This ligand, designated Vascular Endothelial Growth Factor-C (VEGF-C), has been characterized in detail in PCT Patent Application PCT/FI96/00427, filed Aug. 1, 1996, and published as International Publication WO 97/05250, and in the U.S. Patent Application priority documents relied upon therein for priority, all of which are incorporated herein by reference in their entirety.

ANTI-Flt4 ANTIBODIES

Also within the scope of the invention is the production of polyclonal and monoclonal antibodies which recognize Flt4 or related proteins.

Various procedures known in the art may be used for the production of polyclonal antibodies to epitopes of Flt4. For the production of antibodies, various host animals (including

but not limited to rabbits, mice, rats, etc.) can be immunized by injection with Flt4, or a synthetic Flt4 peptide. Various adjuvants may be used to increase the immunological response, depending on the host species, including but not limited to Freund's (complete and incomplete) adjuvant, mineral gels such as aluminum hydroxide, surface active substances such as lysolecithin, pluronic polyols, polyanions, oil emulsions, keyhole limpet hemocyanins, dinitrophenol, and potentially useful human adjuvants such as BCG (*Bacillus Calmette-Guerin*) and *Corynebacterium parvum*.

A monoclonal antibody to an epitope of Flt4 may be prepared by using any technique which provides for the production of antibody molecules by continuous cell lines in culture. These include but are not limited to the hybridoma technique originally described by Köhler et al., *Nature*, 256: 495-497 (1975), and the more recent human B-cell hybridoma technique [Kosbor et al., *Immunology Today*, 4: 72 (1983)] and the EBV-hybridoma technique [Cole et al., *Monoclonal Antibodies and Cancer Therapy*, Alan R Liss, Inc., pp. 77-96 (1985)]. Antibodies against Flt4 also may be produced in bacteria from cloned immunoglobulin cDNAs. With the use of the recombinant phage antibody system it may be possible to quickly produce and select antibodies in bacterial cultures and to genetically manipulate their structure.

Antibody fragments which contain the idiotype of the molecule may be generated by known techniques. For example, such fragments include but are not limited to: the F(ab')₂ fragment which may be produced by pepsin digestion of the antibody molecule; the Fab' fragments which may be generated by reducing the disulfide bridges of the F(ab')₂ fragment, and the two Fab fragments which may be generated by treating the antibody molecule with papain and a reducing agent.

Antibodies to Flt4 may be used in the qualitative and quantitative detection of mature Flt4 and Flt4 precursor and subcomponent forms, in the affinity purification of Flt4 polypeptides, and in the elucidation of Flt4 biosynthesis, metabolism and function. Detection of Flt4 tyrosine kinase activity may be used as an enzymatic means of generating and amplifying a Flt4 specific signal in such assays.

Antibodies to Flt4 may also be useful as diagnostic and therapeutic agents.

USES OF Flt4, Flt4-ENCODING NUCLEIC ACID MOLECULES, AND ANTI-Flt4 ANTIBODIES

Applicants envision a wide variety of uses for the compositions of the present invention, including diagnostic and/or therapeutic uses of Flt4, Flt4 analogues and derivatives, Flt4-encoding nucleic acid molecules, antisense nucleic acid molecules and anti-Flt4 antibodies.

Flt4-encoding nucleic acid molecules or fragments thereof may be used as probes to detect and quantify mRNAs encoding Flt4. Assays which utilize nucleic acid probes to detect sequences comprising all or part of a known gene sequence are well known in the art. Flt4 mRNA levels may indicate emerging and/or existing neoplasias as well as the onset and/or progression of other human diseases. Therefore, assays which can detect and quantify Flt4 mRNA may provide a valuable diagnostic tool.

Anti-sense Flt4 RNA molecules are useful therapeutically to inhibit the translation of Flt4-encoding mRNAs where the therapeutic objective involves a desire to eliminate the presence of Flt4 or to downregulate its levels. Flt4 anti-sense RNA, for example, could be useful as a Flt4 antagonizing

agent in the treatment of diseases in which Flt4 is involved as a causative agent, for example due to its overexpression.

Additionally, Flt4 anti-sense RNAs are useful in elucidating Flt4 functional mechanisms. Flt4-encoding nucleic acid molecules may be used for the production of recombinant Flt4 proteins and related molecules as separately discussed in this application.

Anti-Flt4 antibodies may be used to diagnose and quantify Flt4 in various contexts. For example, antibodies against various domains of Flt4 may be used as a basis for Flt4 immunoassays or immunohistochemical assessment of Flt4. Tyrosine kinase activity of Flt4 may be useful in these assays as an enzymatic amplification reaction for the generation of a Flt4 signal. Anti-Flt4 antibodies may also be useful in studying the amount of Flt4 on cell surfaces.

Antibodies may be produced which function as Flt4 ligand agonists or antagonists whereby the regulation of Flt4 activity becomes possible. Also, random peptides may be produced by synthetic means or by recombinant means from random oligonucleotides and the ones showing specific binding to the Flt4 receptor may be selected with the aid of the Flt4 extracellular domain. Such peptide segments also may be selected from a phage display library using the extracellular domain of Flt4, using methods standard in the art. Such peptides may have agonistic or antagonistic activity. Flt4 antibodies may also provide valuable diagnostic tools after conjugation to various compounds for in vivo imaging of Flt4 expressing cells and tissues or tumors.

Monoclonal antibodies against Flt4 may be coupled either covalently or noncovalently to a suitable supramagnetic, paramagnetic, electron-dense, echogenic or radioactive agent to produce a targeted imaging agent. Antibody fragments generated by proteolysis or chemical treatments or molecules produced by using the epitope binding domains of the monoclonal antibodies could be substituted for the intact antibody. This imaging agent would then serve as a contrast reagent for X-ray, magnetic resonance, sonographic or scintigraphic imaging of the human body for diagnostic purposes.

MOLECULAR BIOLOGY OF Flt4

The complete sequences of the Flt4 cDNA clones set forth in SEQ ID NOs: 1 and 3 extend for 4195 or 4795 nucleotides and contain open reading frames of 1298 or 1363 amino acids, depending on alternative splicing. The nucleotide and deduced Flt4 amino acid sequence (short form) is shown in SEQ ID NOs: 1 and 2. FIG. 2 depicts a comparison of the Flt4 amino acid sequence with that of the Flt1 tyrosine kinase amino acid sequence. See Shibuya et al., *Oncogene*, 5: 519-524 (1990).

A putative signal peptide sequence of mostly hydrophobic amino acids follows the initiator methionine. The sequence surrounding the corresponding ATG is in agreement with the consensus translation initiation sequence [Kozak, *Nucl. Acids Res.*, 15: 8125-8135 (1987)]. The predicted extracellular portion of both Flt4 polypeptides is 775 amino acids long and contains twelve potential sites for asparagine-linked glycosylation (NXS/T). It also contains several amino acid residues exhibiting a pattern of spacing described for members of the immunoglobulin superfamily of proteins [Williams et al., *Annu. Rev. Immunol.*, 6: 381-405 (1988)]. It has 12 cysteine residues and it can be organized in seven immunoglobulin-like domains. The predicted Ig-like domain IV lacks cysteine residues. FIG. 2 also shows the extracellular domain of Flt1 (SEQ. ID No. 5), which is the closest human homologue of Flt4. From this figure one can

see the alignment of the cysteine residues and the very similar composition of the Ig-like regions.

The cytoplasmic domain of Flt4 is separated from the extracellular part by a putative transmembrane region of 23 hydrophobic amino acid residues. This sequence is flanked on the cytoplasmic side by a basic region suggesting the junction between the transmembrane and cytoplasmic domains. The tyrosine kinase homologous domain begins at residue 843 and includes an ATP-binding pocket and a putative autophosphorylation site homologous to Y416 of c-src at Y1068 (FIG. 2). The tyrosine kinase catalytic domain of Flt4 is divided into two subdomains by a 65 amino acid sequence (aa 944–1008) which is mostly hydrophilic and does not show homology to Flt1. Unlike Flt1, Flt4 does not contain tyrosine residues in its kinase insert.

A second species of Flt4 mRNA has an alternative 3' end which encodes a longer form of the Flt4 protein.

In FIGS. 3A–C, production of short and long forms of the Flt4 mRNA by alternative splicing is illustrated. FIG. 3A shows the schematic structure of the 3' ends of the cDNA inserts of clones J.1.1 and I.1.1. The TAG stop codon of clone J.1.1 as well as the polyadenylation site (polyA) are indicated. Clone I.1.1 differs from clone J.1.1 in the shaded segment (the long and short forms of Flt4 mRNA, respectively). TAA and polyA indicate the stop codon and polyadenylation site of clone I.1.1. In addition, the restriction endonuclease cleavage sites for EcoRI and Aval have been indicated. Shown below is the 256 bp EcoRI-Aval insert of clone I.1.1. used for cRNA protection analysis. The most heavily-shaded segment indicates sequences from the polylinker in the linearized sense RNA template for transcription of the antisense strand in vitro. Also shown are the schematic structures of the protected fragments after RNase protection analysis. FIGS. 3B and 3C, show autoradiograms of the 256 bp ³⁵S-labeled antisense RNA probe and the 211 and 124 bp digested fragments representing the long and short forms of Flt4 RNA when protected by polyadenylated RNA from the indicated cell lines (Tera-2 is a teratocarcinoma cell line, which has been analyzed here with or without retinoic acid (RA) treatment for 10 days.) The (negative) control lane shows results of protection with transfer RNA. Note the downregulation of Flt4 mRNAs during the differentiation of the Tera-2 cells. Tera-2 cells of clone 13 were provided by Dr. C. F. Graham (Department of Zoology, University of Oxford, UK). Cells between passages 1840 were used in this study. The cells were maintained in Eagle's minimum essential medium (MEM) supplemented with 10% fetal calf serum and antibiotics. To induce differentiation, the cells were plated on gelatin-coated tissue-culture grade dishes at a density of 1.5×10⁵ cells/cm². On the following day, 2×10⁻⁶ M RA was added to the medium. The cells were cultured in the presence of RA for up to 10 days.

Results shown in FIGS. 3A–C illustrate the generation of carboxy termini of these two Flt4 (short and long) forms generated by alternative splicing.

According to its deduced amino acid sequence, Flt4 belongs to class III RTKs. More specifically, Flt4 belongs to a subfamily of RTKs, which contain seven Ig-loops in their extracellular part and thus it differs from other members of class III RTKs which contain five Ig-loops. Flt4 is most closely homologous with the prototype receptor of the FLT family, Flt1, which was cloned as a v-ros-related DNA from a human genomic DNA library [Shibuya et al., *Oncogene*, 5: 519–524 (1990)] and with the mouse FLK1 receptor, which was cloned from hematopoietic stem cell-enriched fractions

of mouse liver [Matthews et al., *Cell*, 65: 1143–1152 (1991); Matthews et al., *Proc. Natl. Acad. Sci. USA*, 88: 9026–9030 (1991)]. The extracellular domain of Flt4 shows 33% and 37% amino acid sequence identity with human Flt1, and mouse FLK1, respectively. Flt1 and FLK1, like Flt4, are widely expressed in various normal tissues, such as lung, heart, and kidney. In addition, a recently identified human endothelial cell receptor tyrosine kinase KDR [Terman et al., *Oncogene*, 6: 1677–1683 (1991)] shows considerable homology with Flt4 and Flt1 family members. From the available sequence data one may calculate that KDR is 81% identical with Flt4 in the tyrosine kinase (TK) domain. In addition, the extracellular domain of KDR also has a seven Ig-loop structure and its TK1 and TK2 domains are 95% and 97% identical with the corresponding domains of mouse FLK1 receptor. This suggests that KDR is the human homologue of mouse FLK1.

While the Flt4 TK domain is about 80% identical with the TK domains of Flt1 and FLK1/KDR, it is only about 60% identical with the TK domains of other receptors of the RTK class III. As these other receptors also have only five Ig-like domains in the extracellular region, one can classify Flt4, Flt1 and FLK1/KDR in a separate FLT subfamily within class III RTKs.

The tyrosine residue located in the sequence D/E-D/E-Y-M/V-P/D/E-M [Cantley, et al., *Cell*, 64: 281–302 (1991)] (SEQ. ID NO. 6) in kinase inserts of PDGFRs, c-fms and c-kit is an autophosphorylation site, which, when phosphorylated, binds the SH2 domain of phosphatidylinositol 3'-kinase (PI-3K) [Reedijk et al., *EMBO J.*, 11: 1365–1372 (1992)]. Interestingly, unlike these class III RTKs, members of the FLT subfamily or the Flt3/FLK2 receptor do not contain such consensus motifs.

The eight human class III RTK genes are clustered in three different chromosomes. Chromosome 4 contains the c-kit, PDGFR- α and KDR genes [Yarden et al., *ELBO J.*, 6: 3341–3351 (1987); Stenman et al., *Genes, Chromosomes, Cancer*, 1: 155–158 (1989); Terman et al., *Oncogene*, 6: 1677–1683 (1991)]. The Flt1 and Flt3 genes are located in chromosome 13q12 [Sato et al., *Jpn. J. Cancer Res.*, 78: 772–775 (1987); Rosnet et al., *Genomics*, 9: 380–385 (1991)], while Flt4 is localized in chromosome 5 band q35 [Aprelikova et al., *Cancer Res.*, 52: 746–748 (1992)]; close to the fms and PDGFR- β genes [Warrington et al., *Genomics*, 11: 701–708 (1991)]. The long arm of chromosome 5 is involved in translocations found in leukemia cells. Deletions of part of the long arm of chromosome 5 were found in the bone marrow cells of patients with refractory anemia and macrocytosis [Van Den Berghe et al., *Nature*, 251: 437–439 (1974)]. An abnormal 5q chromosome is found in a few other myeloproliferative diseases, such as refractory anemia with excess blasts [Swolin et al., *Blood*, 58: 986–993 (1981)], agnogenic myeloid metaplasia [Whang-Peng et al., *Leuk. Res.*, 2: 41–48 (1978)], chronic myelogenous leukemia [Tomiyasu et al., *Cancer Genet. Cytogenet.*, 2: 309–315 (1980)], polycythemia vera [Van Den Berghe et al., *Cancer Genet. Cytogenet.*, 1: 157–162 (1979)] and essential thrombocythemia [Nowell et al., *Cancer*, 42: 2254–2260 (1978)].

The findings on Flt4 mRNA expression suggest that its protein product is characteristic for certain leukemia cells. Several differentiation antigens shared between megakaryoblastic and endothelial cells have been shown to exist, one example being the platelet glycoprotein IIIa [Ylänne et al., *Blood*, 72: 1478–1486 (1988); Kieffer et al., *Blood*, 72: 1209–1215 (1988); Berridge et al., *Blood*, 66: 76–85 (1985)]. In addition, Flt4 is expressed by certain endothelial cells of, e.g., the lung and kidney during the fetal period.

To further understand the role of Flt4 during development, partial cDNAs for mouse Flt4 were cloned. Using these probes in in situ hybridization, Flt4 mRNA expression during mouse development was analyzed. It was determined that Flt4 is expressed during vasculogenesis and angiogenesis of the lymphatic system. The relevance of these findings was also confirmed in normal and pathological human adult tissues, as Flt4 was found in lymphatic endothelial cells of human adult tissues both in normal and pathological conditions, as well as in some high endothelial venules (HEVs).

The cloning of mouse Flt4 cDNA fragments showed that their deduced amino acid sequence is almost identical with the corresponding human sequence (amino acid identity about 96% in both segments studied). Further evidence for the identity of the mouse Flt4 cDNA was obtained from Northern hybridization studies, wherein probes from both species yielded the typical 5.8 kb mRNA signal from mouse tissues. Analysis of RNA isolated from various tissues of adult mice showed Flt4 expression in the liver, lung, heart, spleen and kidney, with no or very little hybridization in the brain and testes. This pattern is similar to the pattern reported earlier by Galland et al., *Oncogene*, 8: 1233 (1993). The results of RNase protection suggested that the Flt4 gene is needed during mouse development, starting from 8.5 day p.c. embryos, and the relative expression levels appeared quite stable.

For the in situ hybridization, two fragments of mouse Flt4 cDNA were selected which encode sequences of the extracellular domain. This allowed a clear distinction of the hybridization pattern from the related FLK-1 and Flt1 receptor patterns, which show only a very low degree of sequence identity with Flt4 in the extracellular region. See Millauer et al., *Cell*, 72: 835 (1993); Yamaguchi et al., *Development*, 118:489 (1993); Peters et al., *Proc. Natl. Acad. Sci. USA*, 90: 8915 (1993); Finnerty et al., *Oncogene*, 8: 2293 (1993).

Flt4, similar to the FLK-1, Flt1, Tie and Tek endothelial receptor tyrosine kinase genes, was not expressed in 7.5 day post-coitum (p.c.) embryos. In a 8.5-day p.c. embryo, the strongest Flt4 signals were localized in the allantois, the angioblasts of head mesenchyme, the dorsal aortae, and the cardinal vein. Weak signals were seen in the endocardium. In contrast, angioblasts of the yolk sac were negative, unlike for FLK-1 and Flt1, Tie and Tek. See Korhonen et al., *Oncogene*, 8:395 (1993); and Peters et al., *Proc. Natl. Acad. Sci. USA*, 90: 8915 (1993). The restriction of Flt4 expression to the venous system was even more clear in samples from 11.5 day mouse embryos, where the Tie mRNA was expressed also in arteries. In 12.5-day p.c. embryos the Flt4 signal decorated developing venous and presumptive lymphatic endothelia, but unlike for the endothelial Tie receptor tyrosine kinase, arterial endothelia were negative. During later stages of development, Flt4 mRNA became restricted to vascular plexuses devoid of blood cells, representing developing lymphatic vessels. Only the lymphatic endothelium and some high endothelial venules expressed Flt4 mRNA in adult human tissues. Increased expression occurred in lymphatic sinuses and high endothelial venules, in metastatic lymph nodes, and in lymphangioma.

Due to difficulties in the interpretation of data from mouse embryos, human endothelia were studied, because the lymphatic system is much better defined in humans. Also, cells established from various endothelia could be studied in cell culture to see if the specificity of Flt4 expression persists in in vitro conditions. Endothelial cells lines are known to lose differentiated features upon in vitro culture. Therefore, it was not unexpected that they were negative for Flt4 mRNA.

Cultured aortic endothelial cells were also devoid of Flt4 mRNA. However, signals were obtained from human endothelial cells grown from the microvasculature and from femoral and umbilical veins. Thus, at least some of the specificity of Flt4 expression was retained in cell culture.

In situ hybridization analysis of adult human tissues confirmed the restriction of Flt4 to the lymphatic system seen in the developing mouse embryos. Flt4 expression was seen in the lymphatic endothelia and in the sinuses of human lymph nodes. Interestingly, also some of the HEVs, which have a cuboidal endothelium, shown to function in the trafficking of leukocytes to the lymph nodes, were Flt4-positive. Furthermore, a parallel hybridization analysis showed that Flt4 mRNA levels were enhanced in these structures in metastatic as compared to normal lymph nodes. Flt4 was also very prominent in lymphangiomas, which are benign tumours composed of connective tissue stroma and growing, endothelial-lined lymphatic channels. Flt4 mRNA was restricted to the lymphatic endothelium of these tumors and absent from their arteries, veins and capillaries. In the human lung, lymphatic structures were the only Flt4-positive vessels identified.

The foregoing results indicate that Flt4 is a novel marker for lymphatic vessels and some high endothelial venules in human adult tissues. The results also support the theory on the venous origin of lymphatic vessels. Flt4, as a growth factor receptor, may be involved in the differentiation and functions of these vessels. A detailed characterization of biological effects mediated through Flt4 via the Flt4 ligand, VEGF-C, is provided in PCT Patent Application PCT/EP96/00427, filed Aug. 1, 1996, and published as International Publication WO 97/05250.

These results, combined with the Flt4-binding compounds according to the present invention, allows a selective labeling of lymphatic endothelium, especially by using antibodies of the present invention coupled to radioactive, electron-dense or other reporter substances, which can be visualized. It may be possible to inject into the lymphatic system substances, containing Flt4 receptor internalization-inducing monoclonal antibodies or ligands, and thereby transport predefined molecules into the lymphatic endothelium. Also, it may be possible to use Flt4-binding compounds according to the invention for the detection of high endothelial venules, especially activated HEVs, which express enhanced levels of the Flt4 receptor. To our knowledge, no such specific markers are currently available for lymphatic endothelium.

The following examples are given merely to illustrate the present invention and not in any way to limit its scope.

EXAMPLE 1

Isolation and Characterization of cDNA Clones Encoding Flt4

MATERIALS AND METHODS

An oligo-dT primed human HEL cell cDNA library in bacteriophage lambda gt11 [A kind gift from Dr. Mortimer Poncz, Childrens Hospital of Philadelphia, Pa.; Poncz et al., *Blood*, 69: 219-223 (1987)] was screened with a cDNA fragment PCR-amplified from the same library [Aprelikova et al., *Cancer Res.*, 52: 746-748 (1992)]. Positive plaques were identified and purified as described [Sambrook et al., *Molecular Cloning—A Laboratory Manual*, Cold Spring Harbor Laboratory Press, (1989)]. cDNA inserts of bacteriophage lambda were isolated as EcoRI fragments and

subcloned into a GEM3Zf(+) plasmid (Promega). The entire Flt4 protein coding region was isolated. Three overlapping clones isolated from the HEL-library (as illustrated in FIG. 1) were sequenced using the dideoxy chain termination method with oligonucleotide primers designed according to the sequences obtained. All portions of the cDNAs were sequenced on both strands. Sequence analyses were performed using the GCG package programs [Devereux et al., *Nucleic Acids Res.*, 12: 387-395 (1984) and the Prosite program for Apple Macintosh].

FIG. 1A illustrates a schematic structure of the Flt4 cDNA clones analyzed. Arrows delineate subcloned restriction fragments (whose sizes are shown in kb) used for probing Northern blots depicted in FIG. 1B. E=EcoRI site, S=SphI site. FIG. 1B illustrates Northern hybridization analysis of DAMI and HEL leukemia cell RNAs with the probes shown in FIG. 1A.

RESULTS

A 210 bp long Flt4 cDNA fragment isolated by a PCR cloning method from a HEL cell cDNA library was used as a molecular probe to screen an oligo-dT-primed human erythroleukemia cell cDNA library.

Nucleotide sequence analysis of clones revealed an open reading frame of 1298 amino acid (aa) residues (SEQ ID NO: 2, FIG. 2). The translational initiator methionine marked in the figure is surrounded by a typical consensus sequence [Kozak, *Nucleic Acids Res.*, 12: 857-872 (1984)] and followed by a hydrophobic amino acid sequence characteristic of signal sequences for translocation into the endoplasmic reticulum.

The extracellular domain of Flt4 can be aligned into seven immunoglobulin-like loops (FIG. 2). The figure also shows the comparison of Flt4 with Flt1, which contains very similar structures. The amino acid sequence of Flt1 is set forth as SEQ. ID NO: 5.

Amino acid residues 775-798 form a hydrophobic stretch of sequence, which is likely to function as the transmembrane domain of the receptor, followed by several basic residues on the putative cytoplasmic side of the polypeptide. The juxtamembrane domain is 44 residues long before the beginning of a tyrosine kinase sequence homology at aa 842. With the interruption of homology in the kinase insert sequence of 65 aa, this homology is first lost at 1175 aa at carboxyl terminal tail of the receptor. A search for related tyrosine kinase domains in the amino acid sequence database (Swissprot and NBRF) identifies the Flt1 and PDGFRB tyrosine kinases with homology of about 80 and 60% in the catalytic tyrosine kinase regions respectively.

EXAMPLE 2

Preparation of an anti-Flt4 Antisera

A 657 base pair EcoRI fragment encoding the predicted C-terminus of Flt4 short form was cloned in-frame with the glutathione-S-transferase coding region in the pGEX-III bacterial expression vector (Pharmacia) to produce a GST-Flt4 fusion protein in *E. coli*. The resulting fusion protein was produced in bacteria and partially purified by glutathione affinity chromatography according to the manufacturer's instructions. This protein was used in immunization of rabbits in order to produce polyclonal antibodies against Flt4. Antisera were used after the third booster immunization.

EXAMPLE 3

Expression of Flt4 in COS Cells

MATERIALS AND METHODS

The full-length Flt4 protein coding sequence (combined from three clones, FIG. 1) was inserted into the HindIII-

BamHI site of SVpoly mammalian expression vector [Stacey et al., *Nucleic Acids Res.*, 18: 2829 (1990)] construct SV14-2. The expression vectors (SV-FLT4 short and SV-FLT4 long, containing the respective forms of Flt4 cDNA) were introduced into COS cells by DEAE-dextran transfection method [McCutchan et al., *J. Natl. Cancer Inst.*, 41: 351-357 (1968)]. Two days after transfection, the cells were washed with phosphate-buffered saline (PBS) and scraped into immunoprecipitation buffer (10 mM Tris pH 7.5, 50 mM NaCl, 0.5% sodium deoxycholate, 0.5% Nonidet P40, 0.1% SDS, 0.1 TIU/ml Aprotinin). The lysates were sonicated, centrifuged for 15' at 10,000xg and incubated overnight on ice with 3 ml of the antisera. Protein A sepharose (Pharmacia) was added and the incubation was continued for 30' with rotation. The precipitates were washed four times with the immunoprecipitation buffer, once with PBS and once with aqua before analysis in SDS-PAGE.

RESULTS

The structural predictions of the Flt4 cDNA sequence were tested by cloning the full-length Flt4 short and long protein-coding regions into the HindIII-BamHI sites of the pSVpoly expression vector and transfecting these expression vectors into COS cells. The proteins produced by these two constructs differ in their C-terminus: the longer form contains an additional 65 amino acids. Two days after transfection, the cells were lysed and immunoprecipitated using antibodies generated against the GST-Flt4 fusion protein containing 40 carboxyl terminal amino acid residues of the short form of the predicted Flt4 protein (i.e., a portion common to both the short and long forms of Flt4). Immunoprecipitated polypeptides were analyzed by SDS-polyacrylamide gel electrophoresis. The preimmune serum did not reveal any specific bands, whereas the Flt4-specific antibodies recognize two bands of about 170 and 190 KD. These two bands may represent differentially glycosylated forms of Flt4 protein.

EXAMPLE 4

Expression of Flt4 in NIH3T3 Cells

MATERIALS AND METHODS

The full-length Flt4 cDNA (short form) was subcloned into the LTRpoly vector (see Makela, et al., *Gene*, 118:293-294 (1992), disclosing plasmid vector pLTRpoly, having ATCC accession number 77109 and GeneBank accession number X60280) containing the Moloney murine leukemia virus long terminal repeat promoter. This LTR-FLT4 expression vector was used with pSV2neo marker plasmid to co-transfect NIH3T3 cells, and G418 resistant clones were analyzed for Flt4 expression.

For Western immunoblotting analyses, cells on one confluent large plate were lysed in 2.5% SDS, 125 mM Tris, pH 6.5. Cell lysates were electrophoresed on SDS-page and electroblotted onto a nitrocellulose membrane. The membrane was incubated with the antiserum raised against the Flt4 carboxy-terminus peptide, and bound antibodies were visualized using horseradish peroxidase conjugated swine anti-rabbit antiserum (Dako) and ECL reagents (Amersham). For metabolic labeling, the cultures were labeled with 100 μ Ci/ml 35 S-methionine for one hour. After labelling, cells were washed twice and incubated in their growth medium for 1 or 2 hours, lysed, immunoprecipitated with anti-Flt4 antibodies, and analyzed by SDS-PAGE and autofluorography.

25

RESULTS

The 170 and 190 KD polypeptides could be detected in the Flt4 short form-transfected into NIH3T3 cells, but not in cells transfected with pSV2neo only. In addition to these two bands, a major band of about 120 Kd was observed in the clones producing Flt4. Metabolic labeling and pulse-chase experiments showed that this protein is generated as a result of post-translational processing of the short form Flt4 polypeptides.

EXAMPLE 5

Chromosomal Mapping of the Flt4 Locus

Because some clustering of class III receptor genes has been observed, it is of great interest to determine the chromosomal localization of Flt4. Thus, rodent-human cell hybrids were analyzed, indicating linkage of Flt4 to human chromosome 5.

Localization of the Flt4 gene in the region 5q33→5qter was determined using hybrids carrying partial chromosome 5s. These hybrids were tested for presence of the Flt4 locus by filter hybridization. The region of chromosome 5 common to Flt4-positive hybrids and absent from the Flt4-negative hybrids was 5q33.1-qter. The presence of human chromosome 5q33-qter in the hybrids is thus correlated with the presence of Flt4 sequences. The regional mapping results indicated that the Flt4 locus is telomeric to the CSF1R/platelet-derived growth factor receptor- β (PDGFRB) locus as well as to the β -adrenergic receptor (ADRB) locus since these loci are all present in the hybrid GB 13, which was negative for Flt4.

EXAMPLE 6

Expression of the Flt4 mRNA in Tumor Cell Lines and Endothelial Cells

The leukemia cell lines (K562) used in this study have been reported in several previous publications; [Lozzio et al., *Blood*, 45: 321-334 (1975)], HL-60 [Collins et al., *Nature*, 270: 347-349 (1977)], HEL [Martin et al., *Science*, 216: 1233-1235 (1982)], DAMI [Greenberg et al., *Blood*, 72: 1968-1977 (1988)], MOLT4 [Minowada et al., *J. Natl. Cancer Inst.*, 49: 891-895 (1972)], Jurkat [Schwenk et al., *Blut*, 31: 299-306 (1975)], U937 [Sundström et al., *Int. J. Cancer*, 17: 565-577 (1976)], KG-1 [Koeffler et al., *Science*, 200: 1153-1154 (1978)], JOK-1 [Andersson et al., 1982, in R. F. Revoltella (ed.), *Expression of Differentiated Functions in Cancer Cells*, 239-245, Raven Press, New York] and ML-2 [Gahmberg et al., 1985, in L. C. Andersson, et al. (ed.), *Gene Expression During Normal and Malignant Differentiation*, 107-123, Academic Press, London]. The following tumor cell lines, obtained from the American Type Culture Collection also were analyzed: JEG-3, a choriocarcinoma; A204, a rhabdomyosarcoma; SK-NEP-1, a neuroblastoma; BT474, a breast carcinoma; Y79, a retinoblastoma. The leukemia cells were grown in RPMI containing 10% fetal calf serum (FCS) and antibiotics. Dami cells were cultivated in Iscove's modified DMEM with 10% horse serum. A permanent endothelial hybrid cell line (EAhy926) obtained by fusing first-passage human umbilical vein endothelial cells with the A549 lung carcinoma cells [Edgell et al., *Proc. Natl. Acad. Sci. USA*, 50: 3734-3737 (1983)] was cultured in DMEM-HAT medium containing 10% FCS and antibiotics.

Poly(A)⁺ RNA was extracted from the cell lines as described [Sambrook et al., see above]. 5 μ g of the Poly(A)⁺

26

RNA samples were electrophoresed in agarose gels containing formaldehyde and blotted using standard conditions [Sambrook et al., see above]. The inserts of the Flt4 cDNA clones were labelled by the random priming method and hybridized to the blots. Hybridization was carried out in 50% formamide, 5 \times Denhardt's solution (100 \times Denhardt's solution is 2% each of Ficoll, polyvinylpyrrolidone and bovine serum albumin), 5 \times SSPE (3M NaCl, 200 mM NaH₂PO₄, H₂O, 20 mM EDTA, pH 7.0), 0.1% SDS (sodium dodecyl sulphate), and 0.1 mg/ml of sonicated salmon sperm DNA at 42° C. for 18-24 h. The filters were washed at 65° C. in 1 \times SSC (150 mM NaCl, 15 mM sodium citrate, pH 7.0), 0.1% SDS and exposed to Kodak XAR-5 film.

Northern analyses were performed with the extracted poly(A)⁺ RNA from eight leukemia cell lines (HEL, K562, DAMI, U937, MOLT4, HL60, Jurkat, and KG-1) and the endothelial hybrid cell line (EAhy926). Hybridization with the GAPDH probe was used as an internal control for the loading of even amounts of RNA to the analysis. Only the HEL erythroleukemia cells, and DAMI megakaryoblastic leukemia cells expressed 5.8 kb and 4.5 kb Flt4 mRNA. The K562 erythroleukemia, Jurkat and MOLT-4 T-cell leukemias, as well as HL-60 promyelocytic leukemia, U937 monocytic leukemia, and KG-1 myeloid leukemia cells were negative for the Flt4 mRNA.

Northern analyses were performed with the extracted poly(A)⁺ RNA from five tumor cell lines (JEG-3, A-204, SK-NEP-1, BT-474, and Y79) and two of the aforementioned leukemia cell lines (JOK-1, MOLT4). The labeled S2.5 cDNA clone (see FIG. 1) was used as the hybridization probe. Hybridization with a β -actin probe was used as an internal control for the loading of even amounts of RNA to the analysis. Only the SK-NEP-1 neuroblastoma and Y79 retinoblastoma cells were observed to contain Flt4 transcripts.

Tera-2 teratocarcinoma cells were analyzed after a 10 day treatment with vehicle (-) or retinoic acid (+) to induce neuronal differentiation [Thompson et al., *J. Cell Sci.*, 72: 37-64 (1984)]. In Northern blotting analysis of poly(A)⁺ RNA isolated from the cells it was found that the undifferentiated cells expressed 5.8 kb and 4.7 kb mRNAs for Flt4, but after the 10 day differentiation, no Flt4 mRNA could be detected in Northern blotting and hybridization. These results indicate that Flt4 was downregulated during the differentiation of these cells.

Flt4 mRNA expression also was analyzed in undifferentiated and TPA-differentiated HEL cells. Both the HEL and DAMI cell lines possess a dual erythroid/megakaryoblastic phenotype and can be induced to further expression of megakaryoblastic markers by treatment with the tumor promoter 12-O-tetradecanoylphorbol-13-acetate (TPA). We analyzed whether Flt4 expression is stimulated in these cells during their differentiation. HEL cells were analyzed 2 days after treatment with TPA or with DMSO used to dissolve it. After stripping off the Flt4 signal, the filter was probed with Rb-1 and β -actin cDNAs to confirm an even loading of the lanes. On the basis of densitometric scanning analysis of the autoradiograph and normalization against the constitutive expression of the GAPDH gene, it was determined that the Flt4 mRNA level was increased about 3.4 fold in TPA-induced HEL cells, when the cells undergo megakaryoblastic differentiation.

EXAMPLE 7

Expression of Flt4 in Fetal Lung

In situ hybridization: Lung tissue from a 15 week-old human fetus was obtained with the permission of joint

ethical committee of the University Central Hospital and the University of Turku, Finland. The sample was fixed in 10% formalin for 18 hours at 4° C., dehydrated, embedded in wax, and cut into 6 μ m sections. The RNA probes of 206 and 157 bases (antisense and sense) were generated from linearized plasmid DNA using SP6 and T7 polymerases and [³⁵S]-UTP. In situ hybridization of sections was performed according to Wilkinson et al., *Development*, 99:493-500 (1987); Wilkinson, *Cell*, 50:79-88 (1987), with the following modifications: 1) instead of toluene, xylene was used before embedding in paraffin wax; 2) 6 μ m sections were cut, placed on a layer of diethyl pyrocarbonate-treated water on the surface of glass slides pretreated with 2% 3-aminopropyl-triethoxysilane (Sigma); 3) alkaline hydrolysis of the probes was omitted; 4) the hybridization mixture contained 60% deionized formamide; 5) the high stringency wash was for 80 minutes at 65° C. in a solution containing 50 mM DTT and 1xSSC; 6) the sections were covered with NTB-2 emulsion (Kodak) and stored at 4° C. After an exposure time of 14 days, the slides were developed for 2.5 minutes in a Kodak D-19 developer and fixed for 5 minutes with Unifix (Kodak). The sections were stained with hematoxylin in water.

In the hybridization studies using the anti-sense probe, Flt4 mRNA was observed mainly in certain endothelial cells of the lungs of a 15 week old fetus. Control hybridizations with the sense strand probe and with RNase A-treated sections did not give a signal above background.

For immunoperoxidase staining, a 1:100 dilution of the anti-Flt4 antibody, peroxidase-conjugated swine anti-rabbit antibodies and methods standard in the art were used. Control stainings with preimmune serum or immunogen-blocked serum did not give a signal. Lung tissue from seventeen-week old human fetuses were analyzed, and the results were consistent with those of the mRNA in situ hybridization experiments: the endothelium of certain large vessels of the lung were stained positive with the rabbit anti-Flt4 antiserum.

EXAMPLE 8

Identification of Flt4 Genes in Non-human Mammalian Species

In FIG. 4 the results of an experiment examining the presence of Flt4 sequences in DNA from different species is shown. In order to reveal how well the Flt4 gene has been conserved in evolution, the 2.5 kb cDNA fragment (see FIG. 1) was hybridized to genomic DNAs purified from different animals and from yeast and digested with EcoRI. The hybridization solution comprised 50% formamide, 5xDenhardt's solution, (100xDenhardt's solution is 2% each of Ficoll, polyvinylpyrrolidone and bovine serum albumin), 5xsaline-sodium phosphate-EDTA (3M NaCl, 200 mM NaH₂PO₄-H₂O, 20 mM EDTA, pH 7.0), 0.1% sodium dodecyl sulfate, and 0.1 mg/ml sonicated salmon sperm DNA. Hybridization was performed at 42° C. for 24 hours. The filter was washed at 65° C. in 1xstandard saline citrate (150 mM NaCl, 15 mM sodium citrate, pH 7.0) and 0.1% sodium dodecyl sulfate and exposed to Kodak XAR-5 film. Specific bands were found in monkey, rat, mouse, dog, cow, rabbit, and chick DNAs, but the yeast DNA did not give a signal. The Flt4 cDNA has been isolated from quails. See Eichmann et al., *Gene*, 174(1): 3-8 (Sep. 26, 1996) and Genbank accession number X83287.

EXAMPLE 9

Flt4 Gene Expression in Adult Human Tissues

Flt4 mRNA expression in adult human tissues was analyzed using 2 μ g of poly(A)⁺ RNA from heart, brain,

placenta, lung, liver, skeletal muscle, kidney, and pancreas tissues (Multiple Tissue Northern Blot, Clontech Inc.) by hybridization with the Flt4 cDNA probe. Control hybridizations with probes for constitutively expressed genes showed an even loading of the lanes.

Hybridization of poly(A)⁺ RNA from various human tissues with the Flt4 cDNA fragment showed mRNA bands of 5.8 and 4.5 kb mobility and a weakly labeled band of 6.2 kb in placenta, lung, heart and kidney. Faint mRNA bands were seen in the liver and skeletal muscle, whereas the pancreas and brain appeared to contain very little if any Flt4 RNA.

EXAMPLE 10

Flt4 Expression in Human Fetal Tissues

To examine Flt4 mRNA expression in human fetal tissues, a Northern blot containing total RNA from the below-listed tissues of 16-19 week human fetuses was hybridized with the 1.9 kb Flt4 cDNA fragment (see FIG. 1) and the resulting autoradiograph was scanned with a densitometer. The results were normalized for the amount of RNA estimated from a UV picture of the corresponding ethidium bromide (EtBr) stained gel. The following symbols denote mRNA levels in an increasing order: -, +, ++, +++.

TABLE 1

Fetal tissue	mRNA
<u>Brain</u>	
Meninges	+
Cortical plate	++
Intermediate zone	+++
Ependymal zone	+
Cerebellum	++
Choroid plexus	+
Liver	+
Pancreas	+
Small intestine	-
Heart	+
Lung	+++
Kidney	++
Adrenal	++
Skin	++
Spleen	+++
Thymus	-

Analysis of human fetal tissues showed that all except the thymus and small intestine contain Flt4 transcripts. The highest expression levels were found in lung and spleen.

EXAMPLE 11

Flt4 Expression Vector

Full-length Flt4 cDNA (short form) was produced by a) ligation of a SphI-cleaved Flt4 PCR fragment [amplified from the S2.5 kb clone (see FIG. 1) using the primer oligonucleotides 5'-ACATGCATGC CACCATGCAG CGGGGCGCCG CGCTGTGCCT GCGACTGTGG CTCT-GCCTGG GACTCCTGGA-3'(SEQ. ID NO. 7) (forward) and 5'-ACATGCATGC CCCGCCGGT CATCC-3' (reverse)] (SEQ. ID NO. 8) to the 5' end of the S2.5 kb fragment, subcloned into the pSP73 vector 5 (Promega), using two SphI sites; b) ligation of a PCR fragment containing the last 138 bps amplified from the 0.6 kb EcoRI fragment (see FIG. 1) with the oligonucleotide primers 5'-CGGAATTCAC CAIGACCCCA AC-3'(SEQ. ID NO. 9) (forward) and 5'-CCATCGATGG ATCCTACCTG AAGC-

CGCTTT CTT-3' (SEQ. ID NO. 10) (reverse) to the 3' end of construct a) using the EcoRI and BamHI sites; c) ligation of a 1.2 kb EcoRI fragment in the EcoRI site of construct b); d) ligation of the resulting full length HindIII-BamHI fragment into the HindIII-BamHI cleaved SV-poly expression vector [Stacey et al., *Nucl. Acids Res.*, 18: 2829 (1990)].

EXAMPLE 12

Identification of an Flt4 Ligand

Conditioned media from the PC-3 prostatic adenocarcinoma cell line (ATCC CRL 1435) cultured for 7 days in F12 medium in the absence of fetal bovine serum (FBS) was cleared by centrifugation at 16,000×g for 20 minutes and screened for the ability to induce tyrosine phosphorylation of Flt4.

NIH3T3-cells recombinantly expressing Flt4 (see Example 13) were reseeded on 5 cm diameter cell culture dishes and grown to confluency in Dulbecco's modified minimal essential medium (DMEM) containing 10% fetal bovine serum and antibiotics. The confluent cells were washed twice in phosphate-buffered saline (PBS) and starved in DMEM/0.2% bovine serum albumin overnight. For stimulation, the starvation medium was replaced by 1 ml of the conditioned medium and the cells were incubated at 37° C. for 5 minutes.

After stimulation with the PC-3 conditioned medium, the culture plates containing the cells were put on ice and washed twice with Tris-HCl, pH 7.4, 150 mM NaCl containing 100 mM NaVO₄. The washing solution was removed from the dishes and the cells were lysed in RIPA buffer [10 mM Tris-HCl pH 7.5, 50 mM NaCl, 0.5% sodium deoxycholate, 0.5% Nonidet P40, 0.1% sodium dodecyl sulphate (SDS)] containing aprotinin, 1 mM PMSF and 1 mM NaVO₄, and the lysates were sonicated for 10 seconds twice. The lysates were then centrifuged at 16,000 ×g for 30 minutes and the supernatants were transferred to new tubes and used for immunoprecipitation.

The polyclonal antibodies against the Flt4 C-terminus (described above) were used for immunoprecipitation. Supernatants from the cell lysates were incubated for 2 hours on ice with 2 to 4 µl of rabbit polyclonal anti-Flt4 antiserum. About 30 µl of a 50% (vol/vol) solution of protein A-Sepharose (Pharmacia) in PBS was added, and incubation was continued for 45 minutes with rotation at 44° C. The immunoprecipitates were washed three times with the RIPA buffer and once with PBS.

The immunoprecipitates were then subjected to SDS-polyacrylamide gel electrophoresis (SDS-PAGE) in a 7.5% gel and blotted on nitrocellulose. These Western blots were incubated with monoclonal anti-phosphotyrosine (anti-P-Tyr) antibodies (1:2000 dilution of PT-66 Sigma, cat. P-3300) followed by detection with peroxidase-conjugated rabbit anti-mouse antibodies (1:1000 dilution, Dako, cat. P 161) using the chemiluminescence detection system (Amersham). In some cases, the blots were stripped to clear previous signals for 30 minutes at 50° C. in 100 mM 2-mercaptoethanol, 2% SDS, 62.5 mM Tris-HCl pH 6.7 with occasional agitation and re-stained with anti-Flt4 antibodies (1:1000 dilution) followed by staining with peroxidase-conjugated swine anti-rabbit antibodies (1:1000 dilution, Dako, P217). As a positive control for the tyrosine phosphorylation of Flt4, anti-Flt4 immunoprecipitates from the Flt4-expressing NIH3T3 cells treated with 100 mM of the tyrosyl phosphatase inhibitor sodium pervanadate (PerVO₄) for 20 minutes were used. Treatment of cells with Sodium

pervanadate was done by addition of 100 mM (final concentration) of sodium orthovanadate and 2 mM (final concentration) of hydrogen peroxide to the cell medium and incubation of the cells for 20 minutes at 37° C. 5% CO₂. That procedure resulted in the generation of the peroxidized form of vanadate (vanadyl hydroperoxide), which is a very potent inhibitor of the protein tyrosine phosphatases in living cells.

The PC-3 cell conditioned medium stimulated tyrosine phosphorylation of a 120 kD polypeptide which co-migrated with tyrosine phosphorylated, processed mature form of Flt4. Co-migration was confirmed after restaining of the blot with anti-Flt4 antibodies.

To prove that 120 kD polypeptide is not a non-specific component of the conditioned medium, 15 ml of conditioned medium were separated by SDS-PAGE, blotted on nitrocellulose, and the blot was stained with anti-P-Tyr antibodies. Several polypeptides were detected, but none of them comigrated with Flt4, indicating that the 120 kD band is indeed tyrosine-phosphorylated protein immunoprecipitated from the stimulated cells. Analysis of stimulation by PC-3 conditioned medium pretreated with heparin Sepharose CL-6B (Pharmacia) for 2 hours at room temperature (lane 3) shows that the Flt4 ligand does not bind to heparin.

Unconditioned medium did not induce Flt4 autophosphorylation. Also, neither non-transfected NIH3T3 cells nor NIH3T3 cells transfected with the FGFR-4 showed tyrosine phosphorylation of the 120 kD polypeptide upon stimulation with the conditioned medium from PC-3 cells. Stimulating activity was considerably increased when the PC-3 conditioned medium was concentrated fourfold using a Centricon-10 concentrator (Amicon). Also, the flow through obtained after the concentration, containing proteins of less than 10,000 molecular weight (<10,000), did not stimulate phosphorylation of Flt4. Pretreatment of the concentrated conditioned medium of PC-3 cells with 50 ml of the Flt4 extracellular domain (Flt4EC-6×His, see below) coupled to CNBr-activated Sepharose (1 mg/ml) according to the manufacturer's instructions completely abolished the tyrosine phosphorylation of Flt4. Analogous pretreatment of the conditioned medium with Sepharose CL-4B did not affect its stimulatory activity.

These data prove that PC-3 cells produce soluble ligand for Flt4. The above experiments prove that the ligand binds to the recombinant Flt4 EC domain. Thus, that ligand can be purified using the recombinant Flt4 EC domain in affinity chromatography. The purified protein can be electrophoresed in SDS-PAGE, blotted onto polyvinylidene difluoride (PVDF) membranes and its amino terminal sequence can be determined by methods standard in the art. Alternatively, the purified ligand can be digested to peptides for their amino terminal sequence determination. Peptide sequences obtained from the purified protein are used for the synthesis of a mixture of oligonucleotides encoding such sequences. Such oligonucleotides and their complementary DNA strand counterparts can be radioactively labelled by and used for the screening of cDNA libraries made from the PC-3 cells to obtain a cDNA encoding the ligand, all by methods standard in the art (Wen et al., *Cell* 69: 559-572 (1992)). Alternatively, such oligonucleotides and their counterparts can be used as primers in polymerase chain reaction (PCR) to amplify sequences encoding the ligand using cDNA made from PC-3 cell RNA as a template. Such method of cDNA synthesis and PCR (RT-PCR) is standard in the art (Innis et al., 1990, PCR protocols, Academic Press; McPherson, M. J. et al., 1991, PCR, a practical approach, IRL Press; Partanen

et al., *Proc. Nat. Acad. Sci., USA*, 87: 8913-8917 (1990)). Yet another alternative is to clone the Flt4 ligand from the PC-3 cells by using cDNAs cloned into eukaryotic expression vector (e.g. using the Invitrogen Librarian cloning kit and vectors provided, such as pcDNA I or pcDNA III) and screening of such libraries transfected into, e.g., COS cells with Flt4-alkaline phosphatase (Cheng and Flanagan, *Cell*, 79: 157-168, (1994)), Flt4-immunoglobulin (Flt4-Ig) (Lyman et al., *Cell*, 75: 1157-1167 (1993)), or similar affinity reagents, by methods standard in the art.

EXAMPLE 13

Cell Lines and Transfections

NIH3T3 cells and 293-EBNA cells (Invitrogen) were cultured in DMEM containing 10% FCS. For stable expression, NIH3T3 cells were transfected with the LTR-FLT41 vector together with the pSV-neo vector (see Example 4, above) where the Flt4 cDNA is expressed under the control of the Moloney murine leukemia virus LTR promoter, by the lipofection method using the DOTAP transfection reagent (Boehringer-Mannheim). COS-1 cells were transfected by the DEAE dextran method (McClutchan and Pagano, *J. Natl. Cancer Inst.*, 41: 351-35 (1968)). Transfected cells were selected in 500 mg/ml neomycin.

EXAMPLE 14

Construction and Expression of Flt4 Fusion Proteins

The pVTBac-FLT4EC-6xHis fusion construct. The ends of cDNA encoding Flt4 were modified as follows: The 3' end of Flt4 cDNA sequence encoding the extracellular domain (EC) was amplified using oligonucleotides 5'-CTGGA GTCGACTTGGCGGACT-3' (SEQ ID NO: 13, Sall site underlined, containing sequence corresponding to nucleotides 2184-2204 of SEQ ID NO: 1) and 5'-CGC GGATCCCTAGTGAAGGTG AIGGTGATGTCTACCTTCGATCATGCTGCCCTTAT CCTC-3' (SEQ ID NO: 14, BamHI site underlined, containing sequence complementary to nucleotides 2341-2324 of SEQ ID NO: 1) encoding 6 histidine residues for binding to a Ni-NTA column (Qiagen, Hilden, Germany) followed by a stop codon. The amplified fragment was digested with Sall and BamHI and ligated as a Sall-BamHI fragment into the LTR-FLT41 vector (see Example 4), replacing a unique Sall-BamHI fragment containing sequences encoding the Flt4 transmembrane and cytoplasmic domains.

The 5' end of the Flt4 cDNA without the Flt4 signal sequence encoding region was amplified by PCR using oligonucleotides 5'-CCC AAGCTTGGATCCAAGTGGCTACTCCATGACC-3' (SEQ ID NO: 11, HindIII and BamHI sites underlined, containing sequence corresponding to nucleotides 86-103 of SEQ ID NO: 1) and 5'-GTTGCCTGTGATGTGCACCA-3' (SEQ ID NO: 12, containing sequence complementary to nucleotides 700-681 of SEQ ID NO: 1). This amplified fragment (which included nucleotides 86-700 of SEQ ID NO: 1) was digested with HindIII and SphI (the SphI site, corresponding to nucleotides 588-593 of SEQ ID NO: 1, being within the amplified region of the Flt4 cDNA).

The resultant HindIII-SphI fragment was used to replace a HindIII-SphI fragment in the modified LTR-FLT41 vector described immediately above (the HindIII site is in the 5' junction of the Flt4 insert with the pLTRpoly portion of the vector, the SphI site is in the Flt4 cDNA and corresponds to

nucleotides 588-593 of SEQ ID NO: 1). The resultant Flt4EC-6xHis insert was then ligated as a BamHI fragment into the BamHI site in the pVTBac plasmid (Tessier et al., *Gene* 98: 177-183 (1991)). The construct was transfected together with baculovirus genomic DNA into SF-9 cells by lipofection. Recombinant virus was generated and used for infection of High-Five cells (Invitrogen).

The Flt4-AP fusion construct. The 3' end of the sequence encoding the Flt4 EC domain was amplified using oligonucleotides 5'-CTGGAGTCGACTTGGCGGACT-3' (SEQ ID NO: 15) and 5'-CGGGATCCCTCCATGCTGCCCTTATCCT-3' (SEQ ID NO: 16) and ligated as Sall-BamHI fragment into the LTR-FLT41 vector, replacing sequences encoding the transmembrane and cytoplasmic domains. The resulting insert was then ligated as a HindIII-BamHI fragment into the HindIII-BglII sites of plasmid APTag-1 in frame with the alkaline phosphatase coding region (Flanagan and Leder, 1990, *Cell* 63, 185-194). NIH3T3 cells were co-transfected with this Flt4-AP construct and pSV2neo (Southern and Berg, *J. Mol. Appl. Genet.* 1: 327-341 (1982)) by lipofection using the DOTAP transfection reagent (Boehringer) and the transfected cells were selected in the presence of 500 mg/ml neomycin. The recombinant protein produced into the medium was detected by a calorimetric reaction for staining for alkaline phosphatase activity (Cheng and Flanagan, *Cell* 79: 157-168 (1994)).

The Flt4-Ig construct. A recombinant DNA encoding an Flt4-immunoglobulin chimera was constructed as follows. The 5' end of the cDNA encoding Flt4, including Flt4 nucleotides encoding the signal sequence, was amplified by PCR using primers 5'-GGCAAGCTTGAATTCGCCACCATGCAGCGGGG CGCC-3' (SEQ ID NO: 17) and 5'-GTTGCCTGTGATGTGCACCA-3' (SEQ ID NO: 18) and ligated as HindIII-SphI fragment into the LTR-FLT41 vector. The 3' end of Flt4 EC-encoding sequence was amplified using oligonucleotides 5'-CTGGAGTCGACTTGGCGGACT-3' (SEQ ID NO: 19) and 5'-CGCGGATCCAAGCTTACTTACCTTCCATGCTGCCCTTATCCTCG-3' (SEQ ID NO: 20) and ligated as Sall-BamHI fragment into the LTR-FLT41 vector replacing the sequences encoding the transmembrane and cytoplasmic domains. This Flt4EC insert containing a splice donor site was ligated first into pHyCE2 containing exons encoding the human immunoglobulin heavy chain hinge and constant region exons (Kajalainen, K., *TIBTECH*, 9: 109-113 (1991)). The EcoRI-BamHI insert containing the Flt4-Ig chimera was then blunted by methods standard in the art (Klenow) and ligated to the blunted HindIII site in pREP7 (Invitrogen). The construct was transfected into 293-EBNA cells by the calcium-phosphate precipitation method and the conditioned medium was used for the isolation of the Flt4-Ig protein by protein A-Sepharose affinity chromatography.

EXAMPLES 15-17

Purification and Sequencing the Flt4 Ligand

Cell culture supernatants produced by PC-3 cells under serum-depleted conditions are concentrated 30-50 fold using Centrprep filter cartridges and loaded onto a column of immobilized Flt4 extracellular domain. Two affinity matrices are prepared using the alternative constructs and methods. In the first case the Flt4EC-6xHis fusion protein is crosslinked to CNBr-activated Sepharose 4B (Pharmacia) and in the second case the Flt4-Ig fusion protein is coupled to protein A-Sepharose using dimethylpimelidate (Schneider

et al., 1982, *J. Biol. Chem.* 257: 10766–10769). The material eluted from the affinity column is subjected to further purification using ion exchange and reverse-phase high pressure chromatography and SDS-polyacrylamide gel electrophoresis. Chromatography fractions are tested for the ability to stimulate tyrosine phosphorylation of Flt4. The purified biologically active ligand protein is microsequenced and the degenerate oligonucleotides are made based on the amino acid sequence obtained, for the purpose of isolating and cloning a ligand-encoding cDNA; e.g., from a cDNA library generated from poly(A)⁺ RNA isolated from PC-3 cells.

A detailed characterization of an Flt4 ligand, designated VEGF-C, is provided in PCT Patent Application No. PCT/US98/01973, filed Feb. 2, 1998 (published as International Publication No. WO 98/33917); in PCT Patent Application PCT/FI96/00427, filed Aug. 1, 1996 (published as International Publication WO 97/05250); and in the U.S. Patent Application priority documents relied upon therein for priority, all of which are incorporated herein by reference: U.S. patent application Ser. Nos. 08/510,193, filed Aug. 1, 1995; U.S. patent application Ser. No. 08/585,895, filed Jan. 12, 1996; U.S. patent application Ser. No. 08/601,132, filed Feb. 14, 1996; U.S. patent application Ser. No. 08/671,573, filed Jun. 28, 1996; and U.S. patent application Ser. No. 08/795,430, filed Feb. 5, 1997. It will be apparent from the foregoing teachings that the Flt4 ligand described herein may be used in assays as an additional indicia to confirm the identity of human Flt4 allelic variants, and to confirm that non-human gene sequences having homology to the Flt4 sequences taught herein (See, e.g., Example 8 and FIG. 4) are in fact the non-human counterparts to Flt4. The deduced amino acid sequence for prepro-VEGF-C is set forth herein in SEQ ID NO: 21.

A detailed description of a second Flt4 ligand, designated VEGF-D, is provided in Achen, et al., *Proc. Nat'l Acad. Sci. U.S.A.*, 95(2): 548–553 (1998), also incorporated herein by reference. The cDNA and deduced amino acid sequences for prepro-VEGF-D are set forth herein in SEQ ID NO: 22.

EXAMPLE 18

Cloning of Mouse Flt4 cDNA Probes

Approximately 10⁶ plaques from a λFIX@II genomic library from 129SV mice (Stratagene) was screened with the S2.5 human Flt4 receptor cDNA fragment described above, covering the extracellular domain. See also Pajusola et al., *Cancer Res.*, 52:5738 (1992). A 2.5 kb BamHI fragment was subcloned from a positive plaque and sequenced from both ends. From this subclone, polymerase chain reaction was used to amplify and clone into the pBluescript KSII+/– vector (Stratagene) an exon fragment covering nucleotides 1745–2049 of the mouse Flt4 cDNA sequence. See Finnerty et al., *Oncogene*, 8:2293 (1993).

A second fragment covering nucleotides 1–192 was similarly cloned.

EXAMPLE 19

Analysis of Flt4 mRNA in Mouse Tissues

Total RNA was isolated from developing embryos (8–18 days p.c. and one day old mice) according to Chomczynski et al., *Anal. Biochem.*, 162:156 (1987). The sample from 8 day p.c. embryos also included the placenta.

For RNase protection analysis, RNA probe was generated from the linearized murine Flt4 plasmid obtained according

to Example 18 using [³²P]-UTP and T7 polymerase for the antisense orientation. The β-actin probe used corresponds to nucleotides 1188–1279 of the published mouse β-actin sequence. See Tokunaga, et al., *Nucleic. Acid. Res.*, 14:2829 (1986). After purification in a 6% polyacrylamide/7M urea gel, the labelled transcripts were hybridized to 30 μg of total RNA overnight at 52° C. Unhybridized RNA was digested with RNase A (10 U/ml) and T1 (1 mg/ml) at 37° C., pH 7.5 for 1 hour. The RNases were inactivated by proteinase K digestion at 37° C. for 15 minutes and the samples were analysed in a 6% polyacrylamide/7M urea gel.

The pattern of expression of Flt4 analysed in this experiment showed that very weak mRNA signals were obtained from lung, liver, heart, kidney, skeletal muscle and spleen, whereas testis and brain were apparently without specific signal. Analysis of a series of RNAs collected during different phases of mouse development by RNase protection assay showed that the Flt4 mRNA was expressed throughout embryogenesis from day 8 p.c. to newborn mice without great variations in signal intensity.

EXAMPLE 20

In Situ Hybridization for Flt4 in Mouse Embryos

To better assign Flt4 transcripts to cells and tissues, sections of 7.5 and 8.5 day p.c. mouse embryos were hybridized with labelled Flt4 RNAs. Mouse embryos were derived from matings of CBA and NMRI mice. Pregnant mice were killed by cervical dislocation and the embryos were either immediately frozen or transferred via phosphate buffered saline into 4% paraformaldehyde. The embryos and isolated mouse organs were fixed for 18 hours at 4° C., dehydrated, embedded in paraffin, and cut into 6 μm sections.

RNA probes (antisense and sense) of 192 and 305 nucleotides (see Example 18) were generated from linearized plasmids using [³⁵S]-UTP. In situ hybridization of sections was performed according to Wilkinson et al., *Development*, 99:493 (1987); and Wilkinson et al., *Cell*, 50:79 (1987), incorporated by reference herein, with the following modifications: 1) instead of toluene, xylene was used before embedding in paraffin wax; 2) 6 μm sections were cut, placed on a layer of diethyl pyrocarbonate-treated water on the surface of glass slides pretreated with 2% 3-triethoxysilylpropylamine; 3) alkaline hydrolysis of the probes was omitted; and 4) the high stringency wash was for 80 minutes at 65° C. in a solution containing 30 mM DTT and 1×SSC. The sections were covered with NTB-2 emulsion (Kodak) and stored at 4° C. The slides were exposed for 14 days, developed, and stained with hematoxylin. Control hybridizations with sense strand and RNase A-treated sections did not give a specific signal above background.

Flt4 mRNA expression was not detected in 7.5 day p.c. mouse embryos, but bright signals were detected in the developing aortae on day 8.5 of development. In contrast, the developing yolk sac was Flt4-negative. In the extraembryonic tissues, Flt4 was prominently expressed in the allantois, whereas developing blood islands of the yolk sac were negative. On the other hand, angioblasts of the head mesenchyme were strongly Flt4-positive. In the developing placenta, Flt4 signal was first seen in peripheral sinusoidal veins. In 9.5 day p.c. placenta, the endothelium of venous lacunae and the giant cells partially fused to the Reichert's membrane expressed Flt4 mRNA.

Thus, although Flt4 expression was very prominent in the earliest endothelial cell precursors, the angioblasts, it

appeared to be restricted only to certain vessels of 8.5 day p.c. embryos. The Tie receptor is known to be expressed in all endothelial cells of developing mouse embryos and thus provides a marker for these cells. See Korhonen, et al. *Oncogene*, 8:395 (1993); and Korhonen et al., *Blood*, 80: 2548-2555 (1992). Notably, in contrast to the Tie probe, the Flt4 probe hybridized very weakly if at all with arterial endothelia of 11.5 day p.c. embryos, e.g. with the endothelium of the developing dorsal aorta or the carotid arteries. Instead, Flt4 signal was much more prominent in the developing veins. For example, Flt4 signal was detected in veins surrounding the developing metanephros, while the Tie probe predominantly recognized capillaries within the metanephros.

Flt4 mRNA was observed to be distributed in several regions of a 12.5 day p.c. mouse embryo, being particularly prominent in the dilated vessel of the axillar region. A similar Flt4-positive vessel structure was seen in the mid-sagittal section in the jugular area (data not shown). A plexus-like pattern of Flt4-expressing vessels appeared in the periorbital region and surrounding the developing vertebrae. Also, just beneath the developing skin, a Flt4-positive vascular network was evident. Weaker capillary signals were obtained from several regions, including the developing brain. Flt4 mRNA could also be detected in small vessels of the neck region, of the developing snout and at the base of the developing tongue as well as in the tail region. Additionally, the liver was strongly positive for Flt4 mRNA in a spotlike pattern.

During further development, Flt4 mRNA appeared to become more restricted to certain vessels of the embryo. A 14.5 day p.c. embryo shows nicely this restricted pattern of expression. In the midsagittal section from such an embryo, the most prominent Flt4 signal was observed along the developing vertebral column in its anterior part. This signal was considered to originate from endothelial cells of the thoracic duct, which is the largest lymphatic vessel formed at this time of development. In contrast, the dorsal aorta and inferior vena cava were negative. Dilated vessels in the mesenteric region were also strongly positive for Flt4. Furthermore, as in the 12.5 day p.c. embryos, vessel networks along anatomical boundaries in the periorbital, lower jaw, as well as in the neck regions contained Flt4-positive endothelia. Similar structures were present in the pericardial space and throughout the subcutaneous tissue. Notably, in contrast to Flt4-negative vessels, all Flt4-positive vessels were devoid of blood cells in their lumen. These expression patterns suggest that Flt4 becomes confined to the endothelia of lymphatic vessels at this time of development. An additional site where we observed Flt4 expression was in the sinusoids of the developing bone marrow.

A transverse section of the upper thorax of a 16.5 day p.c. embryo hybridized with the Flt4 probe also was analyzed. Hematoxylin-eosin staining was performed to visualize the different types of vessels in this area. These include the carotid and brachiocephalic arteries, the vena cava, and the thoracic duct, which is smaller in size and lacks surrounding muscular and connective tissue. Under higher magnification endothelial cells of the thoracic duct as well as a small vessel in the vicinity were observed to hybridize with the Flt4 probe.

EXAMPLE 21

Analysis of Flt4 mRNA in Cultured Endothelial Cells

The in situ hybridization results described in Example 20 showed that Flt4 is expressed in venous endothelial cells and

later in lymphatic vessels and some venous endothelial cells, but not in arterial endothelia. In order to determine if such regulation was maintained in vitro, we studied cultured endothelial cells using Northern blotting and hybridization analysis.

Endothelial cells from human aorta, femoral vein, umbilical vein, and from foreskin microvessels were isolated, cultured, and characterized as previously described in the art. See Van Hinsberg et al., *Arteriosclerosis*, 7:389 (1987); and Van Hinsberg, et al., *Thromb. Haemostas*, 57:148 (1987). They were used at confluent density after five to eight passages (split ratio 1:3) for the isolation of polyadenylated RNA.

The endothelial cell lines EAhy926 (Edgell et al., *Proc. Natl. Acad. Sci.*, 80: 3734-3737 (1983)), BCE (Folkman et al., *Proc. Natl. Acad. Sci.*, 76: 5217-5221 (1979)) and LEH (Schreiber et al., *Proc. Natl. Acad. Sci.*, 82: 6138 (1985)) did not express Flt4. However, cultured human microvascular, venous and umbilical vein endothelial cells were positive for the Flt4-specific 5.8 and 4.5 kb mRNAs, whereas the aortic endothelial cells were negative. In contrast, another endothelial receptor tyrosine kinase gene, tie, was expressed as a 4.4 kb mRNA in all endothelial cell types studied.

EXAMPLE 22

Flt4 mRNA in Adult Human Tissues

The results obtained in Example 20 indicated that the Flt4 mRNA becomes largely confined to the endothelium of lymphatic vessels during development. Because of the potential significance of this finding in humans, we also studied Flt4 expression in adult human tissues using a human Flt4 probe. The human Flt4 probe used was an EcoRI-SphI fragment covering base pairs 1-595 of the cDNA (SEQ ID NO: 1). See also Pajusola et al., *Cancer Res.*, 52:5738 (1992). The von Willebrand factor probe was an EcoRI-HindIII fragment covering base pairs 1-2334. Bonthron, et al., *Nucleic Acids Res.*, 141:7125 (1986).

We used routinely fixed material sent for histopathological diagnosis. Normal lung tissue was obtained from a resection of the left inferior lung lobe affected by epidermoid cancer. Mesenterium and mesenterial lymph nodes were obtained from a patient having a colonic adenocarcinoma. A normal lymph node adjacent to the salivary gland was enucleated because of its abnormal size. The tonsils from two patients and the two appendixes had no diagnostic changes. Two lymphangiomyomas and three cystic lymphangiomas were studied with similar results.

For human tissues, which were routine samples fixed with 10% formalin for histopathological diagnosis, the normal in situ protocol gave just background, whereas microwave treatment instead of proteinase K enabled specific hybridization. Shi, et al., *J. Biol. Chem.*, 266:5774 (1991); Catoretti, et al., *J. of Pathol.*, 168:357 (1992).

In the mesenterium, lung and appendix lymphatic endothelia gave Flt4 signals, while veins, arteries, and capillaries were negative. To study whether Flt4 is expressed in the HEVs, the tonsils were studied. Indeed, in the tonsils, Flt4-specific autoradiographic grains were detected in some HEVs.

EXAMPLE 23

Analysis of Flt4 mRNA in Normal and Metastatic Lymph Node and in Lymphangioma

A portion of a human mesenterial lymph node (see Example 22) was analysed for Flt4 expression. Flt4 expres-

sion was observed in the lymphatic sinuses and afferent and efferent lymphatic vessels. The same pattern was observed in a lymph node containing adenocarcinoma metastases. Some HEVs in both normal and metastatic lymph node were also positive. Flt4 expression in a cystic lymphangioma was specific to lymphatic endothelia, as evident from a comparison with the in situ signals for von Willebrandt factor in all blood vessels.

Consistent with these results, immunostaining for Flt4 was strongly positive in the endothelium of cutaneous lymphangiomatosis, a rare disorder characterized by proliferation of presumed lymphatic endothelium. See Lymbousaki et al., *Am. J. Pathol.*, 153(2): 395-403 (August, 1998), incorporated herein by reference in its entirety.

Additionally, immunostaining for Flt4 identified spindle cells within Kaposi's sarcoma cutaneous nodular lesion tissue samples. See Jussila et al., *Cancer Res.*, 58:1599-1604 (April, 1998). In view of the apparent lymphatic specificity of Flt4, These results may be considered consistent with suggestions that certain cells in Kaposi's sarcoma are of lymphatic endothelial origin. See, e.g., Beckstead et al., *Am J. Pathol.*, 119: 294-300(1985); and Dictor et al., *Am J. Pathol.*, 130:411-417(1988).

EXAMPLE 24

Localization of Flt4 in Fetal Endothelial Cells

As described in Example 2, An Flt4 cDNA fragment encoding the 40 carboxy terminal amino acids of the short form was cloned as a 657 bp EcoRI-fragment into the pGEX-1 λ T bacterial expression vector (Pharmacia) in frame with the glutathione-S-transferase coding region. The resultant GST-Flt4 fusion protein was produced in *E. coli* and purified by affinity chromatography using a glutathione-Sepharose 4B column. The purified protein was lyophilized, dissolved in PBS, mixed with Freund's adjuvant, and used for immunization of rabbits. Antisera were used after the third booster immunization.

Tissues from 17 and 20-week-old human fetuses were obtained from legal abortions induced with prostaglandins. The study was approved by the Ethical Committee of the Helsinki University Central Hospital. The gestational age was estimated from the fetal foot length. The fetal tissues were embedded in Tissue-Tek (Miles), frozen immediately, and stored at -70° C.

Anti-Flt4 antiserum was cross-absorbed to a GST-Sepharose column to remove anti-GST-antibodies and then purified by GST-Flt4 affinity chromatography. Several 6 μ m-thick cryostat sections of the tissues were fixed with acetone and treated with 0.3% H₂O₂ in methanol for 30 minutes to block endogenous peroxidase activity. After washing, the sections were incubated with 5% normal swine serum. Sections were then incubated with antibodies against Flt4 and washed. Bound antibodies were detected with peroxidase-conjugated swine anti-rabbit IgG followed by staining for peroxidase activity using 0.2% 3,3'-diaminobenzidine (Amersham) as a substrate. The sections were counterstained in Meyer's hematoxylin.

Anti-Flt4 immunoperoxidase staining of human fetal mesenterium showed Flt4 protein in the endothelium of several vessels, while control stainings with antigen-blocked anti-Flt4 antibodies and preimmune sera were negative. For comparison, sections were stained with an antiserum against the Factor VIII-related antigen, which is specific for vascular endothelial cells. Immunoperoxidase staining for Flt4 was observed over endothelial cells of vessels, which did not

contain red blood cells, while blood vessels were negative. The vessels without red blood cells are likely to be lymphatic endothelial cells; such vessels are particularly frequent in the mesenterium. The antibodies against Factor VIII related antigen stained endothelial cells in all vessels.

EXAMPLE 25

Production of Monoclonal Antibodies Against Flt4 Fusion I

Recombinant Flt4 extracellular domain protein was produced by expressing the Flt4EC-6 \times His-pVTBac plasmid construct (Example 14) in High-Five cells. The Flt4 extracellular domain (Flt4EC) was purified from the culture medium of the infected High-Five cells using Ni-NTA affinity chromatography according to manufacturer's instructions (Qiagen) for binding and elution of the 6 \times His tag encoded in the COOH-terminus of the recombinant Flt4 extracellular domain.

Four month old Balb/c male mice were immunized by intraperitoneal injection of the purified, recombinantly produced Flt4 extracellular domain protein (150 μ g/mouse) emulsified with Freund's complete adjuvant. Booster injections of 150 μ g were given at three to four week intervals and a final booster (10 μ g Flt4 EC in PBS, administered intraperitoneally) was given after another three-week interval. Four days after the final booster dose, the mice were sacrificed and mouse splenic lymphoid cells were fused with SP 2/0 plasmacytoma cells at a 2:1 ratio, respectively.

The fused cells were harvested in 96-well culture plates (NUNC) in Ex-Cell 320 medium (SERALAB) containing 20% fetal calf serum and HAT supplement (hypoxanthine-aminopterin-thymidine; GIBCO, 043-01060H; diluted 50-fold). Cells were cultured at +37° C., in a 5% CO₂ atmosphere. After 10 days, HAT-supplemented medium was changed to HT-supplemented cell culture medium (GIBCO; 043-01065H, diluted 50-fold). HT medium is identical to HAT medium, but lacks aminopterin.

In three weeks, specific antibody production was determined by the antigen-specific ImmunoFluoroMetric Assay, (IFMA), described below in Example 26. The master clones were cloned by limited dilutions as described by Staszewski et al., *Yale Journal of Biology and Medicine*, 57:865-868 (1984). Positive clones were expanded onto 24-well tissue culture plates (NUNC), re-cloned, and re-tested by the same method. Positive clones were tested by fluorescence-activated cell sorting (FACS).

The stable clones secreted immunoglobulins belonging to the IgG₁ class, except one, which produced Ig probably belonging to class IgA. The subclass of monoclonal antibody was determined using rat monoclonal antibody to mouse subclass as biotin conjugate (SEROTEC) in IFMA.

Balb/c mice were used to produce monoclonal antibodies in ascites fluid. The hybridomas described above were intraperitoneally injected into mice after pretreatment of the animals with pristane (2,6,10,14-tetramethylpentadecane 98%, ALDRICH-CHEMIE D7924 Steinheim, Cat.No. T 2,280-2). 0.5 ml of pristane (i.v.) was injected about two weeks prior to the hybridoma cells. The amount of cells injected were approximately 7.5 to 9 \times 10⁶ per mouse. Ascites was collected 10 to 14 days after injection of the hybridomas.

Fusion II

Two month old Balb/c mice (female) were immunized by intraperitoneal injection of the recombinantly produced Flt4 extracellular domain protein (20 μ g/mouse), emulsified with Freund's complete adjuvant. Booster injections of 20 μ g were given at three to four week intervals and a final booster

(10 μ g Flt4 in PBS, administered i.v.) was given after another three-week interval. Four days after the final booster dose, the mice were sacrificed and mouse splenic lymphoid cells were fused with SP 2/0 plasmacytoma cells at a 2:1 ratio, respectively.

The fused cells were harvested in 96-well culture plates (FALCON) in OptiMEM 1 (with Glutamax, 1, 51985-026, GIBCO BRL) medium containing 20% fetal calf serum and HAT supplement (hypoxanthine-aminopterin-thymidine, GIBCO BRL 21060-017; diluted 1:50 fold). Cells were cultured at 37° C., in a 5% CO₂ atmosphere. After 10 days, HAT-supplemented medium was changed to HT-supplemental cell culture medium (GIBCO BRL; 41065-012, diluted 1:50-fold).

In three weeks, specific antibody production was determined by the antigen-specific ImmunoFluoroMetric Assay (IFMA) described below in Example 26. The master clones were cloned by limited dilutions as described by Staszewski et al. (1984). Positive clones were expanded onto 24-well tissue culture plates (FALCON), re-cloned, and re-tested by the same method. Positive clones were tested by FACS.

The 2E11 and 6B2 clones secreted immunoglobulins belonging to the IgG₁ class, and 2B 12 clones produced Ig belonging to subclass IgM. The mouse subclass IgG₁ was determined using rat monoclonal antibody against mouse subclass heavy chain as biotin conjugate (SEROTEC) in IFMA and the mouse subclass IgM was determined with Mouse Monoclonal Antibody Isotyping Kit (Dipstick Format) (19663-012, Life Technologies Inc.).

EXAMPLE 26

Specificity of Monoclonal Antibodies Against Flt4

The purified, recombinant Flt4 extracellular domain-6 \times His fusion product (produced as described in Examples 14 and 25) was labelled with Europium according to Mukkala et al., *Anal. Biochem.*, 176(2):319-325 (1989), with the following modification: a 250 times molar excess of isothiocyanate DTTA-Eu (N1 chelate, WALLAC, Finland) was added to the Flt4 solution (0.5 mg/ml in PBS) and the pH was adjusted to about 9 by adding 0.5 M sodium carbonate buffer, pH 9.8. The labelling was performed overnight at +4° C. Unbound label was removed using PD-10 (PHARMACIA, Sweden) with TSA buffer (50 mM Tris-HCl, pH 7.8, containing 0.15 M NaCl) as eluent.

After purification, 1 mg/ml bovine serum albumin (BSA) was added to the labelled Flt4 and the label was stored at +4° C. The average number of Europium ions incorporated per Flt4 molecule was 1.9, as determined by measuring the fluorescence in a ratio to that of known EuCl₃ standards (Hemmilä et al., *Anal. Biochem.*, 137:335-343 (1984)).

The antibodies produced in Example 25 were screened using a Sandwich-type immunofluorometric assay, using microtitration strip wells (NUNC, polysorb) coated with rabbit anti-mouse Ig (Z 259, DAKOPATTS). The pre-coated wells were washed once by Platwash 1296-024 (WALLAC) with DELFIA wash solution. The DELFIA assay buffer was used as a dilution buffer for cell culture supernatants and for serum of the splenectomized mouse (at dilutions between 1:1000 to 1:100,000) used as positive control in the preliminary screening assay.

An overnight incubation at +4° C. (or alternatively for 2 hours at room temperature) was begun by shaking on a Plateshake shaker (1296-001, WALLAC) for 5 minutes followed by washing four times with wash solution as described above.

The Europium-labelled Flt4 was added at a dilution of 1:500 in 100 μ l of the assay buffer. After 5 minutes on a Plateshake shaker and one hour incubation at room temperature, the strips were washed as described above.

Enhancement solution (DELFLIA) was added at 200 μ l/well. The plates were then shaken for 5 minutes on a Plateshake shaker and the intensity of fluorescence was measured by ARCUS-1230 (WALLAC) for 10-15 minutes. (Lövgren et al., In: Collins W. P. (Ed.) *Alternative Immunoassays*, John Wiley & Sons Ltd. (1985), pp. 203-216). The DELFLIA results show that all monoclonal antibodies tested bound the Flt4 EC antigen. Monoclonal antibodies reactive with the Flt4 (and the hybridomas which produce the antibodies) were selected for further screening.

The resulting monoclonal antibodies were used in double antibody immunofluorescence staining of NIH3T3 cells expressing the LTR-FLT41 construct and neomycin-resistant transfected NIH3T3 cells. The cells were detached from the culture plates using EDTA, stained, and analysed in a fluorescence-activated cell sorter (FACS). The results of FACS analysis are given as percentages of cells staining positive with the indicated monoclonal antibody (see Table 2, below).

TABLE 2

Mab clones	LTR % ^{a)}	NEO % ^{b)}	DELFLIA-counts
1B1	67.3	1	20625
1B1D11	75	1.2	19694
1B1F8	76.1	1.4	18580
4F6	69.9	1.2	23229
4F6B8G12	75	0.3	24374
4F6B8H11	75.9	0.3	28281
4F6B8E12	74.8	0.4	27097
4F6B8G10	75.3	0.4	26063
9D9	45.1	0.75	17316
9D9D10	71.7	2.3	18230
9D9F9	73	1.8	11904
9D9G6	74.3	2.9	16743
9D9G7	70.7	1.3	17009
10E4	24.2	1.4	39202
10E4B10E12	32.3	0.3	42490
10E4B10G10	36.5	0.3	54815
10E4B10F12	45.6	0.4	43909
10E4B10G12	45.7	0.5	35576
11G2	30.2	1.6	11304
11G2D12	74.4	1.5	14660
11G2G9	74.2	0.9	10283
11G2H7	74.4	2.1	25382

^{a)}FACS results with LTR transfected cells

^{b)}FACS results with NEO cells (control)

The FACS results with LTR-FLT41-transfected cells indicate that the antibodies effectively recognize Flt4-expressing cells. These same antibodies give only background staining of neomycin phosphotransferase-transfected NIH3T3 cells. Thus, the antibodies specifically recognize the Flt4 tyrosine kinase on the cell surface.

One clone, designated anti-Flt4 hybridoma 9D9F9, was found to stably secrete monoclonal antibody which was determined to be of immunoglobulin class IgG₁ by IFMA. Hybridoma 9D9F9 was deposited with the German Collection of Microorganisms and Cell Cultures, Department of Human and Animal Cell Cultures and Viruses, Mascheroder Weg 1b, 3300 Braunschweig, Germany, Mar. 23, 1995, and given accession No. ACC2210.

Fusion II Antibodies

The Europium-labelled Flt4 extracellular domain protein described above also was used to screen the Fusion II antibodies described in Example 25. The antibodies were screened using a Flt4-specific IFMA using microtitration

wells (Nunc, Polysorb) coated with rabbit anti-mouse Ig (Z 259, DAKO). The precoated wells were washed once with wash solution (Wallac) by using DELFIA Plate wash.

The DELFIA assay buffer was used as dilution buffer for cell culture supernatants (dilution 1:2 in preliminary screening) and for serum of the splenectomized mouse (dilutions 1:1000 to 1:100,000) which was used as a positive control. As standard, the purified anti-Flt4 9D9F9 (mouse subclass IgG₁) was used at concentrations between 1.0 ng/ml and 250 ng/ml. Samples were first shaken at room temperature for five minutes on a Plateshake shaker and then incubated approximately 18 hours at +4° C. The frames were first washed four times, then the Eu-labelled Flt4 (1:2000, in 100 μ l assay buffer) was added, and finally the frames were incubated for one hour at room temperature. After washing as described, the enhancement solution (200 μ l/well, Wallac) was added, and the frames were shaken for 5 minutes on the Plateshake shaker. The intensity of fluorescence was measured by ARCUS-1230 (Wallac). Monoclonal antibodies reactive with Flt4 were selected for further screening in the double antibody immunofluorescence staining assay employing Flt4-expressing NIH3T3 cells, as described above.

The resulting Fusion II monoclonal antibodies against Flt4 and corresponding results of FACS analysis (expressed as percentages of cells staining positive with the indicated monoclonal antibody) are summarized in Table 3.

A standard curve for quantitation of anti-Flt4 antibodies was made by using affinity purified anti-Flt4 9D9F9. The linear range reached from 1.0 ng/ml to 250 ng/ml.

Cell lysate of NIH3T3 cells co-transfected with pLTR-FLT41 construct expressing full-length Flt4 on the surface was electrophoresed in 6.5% SDS-PAGE, proteins were transferred onto nitrocellulose nitrate membrane (0.45 μ m, SCHLEICHER & SCHUELL) and immunoblotted with monoclonal antibody-containing hybridoma cell culture supernatants (1:10, 50 mM TRIS -40 mM glycine buffer containing methanol 4%, SDS 0.04%). The specificities of monoclonal antibodies were detected using incubation with HRP-conjugated rabbit antimouse Ig (P 161, DAKO, diluted 1:1000 in 20 mM TRIS buffer, pH 7.5, containing 150 mM saline, 5% milk powder) and ECL (Enhanced chemiluminescence, AMERSHAM).

TABLE 3

Mab clones	LTR % ^{a)}	NEO ^{b)}	approx. Mab production ng/ml/10 ⁶ cells ^{c)}	WB
2B12E10	39.5	6.0	440	+
2E11D11	44.6	8.8	110	+
2E11F9	49.5	4.5	100	+
2E11F12	46.0	4.1	180	+
2E11G8	41.2	7.8	160	+
6B2E12	NF	NF	1390	+
6B2F8	NF	NF	470	+
6B2G6	NF	NF	630	+
6B2H5	NF	NF	740	+
6B2H8	NF	NF	1800	+

^{a)}FACS results with LTR transfected cells

^{b)}FACS results with NEO cells (control)

^{c)}quantitation of Mab production based on affinity-purified antiFLT 9D9F9

antibody used as standard

NF not functioning in FACS

WB Used successfully in Western immunoblotting

EXAMPLE 27

Use of anti-Flt4 Antibodies to Identify Flt4 in Cell Lysates and Expressed in Lymphatic Endothelial Cells in Human Tissue

The monoclonal antibodies produced by hybridoma 9D9 described in the preceding examples were used in immuno-

precipitation and Western blotting of lysates of HEL cells. As reported in Example 6, Flt4 mRNA expression had been previously observed in HEL cells. About 2 \times 10⁷ cultured HEL cells were lysed in RIPA buffer specified in Example 11 and immunoprecipitated with about 2 micrograms of the 9D9 antibody (as described for polyclonal antibodies in example 11). For Western analysis, immunoprecipitates were electrophoresed via SDS-PAGE (6% gel) and electroblotted onto a nitrocellulose membrane. Polypeptide bands of 175 kD and 125 kD, corresponding to Flt4 polypeptides, were detected in the Western blotting analysis of the immunoprecipitates using a 1 microgram/ml dilution of the 9D9 antibody.

Immunostaining of human skin tissue was performed using the 9D9 monoclonal antibodies and an alkaline phosphatase ABC-AP kit (Dako). Briefly, slides containing 6 μ m samples of adult human skin were dried for 30 minutes at room temperature (RT), fixed for ten minutes with cold acetone, and then washed once for five minutes with phosphate-buffered saline (PBS). The samples were then incubated for 30 minutes at RT with 2% horse serum and washed three times for five minutes in PBS.

For immunostaining, the samples were incubated for one hour at RT with the 9D9 primary antibody and washed three times for five minutes with PBS. After washing, the samples were incubated for thirty minutes at RT with biotinylated rabbit anti-mouse secondary antibodies, and again washed three times for five minutes with PBS.

Bound antibodies were detected by incubating the samples for thirty minutes at RT with ABC-AP complex, washing three times with PBS, incubating for fifteen minutes at RT with AP-substrate (Sigma Fast Red TR/Naphtol AS-MX (Cat. No. F-4648)), and rinsing with water. Samples were then counter-stained with Mayer's hematoxylin for thirty seconds and rinsed with water. Aquamount and a coverslip were applied, and the samples were analyzed under a microscope. The 9D9 antibody staining was observed in lymphatic endothelial cells in these human skin sections. Blood vessel endothelia showed extremely weak or no staining. Additional analyses have served to confirm the apparent specificity for lymphatic endothelia. See Lymbousaki et al., *Am. J. Pathol.*, 153(2):395-403 (August, 1998); and Jussila et al., *Cancer Res.*, 58:1599-1604 (April, 1998), both of which are incorporated herein by reference in their entireties.

These results further confirm the utility of Flt4 as a useful marker for lymphatic endothelia and the utility of anti-Flt4 antibodies for identifying and visualizing Flt4 expressed in these cells, in a tissue sample.

EXAMPLE 28

Upregulation of the VEGF-C/VEGFR-3 Signalling Pathway in Breast Cancer Angiogenesis

The foregoing examples demonstrate that Flt4 (VEGFR-3) is useful as a specific antigenic marker for lymphatic endothelia in normal tissues. The following procedures additionally demonstrate that VEGFR-3 is useful as an antigenic marker (e.g., for diagnosis and screening) and as a therapeutic target in malignant breast tumors. A highly elevated number of VEGFR-3 positive vessels was found in invasive breast cancer in comparison to histologically normal breast tissue (P<0.0001).

MATERIALS AND METHODS

Freshly frozen breast tissue samples were retrieved from the files of the Department of Pathology, University of

Helsinki. The samples consisted of ductal carcinoma (n=6), lobular carcinoma (n=6), intraductal carcinoma (n=8), fibroadenoma (n=4), and histologically normal breast tissue (n=12). All samples had been frozen immediately after surgical excision in liquid nitrogen, and stored at -70° C.

Mouse monoclonal antibodies (Mabs) against human Flt4 (VEGFR-3) were produced essentially as described in preceding examples, e.g., Example 25. The VEGFR-3 extracellular protein domain (VEGFR-3EC) was expressed via a recombinant baculovirus in insect cells, purified from the culture medium. Mouse monoclonal antibodies against VEGFR-3EC were then produced using standard methods and the immunoglobulin fraction was purified by protein A affinity chromatography from hybridoma ascites fluid or Tecnomouse® culture supernatants.

Five μ m cryosections of the tissues samples were air-dried and fixed in cold acetone for 10 minutes. The sections were re-hydrated in phosphate buffered saline (PBS) and incubated for 30 minutes in 5% normal horse serum at room temperature. The sections were then incubated for 2 hours in a humid atmosphere at room temperature with the Mabs 9D9F9 (Example 26) at the concentration of 1.0 μ g/ml. Other anti-VEGFR-3 Mab against distinct epitopes of the VEGFR-3EC were also studied; clones 2E11D11 (Example 26) and 7B8F9 (made essentially as described in Example 26) were used at the concentrations of 9.5 and 8.5 μ g/ml, respectively. A subsequent incubation for 30 minutes in biotinylated anti-mouse serum was followed by a 60 minute incubation using reagents of the Vectastain Elite Mouse IgG ABC kit (Vector laboratories, Burlingame, USA). Peroxidase activity was developed with 3-amino-9-ethyl carbazole (AEC, Sigma, St. Louis, USA) for 10 minutes. Finally, the sections were stained with haematoxylin for 20 seconds. Negative controls were performed by omitting the primary antibody, or by using irrelevant primary antibodies of the same isotype. The purified baculoviral immunogen was used to block the binding of the 9D9 antibodies as another negative control. In these experiments, the antibodies were incubated overnight with a 10-fold molar excess of the VEGFR-3EC protein in PBS. After centrifugation for 4 minutes at 4000 rpm, +4° C., the supernatant was carefully collected and then used as primary antibody. The 5 μ m cryosections adjacent to the ones stained with the anti-VEGFR-3 antibodies were immunostained for the blood vascular endothelial marker PAL-E (0.15 μ g/ml, Monosan, Uden, the Netherlands), laminin (1:4000 dilution of the supernatant of clone LAM-89, Sigma, St Louis, Mo.), collagen XVIII (1.9 μ g/ml), α -smooth muscle actin (SMA, 0.5 μ g/ml, clone 1A4, Sigma), VEGFR-1 (1:200 dilution of the supernatant of clone 19) or VEGFR-2 (dilution 1:100).

Pathological examination of all of the samples was performed after the staining procedures. The blood vascular densities were obtained from the slides stained for PAL-E [de Waal et al., *Am. J. Pathol.*, 150: 1951-1957 (1997)], following the guidelines recommended by Gasparini and Harris. [Gasparini G, and Harris A, *J. Clin. Oncol.*, 13: 765-782 (1995).] The VEGFR-3 positive vessel densities were studied in the same way. A slide was first scanned at low magnification, and intratumoral vessel density was then assessed by counting the number of stained vessels per a 400 \times magnification high power field (hpf) in the areas with the highest vascular density ("vascular hotspots") or in the areas with highest VEGFR-3 positive vessel density. A minimum of 5 fields was counted per a slide, after which the 3 highest counts were averaged.

Double staining was performed to differentiate immunohistochemical staining of lymphatic and blood vessels in two

intraductal carcinomas. Acetone-fixed 5 gm cryosections were were incubated for 1 hour with anti-PAL-E antibodies, with biotinylated horse anti-mouse antibody (Vectastain Elite Mouse IgG ABC kit, Vector laboratories, Burlingame, USA) for 30 minutes, with ABC-peroxidase (Vectastain, 1:100) for 45 minutes, and developed finally with AEC for 10 minutes. For the second step, the sections were incubated with anti-VEGFR-3 antibodies for 1 hour (0.14 μ g/ml), followed by biotinylated anti-mouse antibody for 30 minutes (1:200 dilution of the supernatant of clone), ABC-peroxidase for 30 minutes (1:100), biotinylated tyramin solution (1:2.000) containing 0.01% peroxide for 5 minutes, ABC-alkaline phosphatase (1:100) for 20 minutes, and developed with Fast Blue (Sigma, St. Louis, USA) for 20 minutes, according to a procedure previously described in the literature for ISH signal enhancement. [Kerstens et al., *J. Histochem. Cytochem.*, 43: 347-352 (1995).] Cryosections (5 μ m) adjacent to the double-stained sections were also immunostained with VEGFR-3 antibodies only, as described above.

Polyclonal antibodies were produced in rabbits against a synthetic peptide corresponding to the amino acid residues 2-18 of the N-terminus of mature, secreted human vascular endothelial growth factor C (VEGF-C) (residues 104-120 of the VEGF-C prepro-VEGF-C polypeptide) as described in the literature. [Joukov et al., *EMBO J.*, 16: 3898-3911 (1997), incorporated herein by reference in its entirety.] The antisera were affinity-purified using the immunogenic polypeptide coupled to an epoxy-activated sepharose-6B column and tested for specific staining of VEGF-C using cells infected with an adenoviral vector expressing VEGF-C or control β -galactosidase.

The eight intraductal carcinomas and all of the invasive carcinomas analysed for VEGFR-3 were chosen for further analyses of the expression of VEGF-C. Five micrometer cryosections adjacent to the sections stained with the anti-VEGFR-3 antibodies were air-dried and fixed in cold acetone for 10 minutes. The sections were rehydrated in PBS and incubated for 30 minutes in 5% normal goat serum and then for 2 hours in a humid atmosphere at room temperature with the rabbit polyclonal antibodies against human VEGF-C, diluted 1:200 in PBS. A subsequent incubation for 30 minutes in biotinylated anti-rabbit serum was followed by a 60 minutes incubation using reagents of the Vectastain Elite Rabbit IgG ABC kit (Vector laboratories, Burlingame, USA). The sections were further processed as described above. As a negative control, the purified immunogen was used to block the binding of the VEGF-C antibodies. In these experiments, VEGF-C antibodies were incubated overnight with a 10-fold molar excess of the VEGF-C protein in PBS. After centrifugation for 4 minutes at 4000 rpm at +4° C., the supernatant was carefully collected and used in the immunostainings.

Monoclonal antibodies to human type XVIII collagen were generated by DiaBor Ltd. (Oulu, Finland) by immunization of mice with the recombinant polypeptide QH48.18 [Saarela et al., *Matrix Biolog.*, 16: 319-28 (1998)], corresponding to the common region of the N-terminal NC domain of human type XVIII collagen. The clones were screened by ELISA assay and Western analysis using the polypeptide QH48.18, and also by immunofluorescence staining of frozen human tissue sections. The screening of the hybridoma clones resulted in three monoclonal antibodies, which were positive in all three assays mentioned (ELISA, Western, immunofluorescence staining). One of the antibodies which gave the strongest signals, DB 144-N2, was used in subsequent experiments. It gave an

identical staining pattern (e.g., in adult human skin and kidney samples) to that of the polyclonal anti-all hu(XVIII).

RESULTS

A. VEGFR-3 in Histologically Normal Breast Tissue and in Benign Fibroadenomas

Immunohistochemical staining of VEGFR-3 in normal breast tissue showed a very weak staining in capillaries of the interductal stroma. These vessels did not form any specific pattern, but were scattered throughout the stroma. The density of the VEGFR-3 positive vessels in the normal breast tissue samples ranged from 6 to 17 per hpf, median 9 (n=12). Most of such vessels were strongly stained for the blood vascular endothelial marker PAL-E and for the basal lamina component, collagen XVIII, suggesting that VEGFR-3 was expressed weakly in the blood vessels of normal breast tissue. However, some thin vessels in the stroma, which were clearly stained for VEGFR-3 were negative for PAL-E and only weakly positive for the collagen type XVIII, suggesting that they were lymphatic vessels. VEGFR-3 positive vessels were also uniformly found in benign fibroadenomas, where their density (median 8 vessels per hpf; range 3–19; n=4) did not differ from that of the histologically normal breast tissue (median 8 vs. 9; P>0.1, the Mann-Whitney test).

B. VEGFR-3 Positive Vessels in Intraductal Carcinomas

In intraductal carcinomas, a distinctive pattern of strongly-stained VEGFR-3 positive vessels was observed. The vessels formed arch-like structures around the affected ducts (FIG. 5A). This “necklace” pattern also was observed in staining of adjacent sections for the blood vessel endothelial marker, PAL-E (FIG. 5B), suggesting that VEGFR-3 expression was enhanced in capillary endothelium. In order to more definitively differentiate between blood and lymphatic vessels and to search for the presence of smooth muscle cells and pericytes in the vessel walls, additional stainings were done using antibodies against smooth muscle α -actin (SMA) and basal lamina components laminin and type XVIII collagen. According to this staining, the small vessels close to the intraductal carcinomas expressed simultaneously VEGFR-3 and the basal lamina proteins, but stained more weakly for SMA, indicating that they are incompletely covered by pericytes/smooth muscle cells in the vessel wall (black arrows in FIGS. 5C–5F). In contrast, larger blood vessels at some distance from the intraductal lesions were in general negative for VEGFR-3, but positive for laminin, collagen XVIII and SMA (red arrows). In addition, vessels were found, which were positive for VEGFR-3, but only very weakly stained for the basal lamina proteins laminin and type XVIII collagen and not at all for SMA (green arrows). These were considered to represent lymphatic vessels.

C. Differential Double-staining of Blood and Lymphatic Vessels

Two intraductal carcinomas were chosen for the immunohistochemical double-staining procedure to more clearly differentiate lymphatic vessels from blood vessels. [See de Waal et al., *Am. J. Pathol.*, 150: 1951–1957 (1997).] Using this method, the VEGFR-3 positive vessels were stained blue, while the PAL-E positive vessels and basal laminae were stained brown. Both tested samples showed a similar pattern of staining: the vessels lining the tumor filled ducts were predominantly PAL-E positive (arrowhead in FIGS. 5G and 5H) while the presumably lymphatic, VEGFR-3 positive vessels a short distance away in the interductal stroma were PAL-E negative (black arrows in FIGS. 5G and 5H). In order to exclude misinterpretation due to possible double-staining artefacts, adjacent 5 μ m sections were

stained with anti-VEGFR-3 alone. This staining confirmed that several of the PAL-E positive blood vessels are also positive for VEGFR-3.

D. VEGF-C, VEGFR-1 and VEGFR-2 in the Intraductal Carcinoma Cells and its Receptors in Adjacent Vessels

Polyclonal affinity-purified antibodies against human VEGF-C were used to stain the 8 intraductal carcinoma samples. All tested samples contained at least some VEGF-C, but considerable heterogeneity was observed in the intensity of staining and in the expression patterns. In some cases, most of the carcinoma cells were strongly positive for VEGF-C, while in others, only some carcinoma cells gave a staining signal. In contrast, very little or no staining was observed in the normal tissues surrounding the affected ducts, although weak signal was also obtained in unaffected normal ductal epithelium. Antigen blocking experiments indicated that the staining for VEGF-C was specific. The other VEGF-C receptor, VEGFR-2, as well as the other VEGF receptor (VEGFR-1), were both expressed in the same “necklace” vessels adjacent to the intraductal carcinoma cells.

E. VEGFR-3 Positive Vessels and VEGF-C in Invasive Breast Carcinoma

Strongly-stained VEGFR-3 positive vessels were also present in all invasive ductal carcinomas and lobular carcinomas studied. The VEGFR-3 positive vessels did not appear to form any specific distribution pattern; most of these vessels were also immunoreactive for the PAL-E antigen. The intratumoral VEGFR-3 positive vessel density (median 21, range 9–56 vessels per hpf; n=12) was significantly elevated in the invasive breast carcinomas when compared with normal breast tissue (median 21 vs. 9; P<0.0001, the Mann-Whitney test). Occasionally, invasion of the carcinoma cells into the VEGFR-3 positive lymphatic vessels could be observed.

Immunostaining for VEGF-C varied strongly among the invasive carcinomas studied (n=12). Some carcinoma cells were strongly positive for VEGF-C, while others stained very weakly or, in some cases, no staining was observed. Like in the intraductal carcinomas, very little or no staining was observed in the connective tissue in these sections.

The foregoing data reveals that VEGFR-3, which had otherwise appeared to be a predominantly lymphatic endothelial marker in most adult tissues, is very weakly expressed also in capillary endothelium of normal breast tissue. More significantly, in intraductal carcinomas, a “necklace” pattern of strongly-stained VEGFR-3 positive vessels was detected lining the tumor-filled ducts. Most of these vessels expressed the blood vessel endothelial marker PAL-E and the basal lamina components laminin and collagen XVIII, but apparently had less pericytes/smooth muscle cells than blood vessels located further away from the tumor cells, as shown by staining using antibodies against SMA. These features suggest that the “necklace” vessels were undergoing angiogenesis. A second group of vessels lying a distance away from the affected ducts were positive for VEGFR-3 but very weakly positive for the basal lamina components and negative for PAL-E, suggesting that they are lymphatic vessels. These vessels also lacked SMA-positive pericytic components. Also in invasive breast carcinomas, VEGFR-3 was upregulated in PAL-E positive blood vessels, although the vessel patterns seen were more randomly organized in the connective tissue stroma around the tumor cells. The results indicate that VEGFR-3 expression is upregulated in breast carcinomas during angiogenesis associated with tumor growth. The highly elevated number of VEGFR-3 positive vessels found in carcinoma in situ is compatible with the

hypothesis that the carcinoma cells produce factors, which activate the growth of blood vessels in the immediate vicinity of the carcinoma cells.

Since VEGF-C binds both VEGFR-3 and VEGFR-2 with high affinity, and since both intraductal and invasive carcinoma cells often stained positive for VEGF-C protein, this growth factor is a candidate growth factor for the VEGFR-3 and VEGFR-2 positive vessels in the carcinomas. These data are in agreement with another study, in which nearly half of thirty-five unselected malignant invasive tumors (including breast carcinomas, squamous cell carcinomas, lymphomas, melanomas, and sarcomas) contained VEGF-C mRNA in Northern blotting analyses. [See Salven et al., *Am. J. Pathol.*, 153(1): 103-108 (July, 1998), incorporated herein by reference in its entirety.] Collectively, the data reported herein provide an indication for treatment of breast carcinomas and possibly other, non-lymphatic carcinomas with agents that block the VEGF-C mediated stimulation of VEGFR-3 and/or VEGFR-2. Contemplated blocking agents include: anti-VEGF-C antibodies; anti-VEGFR-3 antibodies; anti-VEGFR-2 antibodies; bispecific antibodies that bind to VEGFR-3 and either VEGFR-2 or VEGFR-1; soluble extracellular domain fragments of VEGFR-3 that will bind circulating VEGF-C; VEGF-C fragments and analogs that bind VEGFR-3 and/or VEGFR-2 and that inhibit activation of such receptors; VEGF-C polypeptides, fragments, and analogs that bind VEGFR-3 and/or VEGFR-2 and that are conjugated to a suitable therapeutic agent; VEGFR-3 tyrosine kinase inhibitors; and small molecules that bind and inhibit these receptors. In addition, since VEGF-D binds both VEGFR-3 and VEGFR-2, it is contemplated that anti-VEGF-D antibodies and inhibitory VEGF-D fragments and analogs are suitable blocking agents. Human or humanized antibodies and fragments thereof are preferred, to the extent that antibody agents are selected for human therapy. Additionally, it is contemplated, as an additional aspect of the invention, to use any of the foregoing agents to evaluate mammalian tissue *in vitro* or *in vivo*, e.g., for the purposes of diagnosis and screening for malignancies and the spread of malignancies.

For any of the foregoing agents, it is contemplated that the agent may be further improved for diagnosis and screening by the attachment of a detectable label, including but not limited to radioisotopes (e.g., ^{14}C , ^{133}I and ^{125}I), chromophores (e.g., fluorescein, phycobiliprotein; tetraethyl rhodamine; enzymes which produce a fluorescent or colored product for detection by fluorescence; absorbance, visible color, or agglutination, which produces an electron-dense product for detection by electron microscopy); or electron dense molecules such as ferritin, peroxidase, or gold beads. Likewise, the agents may be further improved for therapeutic purposes by attachment (e.g., conjugation) or co-administration with molecules having anti-neoplastic properties, such as toxins of plant, animal, microbial, or fungal origin; radioisotopes; drugs; enzymes; and/or cytokines and other therapeutic proteins. (See, e.g., Pietersz & McKenzie, "Antibody Conjugates for the treatment of Cancer," *Immunological Reviews*, 129:57-80 (1992), incorporated by reference herein.

EXAMPLE 29

Anti-Fit4 Antibodies for Administration as a Therapeutic to Humans

A. Humanization of Anti-Fit4 Monoclonal Antibodies

The biology of Flt4 as reported herein, e.g., in Example 28, indicates therapeutic uses for Flt4 inhibitors (antagonists) that block ligand-mediated signalling of the

Flt4 receptor. Flt4-neutralizing antibodies comprise one class of therapeutics useful as Flt4 antagonists. Following are protocols to improve the utility of anti-Flt4 monoclonal antibodies as therapeutics in humans, by "humanizing" the monoclonal antibodies to improve their serum half-life and render them less immunogenic in human hosts (i.e., to prevent human antibody response to non-human anti-Flt4 antibodies).

The principles of humanization have been described in the literature and are facilitated by the modular arrangement of antibody proteins. To minimize the possibility of binding complement, a humanized antibody of the IgG4 isotype is preferred.

For example, a level of humanization is achieved by generating chimeric antibodies comprising the variable domains of non-human antibody proteins of interest, such as the anti-Flt4 monoclonal antibodies described herein, with the constant domains of human antibody molecules. (See, e.g., Morrison and Oi, *Adv. Immunol.*, 44:65-92 (1989).) The variable domains of Flt4 neutralizing anti-Flt4 antibodies are cloned from the genomic DNA of a B-cell hybridoma or from cDNA generated from mRNA isolated from the hybridoma of interest. The V region gene fragments are linked to exons encoding human antibody constant domains, and the resultant construct is expressed in suitable mammalian host cells (e.g., myeloma or CHO cells).

To achieve an even greater level of humanization, only those portions of the variable region gene fragments that encode antigen-binding complementarity determining regions ("CDR") of the non-human monoclonal antibody genes are cloned into human antibody sequences. [See, e.g., Jones et al., *Nature*, 321:522-525 (1986); Riechmann et al., *Nature*, 332:323-327 (1988); Verhoeven et al., *Science*, 239:1534x36 (1988); and Tempest et al., *Bio/Technology*, 9:266-71 (1991).] If necessary, the β -sheet framework of the human antibody surrounding the CDR3 regions also is modified to more closely mirror the three dimensional structure of the antigen-binding domain of the original monoclonal antibody. (See Kettleborough et al., *Protein Engin.*, 4:773-783 (1991); and Foote et al., *J. Mol. Biol.*, 224:487-499 (1992).)

In an alternative approach, the surface of a non-human monoclonal antibody of interest is humanized by altering selected surface residues of the non-human antibody, e.g., by site-directed mutagenesis, while retaining all of the interior and contacting residues of the non-human antibody. See Padlan, *Molecular Immunol.*, 28(4/5):489-98 (1991).

The foregoing approaches are employed using Flt4-neutralizing anti-Flt4 monoclonal antibodies and the hybridomas that produce them, such as antibodies 9D9F9, to generate humanized Flt4-neutralizing antibodies useful as therapeutics to treat or palliate conditions wherein Flt4 expression is detrimental.

B. Human Flt4-Neutralizing Antibodies From Phage Display

Human Flt4-neutralizing antibodies are generated by phage display techniques such as those described in Aujame et al., *Human Antibodies*, 8(4):155-168 (1997); Hoogenboom, *TIBTECH*, 15:62-70 (1997); and Rader et al., *Curr. Opin. Biotechnol.*, 8:503-508 (1997), all of which are incorporated by reference. For example, antibody variable regions in the form of Fab fragments or linked single chain Fv fragments are fused to the amino terminus of filamentous phage minor coat protein pIII. Expression of the fusion protein and incorporation thereof into the mature phage coat results in phage particles that present an antibody on their surface and contain the genetic material encoding the anti-

body. A phage library comprising such constructs is expressed in bacteria, and the library is panned (screened) for Flt4-specific phage-antibodies using labelled or immobilized Flt4 as antigen-probe.

C. Human Flt4-neutralizing Antibodies From Transgenic Mice

Human Flt4-neutralizing antibodies are generated in transgenic mice essentially as described in Bruggemann and Neuberger, *Immunol. Today*, 17(8):391-97 (1996) and Bruggemann and Taussig, *Curr. Opin. Biotechnol.*, 8:455-58 (1997). Transgenic mice carrying human V-gene segments in germline configuration and that express these transgenes in their lymphoid tissue are immunized with an Flt4 composition using conventional immunization protocols. Hybridomas are generated using B cells from the immunized mice using conventional protocols and screened to identify hybridomas secreting anti-Flt4 human antibodies (e.g., as described above).

D. Bispecific Antibodies

Bispecific antibodies that specifically bind to Flt4 and that specifically bind to other antigens relevant to pathology and/or treatment are produced, isolated, and tested using standard procedures that have been described in the literature. See, e.g., Pluckthun & Pack, *Immunotechnology*, 3:83-105 (1997); Carter et al., *J. Hematotherapy*, 4: 463-470 (1995); Renner & Pfreundschuh, *Immunological Reviews*, 1995, No. 145, pp. 179-209; Pfreundschuh U.S. Pat. No. 5,643,759; Segal et al., *J. Hematotherapy*, 4: 377-382 (1995); Segal et al., *Immunobiology*, 185: 390-402 (1992); and Bolhuis et al., *Cancer Immunol. Immunother.*,

34: 1-8 (1991), all of which are incorporated herein by reference in their entireties.

EXAMPLE 30

Animal Models to Demonstrate the Efficacy of anti-Flt4 Therapies for Treatment of Cancers

It is contemplated that any accepted animal for cancer therapies would be useful to demonstrate the efficacy of anti-Flt4 therapies for cancer treatment. Exemplary models for demonstrating the efficacy for treatment of breast cancers, using standard dose-response studies, include those described in Tekmal and Durgam, *Cancer Lett.*, 118(1): 21-28 (1997); Moshakis et al., *Br. J. Cancer*, 43: 575-580 (1981); and Williams et al., *J. Nat. Cancer Inst.*, 66: 147-155 (1981). In addition to murine models, dog and pig models are contemplated because at least certain anti-human Flt4 antibodies (e.g., the 9D9 antibodies) also recognize Flt4 from dog and pig. Tumor size and side effects are monitored to demonstrate therapeutic efficacy versus controls.

While the invention here has been described in connection with specific embodiments thereof, it will be understood that it is capable of further modifications and this application is intended to cover any variations, uses, or adaptations of the invention following, in general, the principles of the invention and including such departures from the present disclosure as come within known or customary practice within the art to which the invention pertains and as may be applied to the essential features hereinbefore set forth and as follows in the scope of the appended claims.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 22

<210> SEQ ID NO 1

<211> LENGTH: 4195

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<220> FEATURE:

<221> NAME/KEY: CDS

<222> LOCATION: (20)..(3913)

<400> SEQUENCE: 1

```
ccacgcgcag cgcccgag atg cag cgg gcc gcc gcg ctg tgc ctg cga ctg      52
      Met Gln Arg Gly Ala Ala Leu Cys Leu Arg Leu
      1              5              10

tgg ctc tgc ctg gga ctc ctg gac gcc ctg gtg agt gcc tac tcc atg      100
Trp Leu Cys Leu Gly Leu Leu Asp Gly Leu Val Ser Gly Tyr Ser Met
      15              20              25

acc ccc ccg acc ttg aac atc acg gag gag tca cac gtc atc gac acc      148
Thr Pro Pro Thr Leu Asn Ile Thr Glu Glu Ser His Val Ile Asp Thr
      30              35              40

ggt gac agc ctg tcc atc tcc tgc agg gga cag cac ccc ctc gag tgg      196
Gly Asp Ser Leu Ser Ile Ser Cys Arg Gly Gln His Pro Leu Glu Trp
      45              50              55

gct tgg cca gga gct cag gag gcg cca gcc acc gga gac aag gac agc      244
Ala Trp Pro Gly Ala Gln Glu Ala Pro Ala Thr Gly Asp Lys Asp Ser
      60              65              70              75

gag gac acg ggg gtg gtg cga gac tgc gag gcc aca gac gcc agg ccc      292
Glu Asp Thr Gly Val Val Arg Asp Cys Glu Gly Thr Asp Ala Arg Pro
      80              85              90

tac tgc aag gtg ttg ctg ctg cac gag gta cat gcc aac gac aca gcc      340
```

-continued

Tyr	Cys	Lys	Val	Leu	Leu	Leu	His	Glu	Val	His	Ala	Asn	Asp	Thr	Gly	
			95					100					105			
agc	tac	gtc	tgc	tac	tac	aag	tac	atc	aag	gca	cgc	atc	gag	ggc	acc	388
Ser	Tyr	Val	Cys	Tyr	Tyr	Lys	Tyr	Ile	Lys	Ala	Arg	Ile	Glu	Gly	Thr	
		110					115					120				
acg	gcc	gcc	agc	tcc	tac	gtg	ttc	gtg	aga	gac	ttt	gag	cag	cca	ttc	436
Thr	Ala	Ala	Ser	Ser	Tyr	Val	Phe	Val	Arg	Asp	Phe	Glu	Gln	Pro	Phe	
	125					130				135						
atc	aac	aag	cct	gac	acg	ctc	ttg	gtc	aac	agg	aag	gac	gcc	atg	tgg	484
Ile	Asn	Lys	Pro	Asp	Thr	Leu	Leu	Val	Asn	Arg	Lys	Asp	Ala	Met	Trp	
	140				145				150					155		
gtg	ccc	tgt	ctg	gtg	tcc	atc	ccc	ggc	ctc	aat	gtc	acg	ctg	cgc	tcg	532
Val	Pro	Cys	Leu	Val	Ser	Ile	Pro	Gly	Leu	Asn	Val	Thr	Leu	Arg	Ser	
			160					165					170			
caa	agc	tcg	gtg	ctg	tgg	cca	gac	ggg	cag	gag	gtg	gtg	tgg	gat	gac	580
Gln	Ser	Ser	Val	Leu	Trp	Pro	Asp	Gly	Gln	Glu	Val	Val	Trp	Asp	Asp	
		175					180					185				
cgg	cgg	ggc	atg	ctc	gtg	tcc	acg	cca	ctg	ctg	cac	gat	gcc	ctg	tac	628
Arg	Arg	Gly	Met	Leu	Val	Ser	Thr	Pro	Leu	Leu	His	Asp	Ala	Leu	Tyr	
		190					195					200				
ctg	cag	tgc	gag	acc	acc	tgg	gga	gac	cag	gac	ttc	ctt	tcc	aac	ccc	676
Leu	Gln	Cys	Glu	Thr	Thr	Trp	Gly	Asp	Gln	Asp	Phe	Leu	Ser	Asn	Pro	
	205					210					215					
ttc	ctg	gtg	cac	atc	aca	ggc	aac	gag	ctc	tat	gac	atc	cag	ctg	ttg	724
Phe	Leu	Val	His	Ile	Thr	Gly	Asn	Glu	Leu	Tyr	Asp	Ile	Gln	Leu	Leu	
	220				225				230					235		
ccc	agg	aag	tcg	ctg	gag	ctg	ctg	gta	ggg	gag	aag	ctg	gtc	ctg	aac	772
Pro	Arg	Lys	Ser	Leu	Glu	Leu	Leu	Val	Gly	Glu	Lys	Leu	Val	Leu	Asn	
			240					245				250				
tgc	acc	gtg	tgg	gct	gag	ttt	aac	tca	ggt	gtc	acc	ttt	gac	tgg	gac	820
Cys	Thr	Val	Trp	Ala	Glu	Phe	Asn	Ser	Gly	Val	Thr	Phe	Asp	Trp	Asp	
		255					260					265				
tac	cca	ggg	aag	cag	gca	gag	cgg	ggt	aag	tgg	gtg	ccc	gag	cga	cgc	868
Tyr	Pro	Gly	Lys	Gln	Ala	Glu	Arg	Gly	Lys	Trp	Val	Pro	Glu	Arg	Arg	
		270					275					280				
tcc	cag	cag	acc	cac	aca	gaa	ctc	tcc	agc	atc	ctg	acc	atc	cac	aac	916
Ser	Gln	Gln	Thr	His	Thr	Glu	Leu	Ser	Ser	Ile	Leu	Thr	Ile	His	Asn	
		285				290					295					
gtc	agc	cag	cac	gac	ctg	ggc	tcg	tat	gtg	tgc	aag	gcc	aac	aac	ggc	964
Val	Ser	Gln	His	Asp	Leu	Gly	Ser	Tyr	Val	Cys	Lys	Ala	Asn	Asn	Gly	
	300				305				310					315		
atc	cag	cga	ttt	cgg	gag	agc	acc	gag	gtc	att	gtg	cat	gaa	aat	ccc	1012
Ile	Gln	Arg	Phe	Arg	Glu	Ser	Thr	Glu	Val	Ile	Val	His	Glu	Asn	Pro	
		320						325				330				
ttc	atc	agc	gtc	gag	tgg	ctc	aaa	gga	ccc	atc	ctg	gag	gcc	acg	gca	1060
Phe	Ile	Ser	Val	Glu	Trp	Leu	Lys	Gly	Pro	Ile	Leu	Glu	Ala	Thr	Ala	
		335					340					345				
gga	gac	gag	ctg	gtg	aag	ctg	ccc	gtg	aag	ctg	gca	gcg	tac	ccc	ccg	1108
Gly	Asp	Glu	Leu	Val	Lys	Leu	Pro	Val	Lys	Leu	Ala	Ala	Tyr	Pro	Pro	
		350				355						360				
ccc	gag	ttc	cag	tgg	tac	aag	gat	gga	aag	gca	ctg	tcc	ggg	cgc	cac	1156
Pro	Glu	Phe	Gln	Trp	Tyr	Lys	Asp	Gly	Lys	Ala	Leu	Ser	Gly	Arg	His	
		365				370					375					
agt	cca	cat	gcc	ctg	gtg	ctc	aag	gag	gtg	aca	gag	gcc	agc	aca	ggc	1204
Ser	Pro	His	Ala	Leu	Val	Leu	Lys	Glu	Val	Thr	Glu	Ala	Ser	Thr	Gly	
				385					390					395		
acc	tac	acc	ctc	gcc	ctg	tgg	aac	tcc	gct	gct	ggc	ctg	agg	cgc	aac	1252
Thr	Tyr	Thr	Leu	Ala	Leu	Trp	Asn	Ser	Ala	Ala	Gly	Leu	Arg	Arg	Asn	
			400						405					410		

-continued

atc agc ctg gag ctg gtg gtg aat gtg ccc ccc cag ata cat gag aag Ile Ser Leu Glu Leu Val Val Asn Val Pro Pro Gln Ile His Glu Lys 415 420 425	1300
gag gcc tcc tcc ccc agc atc tac tgg cgt cac agc cgc cag gcc ctc Glu Ala Ser Ser Pro Ser Ile Tyr Ser Arg His Ser Arg Gln Ala Leu 430 435 440	1348
acc tgc acg gcc tac ggg gtg ccc ctg cct ctc agc atc cag tgg cac Thr Cys Thr Ala Tyr Gly Val Pro Leu Pro Leu Ser Ile Gln Trp His 445 450 455	1396
tgg cgg ccc tgg aca ccc tgc aag atg ttt gcc cag cgt agt ctc cgg Trp Arg Pro Trp Thr Pro Cys Lys Met Phe Ala Gln Arg Ser Leu Arg 460 465 470 475	1444
cgg cgg cag cag caa gac ctc atg cca cag tgc cgt gac tgg agg gcg Arg Arg Gln Gln Asp Leu Met Pro Gln Cys Arg Asp Trp Arg Ala 480 485 490	1492
gtg acc acg cag gat gcc gtg aac ccc atc gag agc ctg gac acc tgg Val Thr Thr Gln Asp Ala Val Asn Pro Ile Glu Ser Leu Asp Thr Trp 495 500 505	1540
acc gag ttt gtg gag gga aag aat aag act gtg agc aag ctg gtg atc Thr Glu Phe Val Glu Gly Lys Asn Lys Thr Val Ser Lys Leu Val Ile 510 515 520	1588
cag aat gcc aac gtg tct gcc atg tac aag tgt gtg gtc tcc aac aag Gln Asn Ala Asn Val Ser Ala Met Tyr Lys Cys Val Val Ser Asn Lys 525 530 535	1636
gtg ggc cag gat gag cgg ctc atc tac ttc tat gtg acc acc atc ccc Val Gly Gln Asp Glu Arg Leu Ile Tyr Phe Tyr Val Thr Thr Ile Pro 540 545 550 555	1684
gac ggc ttc acc atc gaa tcc aag cca tcc gag gag cta cta gag ggc Asp Gly Phe Thr Ile Glu Ser Lys Pro Ser Glu Glu Leu Leu Glu Gly 560 565 570	1732
cag ccg gtg ctc ctg agc tgc caa gcc gac agc tac aag tac gag cat Gln Pro Val Leu Leu Ser Cys Gln Ala Asp Ser Tyr Lys Tyr Glu His 575 580 585	1780
ctg cgc tgg tac cgc ctc aac ctg tcc acg ctg cac gat gcg cac ggg Leu Arg Trp Tyr Arg Leu Asn Leu Ser Thr Leu His Asp Ala His Gly 590 595 600	1828
aac ccg ctt ctg ctc gac tgc aag aac gtg cat ctg ttc gcc acc cct Asn Pro Leu Leu Leu Asp Cys Lys Asn Val His Leu Phe Ala Thr Pro 605 610 615	1876
ctg gcc gcc agc ctg gag gag gtg gca cct ggg gcg cgc cac gcc acg Leu Ala Ala Ser Leu Glu Val Ala Pro Gly Ala Arg His Ala Thr 620 625 630 635	1924
ctc agc ctg agt atc ccc cgc gtc gcg ccc gag cac gag ggc cac tat Leu Ser Leu Ser Ile Pro Arg Val Ala Pro Glu His Glu Gly His Tyr 640 645 650	1972
gtg tgc gaa gtg caa gac cgg cgc agc cat gac aag cac tgc cac aag Val Cys Glu Val Gln Asp Arg Arg Ser His Asp Lys His Cys His Lys 655 660 665	2020
aag tac ctg tgg gtg cag gcc ctg gaa gcc cct cgg ctc acg cag aac Lys Tyr Leu Ser Val Gln Ala Leu Glu Ala Pro Arg Leu Thr Gln Asn 670 675 680	2068
ttg acc gac ctc ctg gtg aac gtg agc gac tgg ctg gag atg cag tgc Leu Thr Asp Leu Leu Val Asn Val Ser Asp Ser Leu Glu Met Gln Cys 685 690 695	2116
ttg gtg gcc gga gcg cac gcg ccc agc atc gtg tgg tac aaa gac gag Leu Val Ala Gly Ala His Ala Pro Ser Ile Val Trp Tyr Lys Asp Glu 700 705 710 715	2164
agg ctg ctg gag gaa aag tct gga gtc gac ttg gcg gac tcc aac cag Arg Leu Leu Glu Glu Lys Ser Gly Val Asp Leu Ala Asp Ser Asn Gln 720 725 730	2212

-continued

aag ctg agc atc cag cgc gtg cgc gag gag gat gcg gga cgc tat ctg Lys Leu Ser Ile Gln Arg Val Arg Glu Glu Asp Ala Gly Arg Tyr Leu 735 740 745	2260
tgc agc gtg tgc aac gcc aag ggc tgc gtc aac tcc tcc gcc agc gtg Cys Ser Val Cys Asn Ala Lys Gly Cys Val Asn Ser Ser Ala Ser Val 750 755 760	2308
gcc gtg gaa ggc tcc gag gat aag ggc agc atg gag atc gtg atc ctt Ala Val Glu Gly Ser Glu Asp Lys Gly Ser Met Glu Ile Val Ile Leu 765 770 775	2356
gtc ggt acc ggc gtc atc gct gtc ttc ttc tgg gtc ctc ctc ctc ctc Val Gly Thr Gly Val Ile Ala Val Phe Phe Trp Val Leu Leu Leu Leu 780 785 790 795	2404
atc ttc tgt aac atg agg agg ccg gcc cac gca gac atc aag acg ggc Ile Phe Cys Asn Met Arg Arg Pro Ala His Ala Asp Ile Lys Thr Gly 800 805 810	2452
tac ctg tcc atc atc atg gac ccc ggg gag gtg cct ctg gag gag caa Tyr Leu Ser Ile Ile Met Asp Pro Gly Glu Val Pro Leu Glu Glu Gln 815 820 825	2500
tgc gaa tac ctg tcc tac gat gcc agc cag tgg gaa ttc ccc cga gag Cys Glu Tyr Leu Ser Tyr Asp Ala Ser Gln Trp Glu Phe Pro Arg Glu 830 835 840	2548
cgg ctg cac ctg ggg aga gtg ctc ggc tac ggc gcc ttc ggg aag gtg Arg Leu His Leu Gly Arg Val Leu Gly Tyr Gly Ala Phe Gly Lys Val 845 850 855	2596
gtg gaa gcc tcc gct ttc ggc atc cac aag ggc agc agc tgt gac acc Val Glu Ala Ser Ala Phe Gly Ile His Lys Gly Ser Ser Cys Asp Thr 860 865 870 875	2644
gtg gcc gtg aaa atg ctg aaa gag ggc gcc acg gcc agc gag cac cgc Val Ala Val Lys Met Leu Lys Glu Gly Ala Thr Ala Ser Glu His Arg 880 885 890	2692
gcg ctg atg tcg gag ctc aag atc ctc att cac atc ggc aac cac ctc Ala Leu Met Ser Glu Leu Lys Ile Leu Ile His Ile Gly Asn His Leu 895 900 905	2740
aac gtg gtc aac ctc ctc ggg gcg tgc acc aag ccg cag ggc ccc ctc Asn Val Val Asn Leu Leu Gly Ala Cys Thr Lys Pro Gln Gly Pro Leu 910 915 920	2788
atg gtg atc gtg gag ttc tgc aag tac ggc aac ctc tcc aac ttc ctg Met Val Ile Val Glu Phe Cys Lys Tyr Gly Asn Leu Ser Asn Phe Leu 925 930 935	2836
cgc gcc aag ccg gac gcc ttc agc ccc tgc gcg gag aag tct ccc gag Arg Ala Lys Arg Asp Ala Phe Ser Pro Cys Ala Glu Lys Ser Pro Glu 940 945 950 955	2884
cag cgc gga cgc ttc cgc gcc atg gtg gag ctc gcc agg ctg gat ccg Gln Arg Gly Arg Phe Arg Ala Met Val Glu Leu Ala Arg Leu Asp Arg 960 965 970	2932
agg ccg ccg ggg agc agc gac agg gtc ctc ttc gcg ccg ttc tcg aag Arg Arg Pro Gly Ser Ser Asp Arg Val Leu Phe Ala Arg Phe Ser Lys 975 980 985	2980
acc gag ggc gga gcg agg ccg gct tct cca gac caa gaa gct gag gac Thr Glu Gly Gly Ala Arg Arg Ala Ser Pro Asp Gln Glu Ala Glu Asp 990 995 1000	3028
ctg tgg ctg agc ccg ctg acc atg gaa gat ctt gtc tgc tac agc ttc Leu Trp Leu Ser Pro Leu Thr Met Glu Asp Leu Val Cys Tyr Ser Phe 1005 1010 1015	3076
cag gtg gcc aga ggg atg gag ttc ctg gct tcc cga aag tgc atc cac Gln Val Ala Arg Gly Met Glu Phe Leu Ala Ser Arg Lys Cys Ile His 1020 1025 1030 1035	3124
aga gac ctg gct gct ccg aac att ctg ctg tcg gaa agc gac gtg gtg Arg Asp Leu Ala Ala Arg Asn Ile Leu Leu Ser Glu Ser Asp Val Val	3172

-continued

1040	1045	1050	
aag atc tgt gac ttt ggc ctt gcc cgg gac atc tac aaa gac cct gac Lys Ile Cys Asp Phe Gly Leu Ala Arg Asp Ile Tyr Lys Asp Pro Asp 1055 1060 1065			3220
tac gtc cgc aag ggc agt gcc cgg ctg ccc ctg aag tgg atg gcc cct Tyr Val Arg Lys Gly Ser Ala Arg Leu Pro Leu Lys Trp Met Ala Pro 1070 1075 1080			3268
gaa agc atc ttc gac aag gtg tac acc acg cag agt gac gtg tgg tcc Glu Ser Ile Phe Asp Lys Val Tyr Thr Thr Gln Ser Asp Val Trp Ser 1085 1090 1095			3316
ttt ggg gtg ctt ctg tgg gag atc ttc tct ctg ggg gcc tcc ccg tac Phe Gly Val Leu Leu Trp Glu Ile Phe Ser Leu Gly Ala Ser Pro Tyr 1100 1105 1110 1115			3364
cct ggg gtg cag atc aat gag gag ttc tgc cag cgg ctg aga gac ggc Pro Gly Val Gln Ile Asn Glu Glu Phe Cys Gln Arg Leu Arg Asp Gly 1120 1125 1130			3412
aca agg atg agg gcc ccg gag ctg gcc act ccc gcc ata cgc cgc atc Thr Arg Met Arg Ala Pro Glu Leu Ala Thr Pro Ala Ile Arg Arg Ile 1135 1140 1145			3460
atg ctg aac tgc tgg tcc gga gac ccc aag gcg aga cct gca ttc tcg Met Leu Asn Cys Trp Ser Gly Asp Pro Lys Ala Arg Pro Ala Phe Ser 1150 1155 1160			3508
gag ctg gtg gag atc ctg ggg gac ctg ctg cag gcc agg ggc ctg caa Glu Leu Val Glu Ile Leu Gly Asp Leu Leu Gln Gly Arg Gly Leu Gln 1165 1170 1175			3556
gag gaa gag gag gtc tgc atg gcc ccg cgc agc tct cag agc tca gaa Glu Glu Glu Glu Val Cys Met Ala Pro Arg Ser Ser Gln Ser Ser Glu 1180 1185 1190 1195			3604
gag ggc agc ttc tcg cag gtg tcc acc atg gcc cta cac atc gcc cag Glu Gly Ser Phe Ser Gln Val Ser Thr Met Ala Leu His Ile Ala Gln 1200 1205 1210			3652
gct gac gct gag gac agc ccg cca agc ctg cag cgc cac agc ctg gcc Ala Asp Ala Glu Asp Ser Pro Pro Ser Leu Gln Arg His Ser Leu Ala 1215 1220 1225			3700
gcc agg tat tac aac tgg gtg tcc ttt ccc ggg tgc ctg gcc aga ggg Ala Arg Tyr Tyr Asn Trp Val Ser Phe Pro Gly Cys Leu Ala Arg Gly 1230 1235 1240			3748
gct gag acc cgt ggt tcc tcc agg atg aag aca ttt gag gaa ttc ccc Ala Glu Thr Arg Gly Ser Ser Arg Met Lys Thr Phe Glu Glu Phe Pro 1245 1250 1255			3796
atg acc cca acg acc tac aaa ggc tct gtg gac aac cag aca gac agt Met Thr Pro Thr Thr Tyr Lys Gly Ser Val Asp Asn Gln Thr Asp Ser 1260 1265 1270 1275			3844
ggg atg gtg ctg gcc tcg gag gag ttt gag cag ata gag agc agg cat Gly Met Val Leu Ala Ser Glu Glu Phe Glu Gln Ile Glu Ser Arg His 1280 1285 1290			3892
aga caa gaa agc ggc ttc agg tagctgaagc agagagagag aaggcagcat Arg Gln Glu Ser Gly Phe Arg 1295			3943
acgtcagcat tttcttctct gcacttataa gaaagatcaa agactttaag actttcgcta			4003
tttcttctac tgctatctac tacaaacttc aaagaggaac caggaggaca agaggagcat			4063
gaaagtggac aaggagtgtg accactgaag caccacaggg aaggggttag gcctccggat			4123
gactgcgggc aggcctggat aatatccagc ctcccacaag aagctggtgg agcagagtgt			4183
tccctgactc ct			4195

<210> SEQ ID NO 2

<211> LENGTH: 1298

-continued

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 2

```

Met Gln Arg Gly Ala Ala Leu Cys Leu Arg Leu Trp Leu Cys Leu Gly
 1             5             10             15

Leu Leu Asp Gly Leu Val Ser Gly Tyr Ser Met Thr Pro Pro Thr Leu
      20             25             30

Asn Ile Thr Glu Glu Ser His Val Ile Asp Thr Gly Asp Ser Leu Ser
      35             40             45

Ile Ser Cys Arg Gly Gln His Pro Leu Glu Trp Ala Trp Pro Gly Ala
      50             55             60

Gln Glu Ala Pro Ala Thr Gly Asp Lys Asp Ser Glu Asp Thr Gly Val
      65             70             75             80

Val Arg Asp Cys Glu Gly Thr Asp Ala Arg Pro Tyr Cys Lys Val Leu
      85             90             95

Leu Leu His Glu Val His Ala Asn Asp Thr Gly Ser Tyr Val Cys Tyr
      100            105            110

Tyr Lys Tyr Ile Lys Ala Arg Ile Glu Gly Thr Thr Ala Ala Ser Ser
      115            120            125

Tyr Val Phe Val Arg Asp Phe Glu Gln Pro Phe Ile Asn Lys Pro Asp
      130            135            140

Thr Leu Leu Val Asn Arg Lys Asp Ala Met Trp Val Pro Cys Leu Val
      145            150            155            160

Ser Ile Pro Gly Leu Asn Val Thr Leu Arg Ser Gln Ser Ser Val Leu
      165            170            175

Trp Pro Asp Gly Gln Glu Val Val Trp Asp Asp Arg Arg Gly Met Leu
      180            185            190

Val Ser Thr Pro Leu Leu His Asp Ala Leu Tyr Leu Gln Cys Glu Thr
      195            200            205

Thr Trp Gly Asp Gln Asp Phe Leu Ser Asn Pro Phe Leu Val His Ile
      210            215            220

Thr Gly Asn Glu Leu Tyr Asp Ile Gln Leu Leu Pro Arg Lys Ser Leu
      225            230            235            240

Glu Leu Leu Val Gly Glu Lys Leu Val Leu Asn Cys Thr Val Trp Ala
      245            250            255

Glu Phe Asn Ser Gly Val Thr Phe Asp Trp Asp Tyr Pro Gly Lys Gln
      260            265            270

Ala Glu Arg Gly Lys Trp Val Pro Glu Arg Arg Ser Gln Gln Thr His
      275            280            285

Thr Glu Leu Ser Ser Ile Leu Thr Ile His Asn Val Ser Gln His Asp
      290            295            300

Leu Gly Ser Tyr Val Cys Lys Ala Asn Asn Gly Ile Gln Arg Phe Arg
      305            310            315            320

Glu Ser Thr Glu Val Ile Val His Glu Asn Pro Phe Ile Ser Val Glu
      325            330            335

Trp Leu Lys Gly Pro Ile Leu Glu Ala Thr Ala Gly Asp Glu Leu Val
      340            345            350

Lys Leu Pro Val Lys Leu Ala Ala Tyr Pro Pro Pro Glu Phe Gln Trp
      355            360            365

Tyr Lys Asp Gly Lys Ala Leu Ser Gly Arg His Ser Pro His Ala Leu
      370            375            380

Val Leu Lys Glu Val Thr Glu Ala Ser Thr Gly Thr Tyr Thr Leu Ala
      385            390            395            400

```

-continued

Leu Trp Asn Ser Ala Ala Gly Leu Arg Arg Asn Ile Ser Leu Glu Leu
 405 410 415
 Val Val Asn Val Pro Pro Gln Ile His Glu Lys Glu Ala Ser Ser Pro
 420 425 430
 Ser Ile Tyr Ser Arg His Ser Arg Gln Ala Leu Thr Cys Thr Ala Tyr
 435 440 445
 Gly Val Pro Leu Pro Leu Ser Ile Gln Trp His Trp Arg Pro Trp Thr
 450 455 460
 Pro Cys Lys Met Phe Ala Gln Arg Ser Leu Arg Arg Arg Gln Gln Gln
 465 470 475 480
 Asp Leu Met Pro Gln Cys Arg Asp Trp Arg Ala Val Thr Thr Gln Asp
 485 490 495
 Ala Val Asn Pro Ile Glu Ser Leu Asp Thr Trp Thr Glu Phe Val Glu
 500 505 510
 Gly Lys Asn Lys Thr Val Ser Lys Leu Val Ile Gln Asn Ala Asn Val
 515 520 525
 Ser Ala Met Tyr Lys Cys Val Val Ser Asn Lys Val Gly Gln Asp Glu
 530 535 540
 Arg Leu Ile Tyr Phe Tyr Val Thr Thr Ile Pro Asp Gly Phe Thr Ile
 545 550 555 560
 Glu Ser Lys Pro Ser Glu Glu Leu Leu Glu Gly Gln Pro Val Leu Leu
 565 570 575
 Ser Cys Gln Ala Asp Ser Tyr Lys Tyr Glu His Leu Arg Trp Tyr Arg
 580 585 590
 Leu Asn Leu Ser Thr Leu His Asp Ala His Gly Asn Pro Leu Leu Leu
 595 600 605
 Asp Cys Lys Asn Val His Leu Phe Ala Thr Pro Leu Ala Ala Ser Leu
 610 615 620
 Glu Glu Val Ala Pro Gly Ala Arg His Ala Thr Leu Ser Leu Ser Ile
 625 630 635 640
 Pro Arg Val Ala Pro Glu His Glu Gly His Tyr Val Cys Glu Val Gln
 645 650 655
 Asp Arg Arg Ser His Asp Lys His Cys His Lys Lys Tyr Leu Ser Val
 660 665 670
 Gln Ala Leu Glu Ala Pro Arg Leu Thr Gln Asn Leu Thr Asp Leu Leu
 675 680 685
 Val Asn Val Ser Asp Ser Leu Glu Met Gln Cys Leu Val Ala Gly Ala
 690 695 700
 His Ala Pro Ser Ile Val Trp Tyr Lys Asp Glu Arg Leu Leu Glu Glu
 705 710 715 720
 Lys Ser Gly Val Asp Leu Ala Asp Ser Asn Gln Lys Leu Ser Ile Gln
 725 730 735
 Arg Val Arg Glu Glu Asp Ala Gly Arg Tyr Leu Cys Ser Val Cys Asn
 740 745 750
 Ala Lys Gly Cys Val Asn Ser Ser Ala Ser Val Ala Val Glu Gly Ser
 755 760 765
 Glu Asp Lys Gly Ser Met Glu Ile Val Ile Leu Val Gly Thr Gly Val
 770 775 780
 Ile Ala Val Phe Phe Trp Val Leu Leu Leu Leu Ile Phe Cys Asn Met
 785 790 795 800
 Arg Arg Pro Ala His Ala Asp Ile Lys Thr Gly Tyr Leu Ser Ile Ile
 805 810 815

-continued

Met Asp Pro Gly Glu Val Pro Leu Glu Glu Gln Cys Glu Tyr Leu Ser	820	825	830
Tyr Asp Ala Ser Gln Trp Glu Phe Pro Arg Glu Arg Leu His Leu Gly	835	840	845
Arg Val Leu Gly Tyr Gly Ala Phe Gly Lys Val Val Glu Ala Ser Ala	850	855	860
Phe Gly Ile His Lys Gly Ser Ser Cys Asp Thr Val Ala Val Lys Met	865	870	875
Leu Lys Glu Gly Ala Thr Ala Ser Glu His Arg Ala Leu Met Ser Glu	885	890	895
Leu Lys Ile Leu Ile His Ile Gly Asn His Leu Asn Val Val Asn Leu	900	905	910
Leu Gly Ala Cys Thr Lys Pro Gln Gly Pro Leu Met Val Ile Val Glu	915	920	925
Phe Cys Lys Tyr Gly Asn Leu Ser Asn Phe Leu Arg Ala Lys Arg Asp	930	935	940
Ala Phe Ser Pro Cys Ala Glu Lys Ser Pro Glu Gln Arg Gly Arg Phe	945	950	955
Arg Ala Met Val Glu Leu Ala Arg Leu Asp Arg Arg Arg Pro Gly Ser	965	970	975
Ser Asp Arg Val Leu Phe Ala Arg Phe Ser Lys Thr Glu Gly Gly Ala	980	985	990
Arg Arg Ala Ser Pro Asp Gln Glu Ala Glu Asp Leu Trp Leu Ser Pro	995	1000	1005
Leu Thr Met Glu Asp Leu Val Cys Tyr Ser Phe Gln Val Ala Arg Gly	1010	1015	1020
Met Glu Phe Leu Ala Ser Arg Lys Cys Ile His Arg Asp Leu Ala Ala	1025	1030	1035
Arg Asn Ile Leu Leu Ser Glu Ser Asp Val Val Lys Ile Cys Asp Phe	1045	1050	1055
Gly Leu Ala Arg Asp Ile Tyr Lys Asp Pro Asp Tyr Val Arg Lys Gly	1060	1065	1070
Ser Ala Arg Leu Pro Leu Lys Trp Met Ala Pro Glu Ser Ile Phe Asp	1075	1080	1085
Lys Val Tyr Thr Thr Gln Ser Asp Val Trp Ser Phe Gly Val Leu Leu	1090	1095	1100
Trp Glu Ile Phe Ser Leu Gly Ala Ser Pro Tyr Pro Gly Val Gln Ile	1105	1110	1115
Asn Glu Glu Phe Cys Gln Arg Leu Arg Asp Gly Thr Arg Met Arg Ala	1125	1130	1135
Pro Glu Leu Ala Thr Pro Ala Ile Arg Arg Ile Met Leu Asn Cys Trp	1140	1145	1150
Ser Gly Asp Pro Lys Ala Arg Pro Ala Phe Ser Glu Leu Val Glu Ile	1155	1160	1165
Leu Gly Asp Leu Leu Gln Gly Arg Gly Leu Gln Glu Glu Glu Val	1170	1175	1180
Cys Met Ala Pro Arg Ser Ser Gln Ser Ser Glu Glu Gly Ser Phe Ser	1185	1190	1195
Gln Val Ser Thr Met Ala Leu His Ile Ala Gln Ala Asp Ala Glu Asp	1205	1210	1215
Ser Pro Pro Ser Leu Gln Arg His Ser Leu Ala Ala Arg Tyr Tyr Asn	1220	1225	1230
Trp Val Ser Phe Pro Gly Cys Leu Ala Arg Gly Ala Glu Thr Arg Gly			

-continued

1235	1240	1245	
Ser Ser Arg Met Lys Thr Phe Glu Glu Phe Pro Met Thr Pro Thr Thr			
1250	1255	1260	
Tyr Lys Gly Ser Val Asp Asn Gln Thr Asp Ser Gly Met Val Leu Ala			
265	1270	1275	1280
Ser Glu Glu Phe Glu Gln Ile Glu Ser Arg His Arg Gln Glu Ser Gly			
1285	1290	1295	
Phe Arg			
<210> SEQ ID NO 3			
<211> LENGTH: 4795			
<212> TYPE: DNA			
<213> ORGANISM: Homo sapiens			
<220> FEATURE:			
<221> NAME/KEY: CDS			
<222> LOCATION: (20)..(4108)			
<400> SEQUENCE: 3			
ccacgcgcag cggccggag atg cag cgg ggc gcc gcg ctg tgc ctg cga ctg		52	
Met Gln Arg Gly Ala Ala Leu Cys Leu Arg Leu			
1 5 10			
tgg ctg tgc ctg gga ctc ctg gac ggc ctg gtg agt ggc tac tcc atg		100	
Trp Leu Cys Leu Gly Leu Leu Asp Gly Leu Val Ser Gly Tyr Ser Met			
15 20 25			
acc ccc ccg acc ttg aac atc acg gag gag tca cac gtc atc gac acc		148	
Thr Pro Pro Thr Thr Leu Asn Ile Thr Glu Glu Ser His Val Ile Asp Thr			
30 35 40			
ggt gac agc ctg tcc atc tcc tgc agg gga cag cac ccc ctg gag tgg		196	
Gly Asp Ser Leu Ser Ile Ser Cys Arg Gly Gln His Pro Leu Glu Trp			
45 50 55			
gct tgg cca gga gct cag gag gcg cca gcc acc gga gac aag gac agc		244	
Ala Trp Pro Gly Ala Gln Glu Ala Pro Ala Thr Gly Asp Lys Asp Ser			
60 65 70 75			
gag gac acg ggg gtg gtg cga gac tgc gag ggc aca gac gcc agg ccc		292	
Glu Asp Thr Gly Val Val Arg Asp Cys Glu Gly Thr Asp Ala Arg Pro			
80 85 90			
tac tgc aag gtg ttg ctg ctg cac gag gta cat gcc aac gac aca ggc		340	
Tyr Cys Lys Val Leu Leu Leu His Glu Val His Ala Asn Asp Thr Gly			
95 100 105			
agc tac gtc tgc tac tac aag tac atc aag gca cgc atc gag ggc acc		388	
Ser Tyr Val Cys Tyr Tyr Lys Tyr Ile Lys Ala Arg Ile Glu Gly Thr			
110 115 120			
acg gcc gcc agc tcc tac gtg ttc gtg aga gac ttt gag cag cca ttc		436	
Thr Ala Ala Ser Ser Tyr Val Phe Val Arg Asp Phe Glu Gln Pro Phe			
125 130 135			
atc aac aag cct gac acg ctc ttg gtc aac agg aag gac gcc atg tgg		484	
Ile Asn Lys Pro Asp Thr Leu Leu Val Asn Arg Lys Asp Ala Met Trp			
140 145 150 155			
gtg ccc tgt ctg gtg tcc atc ccc ggc ctc aat gtc acg ctg cgc tcg		532	
Val Pro Cys Leu Val Ser Ile Pro Gly Leu Asn Val Thr Leu Arg Ser			
160 165 170			
caa agc tcg gtg ctg tgg cca gac ggg cag gag gtg gtg tgg gat gac		580	
Gln Ser Ser Val Leu Trp Pro Asp Gly Gln Glu Val Val Trp Asp Asp			
175 180 185			
cgg cgg ggc atg ctc gtg tcc acg cca ctg ctg cac gat gcc ctg tac		628	
Arg Arg Gly Met Leu Val Ser Thr Pro Leu Leu His Asp Ala Leu Tyr			
190 195 200			
ctg cag tgc gag acc acc tgg gga gac cag gac ttc ctt tcc aac ccc		676	
Leu Gln Cys Glu Thr Thr Trp Gly Asp Gln Asp Phe Leu Ser Asn Pro			
205 210 215			

-continued

ttc ctg gtg cac atc aca ggc aac gag ctc tat gac atc cag ctg ttg	724
Phe Leu Val His Ile Thr Gly Asn Glu Leu Tyr Asp Ile Gln Leu Leu	
220 225 230 235	
ccc agg aag tcg ctg gag ctg ctg gta ggg gag aag ctg gtc ctg aac	772
Pro Arg Lys Ser Leu Glu Leu Leu Val Gly Glu Lys Leu Val Leu Asn	
240 245 250	
tgc acc gtg tgg gct gag ttt aac tca ggt gtc acc ttt gac tgg gac	820
Cys Thr Val Trp Ala Glu Phe Asn Ser Gly Val Thr Phe Asp Trp Asp	
255 260 265	
tac cca ggg aag cag gca gag cgg ggt aag tgg gtg ccc gag cga cgc	868
Tyr Pro Gly Lys Gln Ala Glu Arg Gly Lys Trp Val Pro Glu Arg Arg	
270 275 280	
tcc cag cag acc cac aca gaa ctc tcc agc atc ctg acc atc cac aac	916
Ser Gln Gln Thr His Thr Glu Leu Ser Ser Ile Leu Thr Ile His Asn	
285 290 295	
gtc agc cag cac gac ctg ggc tcg tat gtg tgc aag gcc aac aac ggc	964
Val Ser Gln His Asp Leu Gly Ser Tyr Val Cys Lys Ala Asn Asn Gly	
300 305 310 315	
atc cag cga ttt cgg gag agc acc gag gtc att gtg cat gaa aat ccc	1012
Ile Gln Arg Phe Arg Glu Ser Thr Glu Val Ile Val His Glu Asn Pro	
320 325 330	
ttc atc agc gtc gag tgg ctc aaa gga ccc atc ctg gag gcc acg gca	1060
Phe Ile Ser Val Glu Trp Leu Lys Gly Pro Ile Leu Glu Ala Thr Ala	
335 340 345	
gga gac gag ctg gtg aag ctg ccc gtg aag ctg gca gcg tac ccc ccg	1108
Gly Asp Glu Leu Val Lys Leu Pro Val Lys Leu Ala Ala Tyr Pro Pro	
350 355 360	
ccc gag ttc cag tgg tac aag gat gga aag gca ctg tcc ggg cgc cac	1156
Pro Glu Phe Gln Trp Tyr Lys Asp Gly Lys Ala Leu Ser Gly Arg His	
365 370 375	
agt cca cat gcc ctg gtg ctc aag gag gtg aca gag gcc agc aca ggc	1204
Ser Pro His Ala Leu Val Leu Lys Glu Val Thr Glu Ala Ser Thr Gly	
380 385 390 395	
acc tac acc ctc gcc ctg tgg aac tcc gct gct gcc ctg agg cgc aac	1252
Thr Tyr Thr Leu Ala Leu Trp Asn Ser Ala Ala Gly Leu Arg Arg Asn	
400 405 410	
atc agc ctg gag ctg gtg gtg aat gtg ccc ccc cag ata cat gag aag	1300
Ile Ser Leu Glu Leu Val Val Asn Val Pro Pro Gln Ile His Glu Lys	
415 420 425	
gag gcc tcc tcc ccc agc atc tac tcg cgt cac agc cgc cag gcc ctc	1348
Glu Ala Ser Ser Pro Ser Ile Tyr Ser Arg His Ser Arg Gln Ala Leu	
430 435 440	
acc tgc acg gcc tac ggg gtg ccc ctg cct ctc agc atc cag tgg cac	1396
Thr Cys Thr Ala Tyr Gly Val Pro Leu Pro Leu Ser Ile Gln Trp His	
445 450 455	
tgg cgg ccc tgg aca ccc tgc aag atg ttt gcc cag cgt agt ctc cgg	1444
Trp Arg Pro Trp Thr Pro Cys Lys Met Phe Ala Gln Arg Ser Leu Arg	
460 465 470 475	
cgg cgg cag cag caa gac ctc atg cca cag tgc cgt gac tgg agg gcg	1492
Arg Arg Gln Gln Asp Leu Met Pro Gln Cys Arg Asp Trp Arg Ala	
480 485 490	
gtg acc acg cag gat gcc gtg aac ccc atc gag agc ctg gac acc tgg	1540
Val Thr Thr Gln Asp Ala Val Asn Pro Ile Glu Ser Leu Asp Thr Trp	
495 500 505	
acc gag ttt gtg gag gga aag aat aag act gtg agc aag ctg gtg atc	1588
Thr Glu Phe Val Glu Gly Lys Asn Lys Thr Val Ser Lys Leu Val Ile	
510 515 520	
cag aat gcc aac gtg tct gcc atg tac aag tgt gtg gtc tcc aac aag	1636
Gln Asn Ala Asn Val Ser Ala Met Tyr Lys Cys Val Val Ser Asn Lys	

-continued

525	530	535	
gtg ggc cag gat gag cgg ctc atc tac ttc tat gtg acc acc atc ccc Val Gly Gln Asp Glu Arg Leu Ile Tyr Phe Tyr Val Thr Thr Ile Pro 540 545 550 555			1684
gac ggc ttc acc atc gaa tcc aag cca tcc gag gag cta cta gag ggc Asp Gly Phe Thr Ile Glu Ser Lys Pro Ser Glu Glu Leu Leu Glu Gly 560 565 570			1732
cag ccg gtg ctc ctg agc tgc caa gcc gac agc tac aag tac gag cat Gln Pro Val Leu Leu Ser Cys Gln Ala Asp Ser Tyr Lys Tyr Glu His 575 580 585			1780
ctg cgc tgg tac cgc ctc aac ctg tcc acg ctg cac gat gcg cac ggg Leu Arg Trp Tyr Arg Leu Asn Leu Ser Thr Leu His Asp Ala His Gly 590 595 600			1828
aac ccg ctt ctg ctc gac tgc aag aac gtg cat ctg ttc gcc acc cct Asn Pro Leu Leu Leu Asp Cys Lys Asn Val His Leu Phe Ala Thr Pro 605 610 615			1876
ctg gcc gcc agc ctg gag gag gtg gca cct ggg gcg cgc cac gcc acg Leu Ala Ala Ser Leu Glu Glu Val Ala Pro Gly Ala Arg His Ala Thr 620 625 630 635			1924
ctc agc ctg agt atc ccc cgc gtc gcg ccc gag cac gag ggc cac tat Leu Ser Leu Ser Ile Pro Arg Val Ala Pro Glu His Glu Gly His Tyr 640 645 650			1972
gtg tgc gaa gtg caa gac cgg cgc agc cat gac aag cac tgc cac aag Val Cys Glu Val Gln Asp Arg Arg Ser His Asp Lys His Cys His Lys 655 660 665			2020
aag tac ctg tcg gtg cag gcc ctg gaa gcc cct cgg ctc acg cag aac Lys Tyr Leu Ser Val Gln Ala Leu Glu Ala Pro Arg Leu Thr Gln Asn 670 675 680			2068
ttg acc gac ctc ctg gtg aac gtg agc gac tcg ctg gag atg cag tgc Leu Thr Asp Leu Leu Val Asn Val Ser Asp Ser Leu Glu Met Gln Cys 685 690 695			2116
ttg gtg gcc gga gcg cac gcg ccc agc atc gtg tgg tac aaa gac gag Leu Val Ala Gly Ala His Ala Pro Ser Ile Val Trp Tyr Lys Asp Glu 700 705 710 715			2164
agg ctg ctg gag gaa aag tct gga gtc gac ttg gcg gac tcc aac cag Arg Leu Leu Glu Glu Lys Ser Gly Val Asp Leu Ala Asp Ser Asn Gln 720 725 730			2212
aag ctg agc atc cag cgc gtg cgc gag gag gat gcg gga cgc tat ctg Lys Leu Ser Ile Gln Arg Val Arg Glu Glu Asp Ala Gly Arg Tyr Leu 735 740 745			2260
tgc agc gtg tgc aac gcc aag ggc tgc gtc aac tcc tcc gcc agc gtg Cys Ser Val Cys Asn Ala Lys Gly Cys Val Asn Ser Ser Ala Ser Val 750 755 760			2308
gcc gtg gaa ggc tcc gag gat aag ggc agc atg gag atc gtg atc ctt Ala Val Glu Gly Ser Glu Asp Lys Gly Ser Met Glu Ile Val Ile Leu 765 770 775			2356
gtc ggt acc ggc gtc atc gct gtc ttc ttc tgg gtc ctc ctc ctc ctc Val Gly Thr Gly Val Ile Ala Val Phe Phe Trp Val Leu Leu Leu Leu 780 785 790 795			2404
atc ttc tgt aac atg agg agg ccg gcc cac gca gac atc aag acg ggc Ile Phe Cys Asn Met Arg Arg Pro Ala His Ala Asp Ile Lys Thr Gly 800 805 810			2452
tac ctg tcc atc atc atg gac ccc ggg gag gtg cct ctg gag gag caa Tyr Leu Ser Ile Ile Met Asp Pro Gly Glu Val Pro Leu Glu Glu Gln 815 820 825			2500
tgc gaa tac ctg tcc tac gat gcc agc cag tgg gaa ttc ccc cga gag Cys Glu Tyr Leu Ser Tyr Asp Ala Ser Gln Trp Glu Phe Pro Arg Glu 830 835 840			2548
cgg ctg cac ctg ggg aga gtg ctc gcc tac gcc gcc ttc ggg aag gtg			2596

-continued

Arg	Leu	His	Leu	Gly	Arg	Val	Leu	Gly	Tyr	Gly	Ala	Phe	Gly	Lys	Val	
845						850					855					
gtg	gaa	gcc	tcc	gct	ttc	ggc	atc	cac	aag	ggc	agc	agc	tgt	gac	acc	2644
Val	Glu	Ala	Ser	Ala	Phe	Gly	Ile	His	Lys	Gly	Ser	Ser	Cys	Asp	Thr	
860					865					870					875	
gtg	gcc	gtg	aaa	atg	ctg	aaa	gag	ggc	gcc	acg	gcc	agc	gag	cac	cgc	2692
Val	Ala	Val	Lys	Met	Leu	Lys	Glu	Gly	Ala	Thr	Ala	Ser	Glu	His	Arg	
			880						885					890		
gcg	ctg	atg	tcg	gag	ctc	aag	atc	ctc	att	cac	atc	ggc	aac	cac	ctc	2740
Ala	Leu	Met	Ser	Glu	Leu	Lys	Ile	Leu	Ile	His	Ile	Gly	Asn	His	Leu	
			895					900					905			
aac	gtg	gtc	aac	ctc	ctc	ggg	gcg	tgc	acc	aag	ccg	cag	ggc	ccc	ctc	2788
Asn	Val	Val	Asn	Leu	Leu	Gly	Ala	Cys	Thr	Lys	Pro	Gln	Gly	Pro	Leu	
	910					915						920				
atg	gtg	atc	gtg	gag	ttc	tgc	aag	tac	ggc	aac	ctc	tcc	aac	ttc	ctg	2836
Met	Val	Ile	Val	Glu	Phe	Cys	Lys	Tyr	Gly	Asn	Leu	Ser	Asn	Phe	Leu	
	925					930					935					
cgc	gcc	aag	cgg	gac	gcc	ttc	agc	ccc	tgc	gcg	gag	aag	tct	ccc	gag	2884
Arg	Ala	Lys	Arg	Asp	Ala	Phe	Ser	Pro	Cys	Ala	Glu	Lys	Ser	Pro	Glu	
940					945				950						955	
cag	cgc	gga	cgc	ttc	cgc	gcc	atg	gtg	gag	ctc	gcc	agg	ctg	gat	cgg	2932
Gln	Arg	Gly	Arg	Phe	Arg	Ala	Met	Val	Glu	Leu	Ala	Arg	Leu	Asp	Arg	
			960						965					970		
agg	cgg	ccg	ggg	agc	agc	gac	agg	gtc	ctc	ttc	gcg	cgg	ttc	tcg	aag	2980
Arg	Arg	Pro	Gly	Ser	Ser	Asp	Arg	Val	Leu	Phe	Ala	Arg	Phe	Ser	Lys	
			975					980					985			
acc	gag	ggc	gga	gcg	agg	cgg	gct	tct	cca	gac	caa	gaa	gct	gag	gac	3028
Thr	Glu	Gly	Gly	Ala	Arg	Arg	Ala	Ser	Pro	Asp	Gln	Glu	Ala	Glu	Asp	
	990					995					1000					
ctg	tgg	ctg	agc	ccg	ctg	acc	atg	gaa	gat	ctt	gtc	tgc	tac	agc	ttc	3076
Leu	Trp	Leu	Ser	Pro	Leu	Thr	Met	Glu	Asp	Leu	Val	Cys	Tyr	Ser	Phe	
	1005					1010					1015					
cag	gtg	gcc	aga	ggg	atg	gag	ttc	ctg	gct	tcc	cga	aag	tgc	atc	cac	3124
Gln	Val	Ala	Arg	Gly	Met	Glu	Phe	Leu	Ala	Ser	Arg	Lys	Cys	Ile	His	
1020					1025					1030					1035	
aga	gac	ctg	gct	gct	cgg	aac	att	ctg	ctg	tcg	gaa	agc	gac	gtg	gtg	3172
Arg	Asp	Leu	Ala	Ala	Arg	Asn	Ile	Leu	Leu	Ser	Glu	Ser	Asp	Val	Val	
			1040						1045					1050		
aag	atc	tgt	gac	ttt	ggc	ctt	gcc	cgg	gac	atc	tac	aaa	gac	cct	gac	3220
Lys	Ile	Cys	Asp	Phe	Gly	Leu	Ala	Arg	Asp	Ile	Tyr	Lys	Asp	Pro	Asp	
		1055						1060					1065			
tac	gtc	cgc	aag	ggc	agt	gcc	cgg	ctg	ccc	ctg	aag	tgg	atg	gcc	cct	3268
Tyr	Val	Arg	Lys	Gly	Ser	Ala	Arg	Leu	Pro	Leu	Lys	Trp	Met	Ala	Pro	
	1070					1075						1080				
gaa	agc	atc	ttc	gac	aag	gtg	tac	acc	acg	cag	agt	gac	gtg	tgg	tcc	3316
Glu	Ser	Ile	Phe	Asp	Lys	Val	Tyr	Thr	Thr	Gln	Ser	Asp	Val	Trp	Ser	
	1085					1090					1095					
ttt	ggg	gtg	ctt	ctc	tgg	gag	atc	ttc	tct	ctg	ggg	gcc	tcc	ccg	tac	3364
Phe	Gly	Val	Leu	Leu	Trp	Glu	Ile	Phe	Ser	Leu	Gly	Ala	Ser	Pro	Tyr	
1100					1105					1110					1115	
cct	ggg	gtg	cag	atc	aat	gag	gag	ttc	tgc	cag	cgg	ctg	aga	gac	ggc	3412
Pro	Gly	Val	Gln	Ile	Asn	Glu	Glu	Phe	Cys	Gln	Arg	Leu	Arg	Asp	Gly	
			1120						1125					1130		
aca	agg	atg	agg	gcc	ccg	gag	ctg	gcc	act	ccc	gcc	ata	cgc	cgc	atc	3460
Thr	Arg	Met	Arg	Ala	Pro	Glu	Leu	Ala	Thr	Pro	Ala	Ile	Arg	Arg	Ile	
			1135					1140					1145			
atg	ctg	aac	tgc	tgg	tcc	gga	gac	ccc	aag	gcg	aga	cct	gca	ttc	tcg	3508
Met	Leu	Asn	Cys	Trp	Ser	Gly	Asp	Pro	Lys	Ala	Arg	Pro	Ala	Phe	Ser	
	1150						1155						1160			

-continued

gag ctg gtg gag atc ctg ggg gac ctg ctc cag ggc agg ggc ctg caa Glu Leu Val Glu Ile Leu Gly Asp Leu Leu Gln Gly Arg Gly Leu Gln 1165 1170 1175	3556
gag gaa gag gag gtc tgc atg gcc ccg cgc agc tct cag agc tca gaa Glu Glu Glu Glu Val Cys Met Ala Pro Arg Ser Ser Gln Ser Ser Glu 1180 1185 1190 1195	3604
gag ggc agc ttc tcg cag gtg tcc acc atg gcc cta cac atc gcc cag Glu Gly Ser Phe Ser Gln Val Ser Thr Met Ala Leu His Ile Ala Gln 1200 1205 1210	3652
gct gac gct gag gac agc ccg cca agc ctg cag cgc cac agc ctg gcc Ala Asp Ala Glu Asp Ser Pro Pro Ser Leu Gln Arg His Ser Leu Ala 1215 1220 1225	3700
gcc agg tat tac aac tgg gtg tcc ttt ccc ggg tgc ctg gcc aga ggg Ala Arg Tyr Tyr Asn Trp Val Ser Phe Pro Gly Cys Leu Ala Arg Gly 1230 1235 1240	3748
gct gag acc cgt ggt tcc tcc agg atg aag aca ttt gag gaa ttc ccc Ala Glu Thr Arg Gly Ser Ser Arg Met Lys Thr Phe Glu Glu Phe Pro 1245 1250 1255	3796
atg acc cca acg acc tac aaa ggc tct gtg gac aac cag aca gac agt Met Thr Pro Thr Thr Tyr Lys Gly Ser Val Asp Asn Gln Thr Asp Ser 1260 1265 1270 1275	3844
ggg atg gtg ctg gcc tcg gag gag ttt gag cag ata gag agc agg cat Gly Met Val Leu Ala Ser Glu Glu Phe Glu Gln Ile Glu Ser Arg His 1280 1285 1290	3892
aga caa gaa agc ggc ttc agc tgt aaa gga cct ggc cag aat gtg gct Arg Gln Glu Ser Gly Phe Ser Cys Lys Gly Pro Gly Gln Asn Val Ala 1295 1300 1305	3940
gtg acc agg gca cac cct gac tcc caa ggg agg cgg cgg cgt gag Val Thr Arg Ala His Pro Asp Ser Gln Gly Arg Arg Arg Arg Pro Glu 1310 1315 1320	3988
cgg ggg gcc cga gga ggc cag gtg ttt tac aac agc gag tat ggg gag Arg Gly Ala Arg Gly Gly Gln Val Phe Tyr Asn Ser Glu Tyr Gly Glu 1325 1330 1335	4036
ctg tcg gag cca agc gag gag gac cac tgc tcc ccg tct gcc cgc gtg Leu Ser Glu Pro Ser Glu Glu Asp His Cys Ser Pro Ser Ala Arg Val 1340 1345 1350 1355	4084
act ttc ttc aca gac aac agc tac taagcagcat cggacaagac cccagcact Thr Phe Phe Thr Asp Asn Ser Tyr 1360	4138
tggggggttca ggcccggcag ggcgggcaga gggctggagg cccaggctgg gaactcatct	4198
gggtgaactc tgggtggcaca ggagtgtcct cttccctctc tgcagacttc ccagctagga	4258
agagcaggac tccaggccca aggcctcccg aattccgtca ccacgactgg ccagggcacg	4318
ctccagctgc cccggccctt ccccttgaga ttcagatgtc atttagttca gcattccgag	4378
gtgctggtcc cggggccagc acttccatgg gaatgtctct ttggcgacct cttttcatca	4438
cactgggtgg tggcctggtc cctgttttcc cagaggaat ctgtgggtct gggagtcaca	4498
cagtgttgga ggttaaggca tacgagagca gaggtctccc aaacgcctt tcctctcag	4558
gcacacagct actctcccca cgagggttg ctggcctcac ccacccctgc acagttgaag	4618
ggaggggctg tgtttccatc tcaaagaagg catttgacg gtctctttct gggcctgacc	4678
aaacagccaa ctagccctg ggggtggcac cagtatgaca gtattatacg ctggcaacac	4738
agaggcagcc cgcacacctg cgctgggtg ttgagagcca tcctgcaagt ctttttc	4795

<210> SEQ ID NO 4

<211> LENGTH: 1363

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

-continued

<400> SEQUENCE: 4

Met Gln Arg Gly Ala Ala Leu Cys Leu Arg Leu Trp Leu Cys Leu Gly
 1 5 10 15
 Leu Leu Asp Gly Leu Val Ser Gly Tyr Ser Met Thr Pro Pro Thr Leu
 20 25 30
 Asn Ile Thr Glu Glu Ser His Val Ile Asp Thr Gly Asp Ser Leu Ser
 35 40 45
 Ile Ser Cys Arg Gly Gln His Pro Leu Glu Trp Ala Trp Pro Gly Ala
 50 55 60
 Gln Glu Ala Pro Ala Thr Gly Asp Lys Asp Ser Glu Asp Thr Gly Val
 65 70 75 80
 Val Arg Asp Cys Glu Gly Thr Asp Ala Arg Pro Tyr Cys Lys Val Leu
 85 90 95
 Leu Leu His Glu Val His Ala Asn Asp Thr Gly Ser Tyr Val Cys Tyr
 100 105 110
 Tyr Lys Tyr Ile Lys Ala Arg Ile Glu Gly Thr Thr Ala Ala Ser Ser
 115 120 125
 Tyr Val Phe Val Arg Asp Phe Glu Gln Pro Phe Ile Asn Lys Pro Asp
 130 135 140
 Thr Leu Leu Val Asn Arg Lys Asp Ala Met Trp Val Pro Cys Leu Val
 145 150 155 160
 Ser Ile Pro Gly Leu Asn Val Thr Leu Arg Ser Gln Ser Ser Val Leu
 165 170 175
 Trp Pro Asp Gly Gln Glu Val Val Trp Asp Asp Arg Arg Gly Met Leu
 180 185 190
 Val Ser Thr Pro Leu Leu His Asp Ala Leu Tyr Leu Gln Cys Glu Thr
 195 200 205
 Thr Trp Gly Asp Gln Asp Phe Leu Ser Asn Pro Phe Leu Val His Ile
 210 215 220
 Thr Gly Asn Glu Leu Tyr Asp Ile Gln Leu Leu Pro Arg Lys Ser Leu
 225 230 235 240
 Glu Leu Leu Val Gly Glu Lys Leu Val Leu Asn Cys Thr Val Trp Ala
 245 250 255
 Glu Phe Asn Ser Gly Val Thr Phe Asp Trp Asp Tyr Pro Gly Lys Gln
 260 265 270
 Ala Glu Arg Gly Lys Trp Val Pro Glu Arg Arg Ser Gln Gln Thr His
 275 280 285
 Thr Glu Leu Ser Ser Ile Leu Thr Ile His Asn Val Ser Gln His Asp
 290 295 300
 Leu Gly Ser Tyr Val Cys Lys Ala Asn Asn Gly Ile Gln Arg Phe Arg
 305 310 315 320
 Glu Ser Thr Glu Val Ile Val His Glu Asn Pro Phe Ile Ser Val Glu
 325 330 335
 Trp Leu Lys Gly Pro Ile Leu Glu Ala Thr Ala Gly Asp Glu Leu Val
 340 345 350
 Lys Leu Pro Val Lys Leu Ala Ala Tyr Pro Pro Pro Glu Phe Gln Trp
 355 360 365
 Tyr Lys Asp Gly Lys Ala Leu Ser Gly Arg His Ser Pro His Ala Leu
 370 375 380
 Val Leu Lys Glu Val Thr Glu Ala Ser Thr Gly Thr Tyr Thr Leu Ala
 385 390 395 400
 Leu Trp Asn Ser Ala Ala Gly Leu Arg Arg Asn Ile Ser Leu Glu Leu

-continued

405					410					415					
Val	Val	Asn	Val	Pro	Pro	Gln	Ile	His	Glu	Lys	Glu	Ala	Ser	Ser	Pro
			420						425					430	
Ser	Ile	Tyr	Ser	Arg	His	Ser	Arg	Gln	Ala	Leu	Thr	Cys	Thr	Ala	Tyr
		435					440						445		
Gly	Val	Pro	Leu	Pro	Leu	Ser	Ile	Gln	Trp	His	Trp	Arg	Pro	Trp	Thr
	450					455						460			
Pro	Cys	Lys	Met	Phe	Ala	Gln	Arg	Ser	Leu	Arg	Arg	Arg	Gln	Gln	Gln
	465					470					475				480
Asp	Leu	Met	Pro	Gln	Cys	Arg	Asp	Trp	Arg	Ala	Val	Thr	Thr	Gln	Asp
				485					490					495	
Ala	Val	Asn	Pro	Ile	Glu	Ser	Leu	Asp	Thr	Trp	Thr	Glu	Phe	Val	Glu
			500					505					510		
Gly	Lys	Asn	Lys	Thr	Val	Ser	Lys	Leu	Val	Ile	Gln	Asn	Ala	Asn	Val
		515					520						525		
Ser	Ala	Met	Tyr	Lys	Cys	Val	Val	Ser	Asn	Lys	Val	Gly	Gln	Asp	Glu
	530						535					540			
Arg	Leu	Ile	Tyr	Phe	Tyr	Val	Thr	Thr	Ile	Pro	Asp	Gly	Phe	Thr	Ile
	545					550					555				560
Glu	Ser	Lys	Pro	Ser	Glu	Glu	Leu	Leu	Glu	Gly	Gln	Pro	Val	Leu	Leu
			565						570					575	
Ser	Cys	Gln	Ala	Asp	Ser	Tyr	Lys	Tyr	Glu	His	Leu	Arg	Trp	Tyr	Arg
			580				585						590		
Leu	Asn	Leu	Ser	Thr	Leu	His	Asp	Ala	His	Gly	Asn	Pro	Leu	Leu	Leu
	595						600					605			
Asp	Cys	Lys	Asn	Val	His	Leu	Phe	Ala	Thr	Pro	Leu	Ala	Ala	Ser	Leu
	610						615					620			
Glu	Glu	Val	Ala	Pro	Gly	Ala	Arg	His	Ala	Thr	Leu	Ser	Leu	Ser	Ile
	625					630					635				640
Pro	Arg	Val	Ala	Pro	Glu	His	Glu	Gly	His	Tyr	Val	Cys	Glu	Val	Gln
			645						650					655	
Asp	Arg	Arg	Ser	His	Asp	Lys	His	Cys	His	Lys	Lys	Tyr	Leu	Ser	Val
			660					665					670		
Gln	Ala	Leu	Glu	Ala	Pro	Arg	Leu	Thr	Gln	Asn	Leu	Thr	Asp	Leu	Leu
		675					680						685		
Val	Asn	Val	Ser	Asp	Ser	Leu	Glu	Met	Gln	Cys	Leu	Val	Ala	Gly	Ala
	690						695					700			
His	Ala	Pro	Ser	Ile	Val	Trp	Tyr	Lys	Asp	Glu	Arg	Leu	Leu	Glu	Glu
	705					710					715				720
Lys	Ser	Gly	Val	Asp	Leu	Ala	Asp	Ser	Asn	Gln	Lys	Leu	Ser	Ile	Gln
			725					730						735	
Arg	Val	Arg	Glu	Glu	Asp	Ala	Gly	Arg	Tyr	Leu	Cys	Ser	Val	Cys	Asn
			740				745						750		
Ala	Lys	Gly	Cys	Val	Asn	Ser	Ser	Ala	Ser	Val	Ala	Val	Glu	Gly	Ser
		755					760					765			
Glu	Asp	Lys	Gly	Ser	Met	Glu	Ile	Val	Ile	Leu	Val	Gly	Thr	Gly	Val
	770					775						780			
Ile	Ala	Val	Phe	Phe	Trp	Val	Leu	Leu	Leu	Ile	Phe	Cys	Asn	Met	
	785					790					795			800	
Arg	Arg	Pro	Ala	His	Ala	Asp	Ile	Lys	Thr	Gly	Tyr	Leu	Ser	Ile	Ile
			805					810						815	
Met	Asp	Pro	Gly	Glu	Val	Pro	Leu	Glu	Glu	Gln	Cys	Glu	Tyr	Leu	Ser
			820				825							830	

-continued

Tyr Asp Ala Ser Gln Trp Glu Phe Pro Arg Glu Arg Leu His Leu Gly
 835 840 845
 Arg Val Leu Gly Tyr Gly Ala Phe Gly Lys Val Val Glu Ala Ser Ala
 850 855 860
 Phe Gly Ile His Lys Gly Ser Ser Cys Asp Thr Val Ala Val Lys Met
 865 870 875 880
 Leu Lys Glu Gly Ala Thr Ala Ser Glu His Arg Ala Leu Met Ser Glu
 885 890 895
 Leu Lys Ile Leu Ile His Ile Gly Asn His Leu Asn Val Val Asn Leu
 900 905 910
 Leu Gly Ala Cys Thr Lys Pro Gln Gly Pro Leu Met Val Ile Val Glu
 915 920 925
 Phe Cys Lys Tyr Gly Asn Leu Ser Asn Phe Leu Arg Ala Lys Arg Asp
 930 935 940
 Ala Phe Ser Pro Cys Ala Glu Lys Ser Pro Glu Gln Arg Gly Arg Phe
 945 950 955 960
 Arg Ala Met Val Glu Leu Ala Arg Leu Asp Arg Arg Arg Pro Gly Ser
 965 970 975
 Ser Asp Arg Val Leu Phe Ala Arg Phe Ser Lys Thr Glu Gly Gly Ala
 980 985 990
 Arg Arg Ala Ser Pro Asp Gln Glu Ala Glu Asp Leu Trp Leu Ser Pro
 995 1000 1005
 Leu Thr Met Glu Asp Leu Val Cys Tyr Ser Phe Gln Val Ala Arg Gly
 1010 1015 1020
 Met Glu Phe Leu Ala Ser Arg Lys Cys Ile His Arg Asp Leu Ala Ala
 1025 1030 1035 1040
 Arg Asn Ile Leu Leu Ser Glu Ser Asp Val Val Lys Ile Cys Asp Phe
 1045 1050 1055
 Gly Leu Ala Arg Asp Ile Tyr Lys Asp Pro Asp Tyr Val Arg Lys Gly
 1060 1065 1070
 Ser Ala Arg Leu Pro Leu Lys Trp Met Ala Pro Glu Ser Ile Phe Asp
 1075 1080 1085
 Lys Val Tyr Thr Thr Gln Ser Asp Val Trp Ser Phe Gly Val Leu Leu
 1090 1095 1100
 Trp Glu Ile Phe Ser Leu Gly Ala Ser Pro Tyr Pro Gly Val Gln Ile
 1105 1110 1115 1120
 Asn Glu Glu Phe Cys Gln Arg Leu Arg Asp Gly Thr Arg Met Arg Ala
 1125 1130 1135
 Pro Glu Leu Ala Thr Pro Ala Ile Arg Arg Ile Met Leu Asn Cys Trp
 1140 1145 1150
 Ser Gly Asp Pro Lys Ala Arg Pro Ala Phe Ser Glu Leu Val Glu Ile
 1155 1160 1165
 Leu Gly Asp Leu Leu Gln Gly Arg Gly Leu Gln Glu Glu Glu Val
 1170 1175 1180
 Cys Met Ala Pro Arg Ser Ser Gln Ser Ser Glu Glu Gly Ser Phe Ser
 1185 1190 1195 1200
 Gln Val Ser Thr Met Ala Leu His Ile Ala Gln Ala Asp Ala Glu Asp
 1205 1210 1215
 Ser Pro Pro Ser Leu Gln Arg His Ser Leu Ala Ala Arg Tyr Tyr Asn
 1220 1225 1230
 Trp Val Ser Phe Pro Gly Cys Leu Ala Arg Gly Ala Glu Thr Arg Gly
 1235 1240 1245

-continued

Ser Ser Arg Met Lys Thr Phe Glu Glu Phe Pro Met Thr Pro Thr Thr
 1250 1255 1260

Tyr Lys Gly Ser Val Asp Asn Gln Thr Asp Ser Gly Met Val Leu Ala
 265 1270 1275 1280

Ser Glu Glu Phe Glu Gln Ile Glu Ser Arg His Arg Gln Glu Ser Gly
 1285 1290 1295

Phe Ser Cys Lys Gly Pro Gly Gln Asn Val Ala Val Thr Arg Ala His
 1300 1305 1310

Pro Asp Ser Gln Gly Arg Arg Arg Arg Pro Glu Arg Gly Ala Arg Gly
 1315 1320 1325

Gly Gln Val Phe Tyr Asn Ser Glu Tyr Gly Glu Leu Ser Glu Pro Ser
 1330 1335 1340

Glu Glu Asp His Cys Ser Pro Ser Ala Arg Val Thr Phe Phe Thr Asp
 345 1350 1355 1360

Asn Ser Tyr

<210> SEQ ID NO 5
 <211> LENGTH: 1311
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens (FLT1)

<400> SEQUENCE: 5

Met Val Ser Tyr Trp Asp Thr Gly Val Leu Leu Cys Ala Leu Leu Ser
 1 5 10 15

Cys Leu Leu Leu Thr Gly Ser Ser Ser Gly Ser Lys Leu Lys Asp Pro
 20 25 30

Glu Leu Ser Leu Lys Gly Thr Gln His Ile Met Gln Ala Gly Gln Thr
 35 40 45

Leu His Leu Gln Cys Arg Gly Glu Ala Ala His Lys Trp Ser Leu Pro
 50 55 60

Glu Asn Asn Asn Asn Asn Asn Met Val Ser Lys Glu Ser Glu Arg Leu
 65 70 75 80

Ser Ile Thr Lys Ser Ala Cys Gly Arg Asn Gly Lys Gln Phe Cys Ser
 85 90 95

Thr Leu Thr Leu Asn Thr Ala Gln Ala Asn His Thr Gly Phe Tyr Ser
 100 105 110

Cys Lys Tyr Leu Ala Val Pro Thr Ser Lys Lys Lys Glu Thr Glu Ser
 115 120 125

Ala Ile Tyr Ile Phe Ile Ser Asp Thr Gly Arg Pro Phe Val Glu Met
 130 135 140

Tyr Ser Glu Ile Pro Glu Ile Ile His Met Thr Glu Gly Arg Glu Leu
 145 150 155 160

Val Ile Pro Cys Arg Val Thr Ser Pro Asn Ile Thr Val Thr Leu Lys
 165 170 175

Lys Phe Pro Leu Asp Thr Leu Ile Pro Asp Gly Lys Arg Ile Ile Trp
 180 185 190

Asp Ser Arg Lys Gly Phe Ile Ile Ser Asn Ala Thr Tyr Lys Glu Ile
 195 200 205

Gly Leu Leu Thr Cys Glu Ala Thr Val Asn Gly His Leu Tyr Lys Thr
 210 215 220

Asn Asn Tyr Leu Thr His Arg Gln Thr Asn Thr Ile Ile Asp Val Gln
 225 230 235 240

Ile Ser Thr Pro Arg Pro Val Lys Leu Leu Arg Gly His Thr Leu Val
 245 250 255

Leu	Asn	Cys	Thr	Ala	Thr	Thr	Pro	Leu	Asn	Thr	Arg	Val	Gln	Met	Thr	
			260					265					270			
Trp	Ser	Tyr	Pro	Asp	Asn	Asn	Asn	Glu	Lys	Asn	Lys	Arg	Ala	Ser	Val	
			275					280					285			
Arg	Arg	Arg	Ile	Asp	Gln	Ser	Asn	Ser	His	Ala	Asn	Ile	Phe	Tyr	Ser	
			290					295					300			
Val	Leu	Thr	Ile	Asp	Lys	Met	Gln	Asn	Lys	Asp	Lys	Gly	Leu	Tyr	Thr	
			305					310					315		320	
Cys	Arg	Val	Arg	Ser	Gly	Pro	Ser	Phe	Lys	Ser	Val	Asn	Thr	Ser	Val	
			325					330					335			
His	Ile	Tyr	Asp	Lys	Ala	Phe	Ile	Thr	Val	Lys	His	Arg	Lys	Gln	Gln	
			340					345					350			
Val	Leu	Glu	Thr	Val	Ala	Gly	Lys	Arg	Ser	Tyr	Arg	Leu	Ser	Met	Lys	
			355					360					365			
Val	Lys	Ala	Phe	Pro	Ser	Pro	Glu	Val	Val	Trp	Leu	Lys	Asp	Gly	Leu	
			370					375					380			
Pro	Ala	Thr	Glu	Lys	Ser	Ala	Arg	Tyr	Leu	Thr	Arg	Gly	Tyr	Ser	Leu	
			385					390					395		400	
Ile	Ile	Lys	Asp	Val	Thr	Glu	Glu	Asp	Ala	Gly	Asn	Tyr	Thr	Ile	Leu	
			405					410					415			
Leu	Ser	Ile	Lys	Gln	Ser	Asn	Val	Phe	Lys	Asn	Leu	Thr	Ala	Thr	Leu	
			420					425					430			
Ile	Val	Asn	Val	Lys	Pro	Gln	Ile	Tyr	Glu	Lys	Ala	Val	Ser	Ser	Phe	
			435					440					445			
Pro	Asp	Pro	Ala	Leu	Tyr	Pro	Leu	Gly	Ser	Arg	Gln	Ile	Leu	Thr	Cys	
			450					455					460			
Thr	Ala	Tyr	Gly	Ile	Pro	Gln	Pro	Asn	Thr	Ile	Lys	Trp	Phe	Trp	His	
			465					470					475		480	
Pro	Cys	Asn	His	Asn	His	Ser	Glu	Ala	Arg	Cys	Asp	Phe	Cys	Ser	Asn	
			485					490					495			
Asn	Glu	Glu	Ser	Phe	Ile	Leu	Asp	Asn	Asn	Asn	Asn	Asn	Asn	Asn	Ala	
			500					505					510			
Asp	Ser	Asn	Met	Gly	Asn	Arg	Ile	Glu	Ser	Ile	Thr	Gln	Arg	Met	Ala	
			515					520					525			
Ile	Ile	Glu	Gly	Lys	Asn	Lys	Met	Ala	Ser	Thr	Leu	Val	Val	Ala	Asp	
			530					535					540			
Ser	Arg	Ile	Ser	Gly	Ile	Tyr	Ile	Cys	Ile	Ala	Ser	Asn	Lys	Val	Gly	
			545					550					555		560	
Thr	Val	Gly	Arg	Asn	Ile	Ser	Phe	Tyr	Ile	Thr	Asp	Val	Pro	Asn	Gly	
			565					570					575			
Phe	His	Val	Asn	Leu	Glu	Lys	Met	Pro	Thr	Asn	Asn	Glu	Gly	Glu	Asp	
			580					585					590			
Leu	Lys	Leu	Ser	Cys	Thr	Val	Asn	Lys	Phe	Leu	Tyr	Arg	Asp	Val	Thr	
			595					600					605			
Trp	Ile	Leu	Leu	Asn	Asn	Asn	Asn	Asn	Asn	Asn	Asn	Asn	Asn	Asn	Asn	
			610					615					620			
Asn	Asn	Asn	Asn	Asn	Arg	Thr	Val	Asn	Asn	Arg	Thr					

-continued

675	680	685
Glu Ile Thr Ile Arg Asp 690	Gln Glu Ala Pro Tyr 695	Leu Leu Arg Asn Leu 700
Ser Asp His Thr Val Ala 705	Ile Ser Ser Ser Thr 710	Thr Thr Leu Asp Cys His 715 720
Ala Asn Gly Val Pro Glu 725	Pro Gln Ile Thr Trp 730	Phe Lys Asn Asn His 735
Lys Ile Gln Gln Glu Pro 740	Gly Ile Ile Leu Gly Pro 745	Gly Ser Ser Thr 750
Leu Phe Ile Glu Arg Val 755	Thr Glu Glu Asp Glu 760	Gly Val Tyr His Cys 765
Lys Ala Thr Asn Gln Lys 770	Gly Ser Val Glu Ser 775	Ser Ala Tyr Leu Thr 780
Val Gln Gly Thr Ser Asp 785	Lys Ser Asn Leu Glu 790	Leu Ile Thr Leu Thr 795 800
Cys Thr Cys Val Ala Ala 805	Thr Leu Phe Trp Leu 810	Leu Leu Thr Leu Leu 815
Ile Arg Lys Met Lys Arg 820	Ser Ser Asn Ser Glu 825	Ile Lys Thr Asp Tyr 830
Leu Ser Ile Ile Met Asp 835	Pro Asp Glu Val Pro 840	Leu Asp Glu Gln Cys 845
Glu Arg Leu Pro Tyr Asp 850	Ala Ser Lys Trp Glu 855	Phe Ala Arg Glu Arg 860
Leu Lys Leu Gly Lys Ser 865	Leu Gly Arg Gly Ala 870	Phe Gly Lys Val Val 875 880
Gln Ala Ser Ala Phe Gly 885	Ile Lys Lys Ser Pro 890	Thr Cys Arg Thr Val 895
Ala Val Lys Met Leu Lys 900	Glu Gly Ala Thr Ala 905	Ser Glu Tyr Lys Ala 910
Leu Met Thr Glu Leu Lys 915	Ile Leu Thr His Ile 920	Gly His His Leu Asn 925
Val Val Asn Leu Leu Gly 930	Ala Cys Thr Lys Gln 935	Gly Gly Pro Leu Met 940
Val Ile Val Glu Tyr Cys 945	Lys Tyr Gly Asn Leu 950	Ser Asn Tyr Leu Lys 955 960
Ser Lys Arg Asp Leu Phe 965	Phe Leu Asn Lys Asp 970	Ala Ala Leu His Met 975
Glu Pro Lys Lys Glu Lys 980	Met Glu Pro Gly Leu 985	Glu Gln Gly Lys Lys 990
Pro Arg Leu Asp Ser Val 995	Thr Ser Ser Glu Ser 1000	Phe Ala Ser Ser Gly 1005
Phe Gln Glu Asp Lys Ser 1010	Leu Ser Asp Val Glu 1015	Glu Glu Glu Asp Ser 1020
Asp Gly Phe Tyr Lys Glu 1025	Pro Ile Thr Met Glu 1030	Asp Leu Ile Ser Tyr 1035 1040
Ser Phe Gln Val Ala Arg 1045	Gly Met Glu Phe Leu 1050	Ser Ser Arg Lys Cys 1055
Ile His Arg Asp Leu Ala 1060	Ala Arg Asn Ile Leu 1065	Leu Ser Glu Asn Asn 1070
Val Val Lys Ile Cys Asp 1075	Phe Gly Leu Ala Arg 1080	Asp Ile Tyr Lys Asn 1085
Pro Asp Tyr Val Arg Lys 1090	Gly Asp Thr Arg Leu 1095	Pro Leu Lys Trp Met 1100

-continued

Ala Pro Glu Ser Ile Phe Asp Lys Ile Tyr Ser Thr Lys Ser Asp Val
 1105 1110 1115 1120

Trp Ser Tyr Gly Val Leu Leu Trp Glu Ile Phe Ser Leu Gly Gly Ser
 1125 1130 1135

Pro Tyr Pro Gly Val Gln Met Asp Glu Asp Phe Cys Ser Arg Leu Arg
 1140 1145 1150

Glu Gly Met Arg Met Arg Ala Pro Glu Tyr Ser Thr Pro Glu Ile Tyr
 1155 1160 1165

Gln Ile Met Leu Asp Cys Trp His Arg Asp Pro Lys Glu Arg Pro Arg
 1170 1175 1180

Phe Ala Glu Leu Val Glu Lys Leu Gly Asp Leu Leu Gln Ala Asn Val
 1185 1190 1195 1200

Gln Gln Asp Gly Lys Asp Tyr Ile Pro Ile Asn Ala Ile Leu Thr Gly
 1205 1210 1215

Asn Ser Gly Phe Thr Tyr Ser Thr Pro Ala Phe Ser Glu Asp Phe Phe
 1220 1225 1230

Lys Glu Ser Ile Ser Ala Pro Lys Phe Asn Ser Gly Ser Ser Asp Asp
 1235 1240 1245

Val Arg Tyr Val Asn Ala Phe Lys Phe Met Ser Leu Glu Arg Ile Lys
 1250 1255 1260

Thr Phe Glu Glu Leu Leu Pro Asn Ala Thr Ser Met Phe Asp Asp Tyr
 1265 1270 1275 1280

Gln Gly Asp Ser Ser Thr Leu Leu Ala Ser Pro Met Leu Lys Arg Phe
 1285 1290 1295

Thr Trp Thr Asp Ser Lys Pro Lys Ala Ser Leu Lys Ile Glu Val
 1300 1305 1310

<210> SEQ ID NO 6
 <211> LENGTH: 6
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: The amino acid at positions 1 and 2 each are
 selected independently from the group consisting of aspartic acid
 and glutamic acid.
 <220> FEATURE:
 <223> OTHER INFORMATION: The amino acid at position 4 is independently
 selected from the group consisting of methionine
 and valine.
 <220> FEATURE:
 <223> OTHER INFORMATION: The amino acid at position 5 is independently
 selected from the group consisting of proline,
 aspartic acid, and glutamic acid.
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: consensus
 sequence

<400> SEQUENCE: 6

Xaa Xaa Tyr Xaa Xaa Met
 1 5

<210> SEQ ID NO 7
 <211> LENGTH: 70
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence:
 oligonucleotide probe

<400> SEQUENCE: 7

acatgcatgc caccatgcag cggggcgccg cgctgtgcct gcgactgtgg ctctgcctgg 60

-continued

gactcctgga 70

<210> SEQ ID NO 8
 <211> LENGTH: 24
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence:
 oligonucleotide probe

<400> SEQUENCE: 8

acatgcatgc cccgccggtc atcc 24

<210> SEQ ID NO 9
 <211> LENGTH: 22
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence:
 oligonucleotide probe

<400> SEQUENCE: 9

cggaattccc catgacccca ac 22

<210> SEQ ID NO 10
 <211> LENGTH: 33
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence:
 oligonucleotide probe

<400> SEQUENCE: 10

ccatcgatgg atcctacctg aagccgcttt ctt 33

<210> SEQ ID NO 11
 <211> LENGTH: 34
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence:
 oligonucleotide probe

<400> SEQUENCE: 11

cccaagcttg gatccaagtg gctactccat gacc 34

<210> SEQ ID NO 12
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence:
 oligonucleotide probe

<400> SEQUENCE: 12

gttgctgtg atgtgcacca 20

<210> SEQ ID NO 13
 <211> LENGTH: 21
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence:
 oligonucleotide probe

<400> SEQUENCE: 13

ctggagtcga cttggcggac t 21

-continued

<210> SEQ ID NO 14
 <211> LENGTH: 60
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence:
 oligonucleotide probe
 <400> SEQUENCE: 14
 cgcggtatccc tagtgatggt gatggtgatg tctacctcg atcatgctgc cttatcctc 60

<210> SEQ ID NO 15
 <211> LENGTH: 21
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence:
 oligonucleotide probe
 <400> SEQUENCE: 15
 ctggagtcga cttggcggac t 21

<210> SEQ ID NO 16
 <211> LENGTH: 28
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence:
 oligonucleotide probe
 <400> SEQUENCE: 16
 cggtatccct ccatgctgcc cttatcct 28

<210> SEQ ID NO 17
 <211> LENGTH: 36
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence:
 oligonucleotide probe
 <400> SEQUENCE: 17
 ggcaagcttg aattcgccac catgcagcgg ggcgcc 36

<210> SEQ ID NO 18
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence:
 oligonucleotide probe
 <400> SEQUENCE: 18
 gttgcctgtg atgtgcacca 20

<210> SEQ ID NO 19
 <211> LENGTH: 21
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence:
 oligonucleotide probe
 <400> SEQUENCE: 19
 ctggagtcga cttggcggac t 21

-continued

<210> SEQ ID NO 20
 <211> LENGTH: 44
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence:
 oligonucleotide probe

<400> SEQUENCE: 20

cgcggatcca agcttactta ccttccatgc tgccttatc ctcg

44

<210> SEQ ID NO 21
 <211> LENGTH: 419
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 21

Met His Leu Leu Gly Phe Phe Ser Val Ala Cys Ser Leu Leu Ala Ala
 1 5 10 15
 Ala Leu Leu Pro Gly Pro Arg Glu Ala Pro Ala Ala Ala Ala Phe
 20 25 30
 Glu Ser Gly Leu Asp Leu Ser Asp Ala Glu Pro Asp Ala Gly Glu Ala
 35 40 45
 Thr Ala Tyr Ala Ser Lys Asp Leu Glu Glu Gln Leu Arg Ser Val Ser
 50 55 60
 Ser Val Asp Glu Leu Met Thr Val Leu Tyr Pro Glu Tyr Trp Lys Met
 65 70 75 80
 Tyr Lys Cys Gln Leu Arg Lys Gly Gly Trp Gln His Asn Arg Glu Gln
 85 90 95
 Ala Asn Leu Asn Ser Arg Thr Glu Glu Thr Ile Lys Phe Ala Ala Ala
 100 105 110
 His Tyr Asn Thr Glu Ile Leu Lys Ser Ile Asp Asn Glu Trp Arg Lys
 115 120 125
 Thr Gln Cys Met Pro Arg Glu Val Cys Ile Asp Val Gly Lys Glu Phe
 130 135 140
 Gly Val Ala Thr Asn Thr Phe Phe Lys Pro Pro Cys Val Ser Val Tyr
 145 150 155 160
 Arg Cys Gly Gly Cys Cys Asn Ser Glu Gly Leu Gln Cys Met Asn Thr
 165 170 175
 Ser Thr Ser Tyr Leu Ser Lys Thr Leu Phe Glu Ile Thr Val Pro Leu
 180 185 190
 Ser Gln Gly Pro Lys Pro Val Thr Ile Ser Phe Ala Asn His Thr Ser
 195 200 205
 Cys Arg Cys Met Ser Lys Leu Asp Val Tyr Arg Gln Val His Ser Ile
 210 215 220
 Ile Arg Arg Ser Leu Pro Ala Thr Leu Pro Gln Cys Gln Ala Ala Asn
 225 230 235 240
 Lys Thr Cys Pro Thr Asn Tyr Met Trp Asn Asn His Ile Cys Arg Cys
 245 250 255
 Leu Ala Gln Glu Asp Phe Met Phe Ser Ser Asp Ala Gly Asp Asp Ser
 260 265 270
 Thr Asp Gly Phe His Asp Ile Cys Gly Pro Asn Lys Glu Leu Asp Glu
 275 280 285
 Glu Thr Cys Gln Cys Val Cys Arg Ala Gly Leu Arg Pro Ala Ser Cys
 290 295 300
 Gly Pro His Lys Glu Leu Asp Arg Asn Ser Cys Gln Cys Val Cys Lys

-continued

305	310	315	320
Asn Lys Leu Phe Pro Ser Gln Cys Gly Ala Asn Arg Glu Phe Asp Glu	325	330	335
Asn Thr Cys Gln Cys Val Cys Lys Arg Thr Cys Pro Arg Asn Gln Pro	340	345	350
Leu Asn Pro Gly Lys Cys Ala Cys Glu Cys Thr Glu Ser Pro Gln Lys	355	360	365
Cys Leu Leu Lys Gly Lys Lys Phe His His Gln Thr Cys Ser Cys Tyr	370	375	380
Arg Arg Pro Cys Thr Asn Arg Gln Lys Ala Cys Glu Pro Gly Phe Ser	385	390	395
Tyr Ser Glu Glu Val Cys Arg Cys Val Pro Ser Tyr Trp Lys Arg Pro	405	410	415
Gln Met Ser			
<210> SEQ ID NO 22			
<211> LENGTH: 354			
<212> TYPE: PRT			
<213> ORGANISM: Homo sapiens			
<400> SEQUENCE: 22			
Met Tyr Arg Glu Trp Val Val Val Asn Val Phe Met Met Leu Tyr Val	1	5	10
Gln Leu Val Gln Gly Ser Ser Asn Glu His Gly Pro Val Lys Arg Ser	20	25	30
Ser Gln Ser Thr Leu Glu Arg Ser Glu Gln Gln Ile Arg Ala Ala Ser	35	40	45
Ser Leu Glu Glu Leu Leu Arg Ile Thr His Ser Glu Asp Trp Lys Leu	50	55	60
Trp Arg Cys Arg Leu Arg Leu Lys Ser Phe Thr Ser Met Asp Ser Arg	65	70	75
Ser Ala Ser His Arg Ser Thr Arg Phe Ala Ala Thr Phe Tyr Asp Ile	85	90	95
Glu Thr Leu Lys Val Ile Asp Glu Glu Trp Gln Arg Thr Gln Cys Ser	100	105	110
Pro Arg Glu Thr Cys Val Glu Val Ala Ser Glu Leu Gly Lys Ser Thr	115	120	125
Asn Thr Phe Phe Lys Pro Pro Cys Val Asn Val Phe Arg Cys Gly Gly	130	135	140
Cys Cys Asn Glu Glu Ser Leu Ile Cys Met Asn Thr Ser Thr Ser Tyr	145	150	155
Ile Ser Lys Gln Leu Phe Glu Ile Ser Val Pro Leu Thr Ser Val Pro	165	170	175
Glu Leu Val Pro Val Lys Val Ala Asn His Thr Gly Cys Lys Cys Leu	180	185	190
Pro Thr Ala Pro Arg His Pro Tyr Ser Ile Ile Arg Arg Ser Ile Gln	195	200	205
Ile Pro Glu Glu Asp Arg Cys Ser His Ser Lys Lys Leu Cys Pro Ile	210	215	220
Asp Met Leu Trp Asp Ser Asn Lys Cys Lys Cys Val Leu Gln Glu Glu	225	230	235
Asn Pro Leu Ala Gly Thr Glu Asp His Ser His Leu Gln Glu Pro Ala	245	250	255
Leu Cys Gly Pro His Met Met Phe Asp Glu Asp Arg Cys Glu Cys Val			

-continued

260				265				270							
Cys	Lys	Thr	Pro	Cys	Pro	Lys	Asp	Leu	Ile	Gln	His	Pro	Lys	Asn	Cys
275				280				285							
Ser	Cys	Phe	Glu	Cys	Lys	Glu	Ser	Leu	Glu	Thr	Cys	Cys	Gln	Lys	His
290				295				300							
Lys	Leu	Phe	His	Pro	Asp	Thr	Cys	Ser	Cys	Glu	Asp	Arg	Cys	Pro	Phe
305				310				315				320			
His	Thr	Arg	Pro	Cys	Ala	Ser	Gly	Lys	Thr	Ala	Cys	Ala	Lys	His	Cys
				325				330				335			
Arg	Phe	Pro	Lys	Glu	Lys	Arg	Ala	Ala	Gln	Gly	Pro	His	Ser	Arg	Lys
				340				345				350			
Asn				Pro											

What is claimed is:

1. A method of inhibiting Flt4 receptor tyrosine kinase (Flt4) function in a mammalian organism, comprising the step of administering to said mammalian organism a composition, said composition comprising an inhibitor of the binding of an Flt4 ligand protein to Flt4 expressed in cells of said organism, thereby inhibiting Flt4 function, wherein said inhibitor comprises a polypeptide selected from the group consisting of:

(a) a polypeptide comprising an antigen binding fragment of an anti Flt4 antibody; and

(b) a polypeptide comprising a soluble Flt4 fragment, wherein said fragment and said polypeptide are capable of binding to an Flt4 ligand.

2. A method according to claim 1 wherein said cells comprise endothelial cells.

3. A method according to claim 2 wherein said organism is human.

4. A method according to claim 3, wherein said composition further comprises a pharmaceutically acceptable diluent, adjuvant, or carrier medium.

5. A method according to claim 3 wherein said inhibitor comprises an anti Flt4 antibody or fragment thereof that binds to Flt4.

6. A method according to claim 5 wherein said inhibitor further comprises an anti-neoplastic agent conjugated to said antibody or antibody fragment.

7. A method according to claim 3 wherein said organism has a neoplastic disease characterized by expression of Flt4 tyrosine kinase (Flt4) in vascular endothelial cells,

wherein said composition comprises an inhibitor of the binding of an Flt4 ligand protein to Flt4 expressed in vascular endothelial cells of said organism, and

wherein said composition is administered in an amount effective to inhibit Flt4-mediated proliferation of said vascular endothelial cells, thereby inhibiting Flt4-mediated proliferation of said vascular endothelial cells.

8. A method according to claim 7 wherein said neoplastic disease is selected from the group consisting of carcinomas, squamous cell carcinomas, lymphomas, melanomas, and sarcomas.

9. A method according to claim 7 wherein said inhibitor comprises a polypeptide comprising an antigen-binding fragment of an anti-Flt4 antibody.

10. A method for antagonizing the function of Flt4 receptor tyrosine kinase (Flt4) in an organism that expresses Flt4,

comprising a step of administering to the organism a composition comprising a Flt4 antibody or Flt4 binding fragment thereof in a pharmaceutically acceptable carrier.

11. A method according to claim 10, wherein the organism is human.

12. A method according to claims 11 wherein said composition further comprises an anti-neoplastic agent conjugated to said antibody or antibody fragment.

13. A method according to claim 11, wherein the organism has a neoplastic disorder characterized by lymphatic vessels comprising lymphatic endothelia that express Flt4.

14. A method for antagonizing the function of Flt4 receptor tyrosine kinase (Flt4) in an organism that expresses Flt4, comprising a step of administering to the organism a composition comprising a soluble fragment of Flt4 in a pharmaceutically acceptable carrier, wherein the fragment binds to a Flt4 ligand.

15. A method according to claim 14, wherein the organism is human.

16. A method according to claim 15, wherein the organism has a neoplastic disorder characterized by lymphatic vessels comprising lymphatic endothelia that express Flt4.

17. A method of inhibiting neoplastic cell growth in a mammalian subject, comprising steps of:

(a) screening a mammalian subject to identify a neoplastic disorder characterized by cells expressing Flt4 receptor tyrosine kinase (Flt4); and

(b) administering a composition to a mammalian subject identified according to step (a) as having a neoplastic disorder characterized by cells expressing Flt4, said composition comprising an inhibitor of the binding of an Flt4 ligand protein to Flt4 expressed in cells of said subject, thereby inhibiting Flt4-mediated proliferation of said Flt4-expressing cells, wherein said inhibitor comprises a polypeptide selected from the group consisting of:

(i) a polypeptide comprising an antigen binding fragment of an anti Flt4 antibody; and

(ii) a polypeptide comprising a soluble Flt4 fragment, wherein said fragment and said polypeptide are capable of binding to an Flt4 ligand.

18. A method according to claim 17, wherein the mammalian subject is human.

19. A method according to claim 18, wherein step (a) comprises screening for a neoplastic disorder characterized by undesirable lymphatic vessels comprising lymphatic endothelia that express Flt4.

20. A method according to claim 18, wherein step (a) comprises screening for a neoplastic disorder characterized by neoplastic cells that express Flt4.

21. A method according to claim 20, wherein the neoplastic cells comprise lymphoma cells that express Flt4.

22. A method according to claim 18 wherein said composition further comprises a pharmaceutically acceptable diluent, adjuvant, or carrier medium.

23. A method according to claim 18 wherein said inhibitor comprises an anti-Flt4 antibody or fragment thereof.

24. A method according to claim 18, wherein the inhibitor comprises an extracellular domain fragment of human Flt4.

25. A method according to claim 18, wherein the screening step comprises:

(a) contacting tissue from the human subject with a composition comprising an Flt4 binding compound; and

(b) screening for a neoplastic disorder characterized by cells expressing Flt4 by detecting said Flt4 binding compound bound to said tissue.

26. A method according to claim 25 wherein said Flt4 binding compound is an antibody that specifically binds Flt4 or an antigen-binding fragment thereof.

27. A method according to claim 26, wherein said antibody or fragment further comprises a detectable label covalently bound thereto.

28. A method according to claim 18, wherein the screening step comprises:

(a) administering a composition to the human subject, said composition comprising an antibody that specifically binds Flt4 or an antigen-binding fragment thereof; and

(b) screening for a neoplastic disorder characterized by cells expressing Flt4 by detecting said antibody or said fragment bound to cells in said human subject, thereby detecting Flt4 expressed on the surface of cells in said human subject.

29. A method according to claim 28, wherein said antibody or antibody fragment further comprises a detectable label.

30. A method according to claim 17, wherein the screening step comprises:

(a) contacting tissue from the mammalian subject with a composition comprising an antibody or antibody fragment that specifically binds Flt4;

(b) detecting said antibody or antibody fragment bound to cells in said tissue; and

(c) screening for a neoplastic disorder from the quantity or distribution of said antibody bound to cells in said tissue.

31. A method according to claim 30, wherein in said screening step, the screening comprises measuring the quantity or distribution of said antibody bound to lymphatic vessels.

32. A method of treating a mammal having breast cancer characterized by endothelial cells that express Flt4 tyrosine kinase (Flt4), comprising a step of administering to said mammal a composition, said composition comprising an inhibitor of binding between Flt4 ligand protein and Flt4 expressed in cells of said organism, thereby inhibiting Flt4 function, wherein the inhibitor comprises a member selected from the group consisting of:

(a) an anti-Flt4 antibody or a polypeptide comprising an antigen binding fragment thereof;

(b) an anti-VEGF-C antibody or a polypeptide comprising an antigen binding fragment thereof;

(c) an anti-VEGF-D antibody or a polypeptide comprising an antigen binding fragment thereof; and

(d) a soluble polypeptide comprising a fragment of Flt4, wherein the polypeptide and the fragment are capable of binding to human VEGF-C (SEQ ID NO: 21).

33. A method according to claim 32, wherein the mammal is human.

34. A method according to claim 33, comprising a screening step preceding the administering step,

wherein the screening step comprises screening a human to identify breast cancer characterized by endothelial cells expressing Flt4; and

wherein the administering step comprises administering the composition to a human identified by the screening step as having breast cancer characterized by endothelial cells expressing Flt4.

35. A method according to any one of claims 33-34 wherein the inhibitor comprises a member selected from the group consisting of:

(a) an anti-Flt4 antibody or a polypeptide comprising an antigen binding fragment thereof; and

(b) a soluble polypeptide comprising a fragment of Flt4, wherein the polypeptide and the fragment are capable of binding to human VEGF-C (SEQ ID NO: 21).

36. A method according to claim 32, wherein the inhibitor further comprises an anti-neoplastic agent conjugated to the antibody or polypeptide.

37. A method according to claim 32, wherein the composition further comprises a pharmaceutically acceptable diluent, adjuvant, or carrier.

38. A method of inhibiting proliferation of cells in a mammalian organism having a disease characterized by expression of Flt4 tyrosine kinase (Flt4) in cells, comprising the step of administering to said mammalian organism a composition, said composition comprising an inhibitor of the binding of an Flt4 ligand protein to Flt4 expressed in cells of said organism, thereby inhibiting Flt4-mediated proliferation of the cells, wherein said inhibitor comprises a polypeptide selected from the group consisting of:

(a) a polypeptide comprising an antigen binding fragment of an anti Flt4 antibody; and

(b) a polypeptide comprising a soluble Flt4 fragment, wherein said fragment and said polypeptide are capable of binding to an Flt4 ligand.

39. A method according to claim 38 wherein said cells comprise endothelial cells.

40. A method according to claim 39, wherein said cells comprise lymphatic endothelial cells, and said composition inhibits lymphatic vascularization.

41. A method according to claim 39 wherein said organism is human.

42. A method according to claim 38 wherein said composition further comprises a pharmaceutically acceptable diluent, adjuvant, or carrier medium.

43. A method according to claim 41 wherein said inhibitor comprises an anti Flt4 antibody or fragment thereof that binds to Flt4.

44. A method according to claim 43 wherein said inhibitor further comprises an anti-neoplastic agent conjugated to said antibody or antibody fragment.

45. A method according to claim 41, wherein the inhibitor comprises an anti-Flt4 antibody.

46. A method according to claim 41 wherein the inhibitor comprises a polypeptide comprising an antigen-binding fragment of an anti-Flt4 antibody.

47. A method according to claim 41 wherein said neoplastic disease is selected from the group consisting of carcinomas, squamous cell carcinomas, lymphomas, melanomas, and sarcomas.

101

48. A method according to claim 42 wherein the inhibitor comprises a polypeptide comprising an antigen-binding fragment of an anti-Flt4 antibody.

49. A method according to claim 41, wherein the disease is a cancer characterized by metastatic lymph nodes.

50. A method of inhibiting genesis of lymphatic vessels in a mammalian organism having a disease characterized by expression of Flt4 tyrosine kinase (Flt4) in lymphatic vessels, comprising the step of administering to said mam-
malian organism a composition, said composition compris-
ing an inhibitor of the binding of an Flt4 ligand protein to
Flt4 expressed in cells of said organism, thereby inhibiting
Flt4-mediated proliferation of the cells, wherein said inhibi-
tor comprises a member selected from the group consisting
of:

- (a) an anti-Flt4 antibody;
- (b) a polypeptide comprising an antigen binding fragment
of an anti Flt4 antibody; and
- (c) a polypeptide comprising a soluble Flt4 fragment,
wherein said fragment and said polypeptide are capable
of binding to an Flt4 ligand.

51. A method according to claim 50 wherein said organ-
ism is human.

102

52. A method according to claim 51, wherein the human has a cancer characterized by lymph node metastases.

53. A method of treating a human having breast cancer characterized by endothelial cells that express Flt4 tyrosine kinase (Flt4), comprising a step of administering to said human a composition, said composition comprising an inhibitor of binding between Flt4 ligand protein and Flt4 expressed in cells of said human, thereby inhibiting Flt4 function,

wherein the inhibitor comprises a polypeptide comprising an Flt4 binding fragment of human prepro-VEGF-C (SEQ ID NO: 21) or human prepro-VEGF-D (SEQ ID NO: 22) conjugated to an antineoplastic agent.

54. A method according to claim 53, comprising a screen-
ing step preceding the administering step,

wherein the screening step comprises screening a human to identify breast cancer characterized by endothelial cells expressing Flt4; and

wherein the administering step comprises administering the composition to a human identified by the screening step as having breast cancer characterized by endothelial cells expressing Flt4.

* * * * *

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 6,824,777 B1
APPLICATION NO. : 09/169079
DATED : November 30, 2004
INVENTOR(S) : Kari Alitalo et al.

Page 1 of 1

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

On the cover page, section (63), please replace the entire section with --This application is a continuation-in-part of United States Patent Application Serial No. 08,901,710, filed July 28, 1997, now U.S. Patent No. 6,107,046; which is a continuation-in-part of both United States Patent Application Serial No. 08/340,011, filed November 14, 1994, now U.S. Patent No. 5,776,755; and United States Patent Application Serial No. 08/257,754, filed June 9, 1994, now abandoned; the latter two of which in turn are continuations-in-part on United States Patent Application Serial No. 07/959,951, filed October 9, 1992, now abandoned.--

At column 97, claim 7, lines 48-58, please delete the entire claim 7.

At column 97, claim 8, line 59, please delete "claim 7" and insert --claim 5--

At column 97, claim 9, line 63, please delete "claim 7" and insert --claim 8--

At column 98, claim 12, line 25, please delete "claims" and insert --claim--

At column 99, claim 24, line 9, please delete "Fit4." and insert --FIIt4.--

Signed and Sealed this

Twenty-fourth Day of July, 2007

A handwritten signature in black ink, appearing to read "Jon W. Dudas". The signature is stylized with a large, looping initial "J" and a distinct "D".

JON W. DUDAS
Director of the United States Patent and Trademark Office